

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

UPPER BOUNDS FOR COURANT-SHARP NEUMANN AND ROBIN EIGENVALUES

Katie Gittins & Corentin Léna

Tome 148
Fascicule 1

2020

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

pages 99-132

Le *Bulletin de la Société Mathématique de France* est un périodique trimestriel
de la Société Mathématique de France.

Fascicule 1, tome 148, mars 2020

Comité de rédaction

| | |
|---------------------|------------------|
| Christine BACHOC | Laurent MANIVEL |
| Yann BUGEAUD | Julien MARCHÉ |
| Jean-François DAT | Kieran O'GRADY |
| Clothilde FERMANIAN | Emmanuel RUSS |
| Pascal HUBERT | Christophe SABOT |

Marc HERZLICH (Dir.)

Diffusion

| | |
|--|--|
| Maison de la SMF | AMS |
| Case 916 - Luminy | P.O. Box 6248 |
| 13288 Marseille Cedex 9 | Providence RI 02940 |
| France | USA |
| commandes@smf.emath.fr | www.ams.org |

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Bulletin de la SMF

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 1 44 27 67 99 • Fax : (33) 1 40 46 90 96

bulletin@smf.emath.fr • smf.emath.fr

© Société Mathématique de France 2020

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Stéphane SEURET

UPPER BOUNDS FOR COURANT-SHARP NEUMANN AND ROBIN EIGENVALUES

BY KATIE GITTINS & CORENTIN LÉNA

ABSTRACT. — We consider the eigenvalues of the Laplacian, with a Neumann or Robin boundary condition, on an open, bounded, connected set in \mathbb{R}^n with a C^2 boundary. We obtain upper bounds for the eigenvalues that have a corresponding eigenfunction that achieves equality in Courant's Nodal Domain theorem. In the case where the set is also assumed to be convex, we obtain explicit upper bounds in terms of some of the geometric quantities of the set.

RÉSUMÉ (*Majoration des valeurs propres Courant strictes de Neumann et Robin*). — Nous considérons les valeurs propres du laplacien sur un ouvert borné connexe de \mathbb{R}^n à bord C^2 , avec condition au bord de Neumann ou de Robin. Nous majorons celles qui ont une fonction propre dont le nombre de domaines nodaux atteint la borne de Courant (dites *Courant strictes*). Lorsque l'ouvert est convexe, nous présentons une majoration explicite en fonction de grandeurs géométriques.

1. Introduction

1.1. Statement of the problem. — Let Ω be an open, bounded, connected set in \mathbb{R}^n , $n \geq 2$, with a Lipschitz boundary $\partial\Omega$. Consider the Neumann Laplacian

Texte reçu le 6 novembre 2018, modifié le 8 mai 2019, accepté le 20 mai 2019.

KATIE GITTINS, Université de Neuchâtel, Institut de Mathématiques, Rue Emile-Argand 11, CH-2000 Neuchâtel, Suisse • *E-mail* : katie.gittins@unine.ch

CORENTIN LÉNA, Matematiska Institutionen, Stockholms Universitet, Kräftriket Hus 6, SE-106 91 Stockholm, Suède • *E-mail* : corentin@math.su.se

Mathematical subject classification (2010). — 35P15, 49R05, 35P05.

Key words and phrases. — Courant's nodal domain theorem, Neumann eigenvalues, Robin eigenvalues, Euclidean domains.

acting on $L^2(\Omega)$ and note that it has discrete spectrum since Ω is bounded. The Neumann eigenvalues of Ω can hence be written in a non-decreasing sequence, counted with multiplicity,

$$0 = \mu_1(\Omega) < \mu_2(\Omega) \leq \cdots \leq \mu_k(\Omega) \leq \dots,$$

where the only accumulation point is $+\infty$.

By Courant's Nodal Domain theorem, any eigenfunction corresponding to $\mu_k(\Omega)$ has at most k nodal domains. If u_k is an eigenfunction corresponding to $\mu_k(\Omega)$ with k nodal domains, then we call it a Courant-sharp eigenfunction. In this case, we also call $\mu_k(\Omega)$ a Courant-sharp eigenvalue of Ω .

The Courant-sharp property was first considered by Pleijel [15] in 1956 for the Dirichlet Laplacian. In particular, Pleijel proved that there are only finitely many Courant-sharp Dirichlet eigenvalues of a bounded, planar domain with a sufficiently regular boundary. See [6, 14] for generalisations of Pleijel's theorem to higher dimensions and other geometric settings. Following from Pleijel's result, for a given domain, natural questions are, how many such eigenvalues are there and how large are they?

The recent articles [5, 2] consider these questions and give upper bounds for the largest Courant-sharp Dirichlet eigenvalue and the number of such eigenvalues in terms of some of the geometric quantities of the underlying domain. In Theorem 1.3 of [5], the authors bound this number using the area, perimeter, maximal curvature and minimal cut-distance to the boundary, for a set in \mathbb{R}^2 which is sufficiently regular but not necessarily convex (the cut-distance will be defined in Section 3). In [2], such geometric upper bounds are obtained for an open set in \mathbb{R}^n with finite Lebesgue measure. In the case where the domain is convex, the upper bound given in Example 1 of [2] could be expressed in terms of the isoperimetric ratio of the domain. From this, one can deduce that if the domain has a large number of Courant-sharp Dirichlet eigenvalues then its isoperimetric ratio is also large.

It was shown recently in [10] that if Ω is an open, bounded, connected set in \mathbb{R}^n with a $C^{1,1}$ boundary, then the Neumann Laplacian acting on $L^2(\Omega)$ has finitely many Courant-sharp eigenvalues (we refer to [10] for a description of prior results). As mentioned in [5], the aforementioned questions are also interesting for the Courant-sharp eigenvalues of the Neumann Laplacian.

1.2. Goal of the article. — The aim of the present article is to obtain upper bounds for the Courant-sharp Neumann eigenvalues in the case where $\Omega \subset \mathbb{R}^n$ is open, bounded, and connected with a C^2 boundary. In the case where Ω is also convex, we obtain explicit upper bounds for the Courant-sharp Neumann eigenvalues of Ω , and for the number of such eigenvalues, in terms of some of the geometric quantities of Ω . These results correspond to some of those mentioned above for the Dirichlet case, with some additional hypotheses due to the difficulties in handling the Neumann boundary condition.

We follow the same strategy that was used in [10]. This involves distinguishing between the nodal domains of a Courant-sharp eigenfunction u for which the majority of the L^2 norm of u comes from the interior (bulk domains) and those for which the majority of the L^2 norm of u comes from near the boundary (boundary domains), and then obtaining upper bounds for the number of each type of nodal domain. In the first case, the argument used by Pleijel [15], which rests upon the Faber-Krahn inequality, can be used as the eigenfunction in a bulk domain almost satisfies a Dirichlet boundary condition. For the boundary domains, it is not possible to employ the same argument as Pleijel as these nodal domains have mixed Dirichlet-Neumann boundary conditions so the Faber-Krahn inequality cannot be employed. The strategy of [10] to deal with the boundary domains is to locally straighten the boundary of the domain Ω and then to reflect the nodal domain in order to obtain a new domain that almost satisfies a Dirichlet boundary condition. One then has to compare the L^2 norm of the gradient of an eigenfunction corresponding to a Courant-sharp eigenvalue on the boundary domain to the L^2 norm of the gradient of the reflected eigenfunction on the reflected domain. See Section 5.

We restrict our attention to Euclidean domains with a C^2 boundary. We can then make use of tubular coordinates in order to set up and describe the reflection procedure explicitly. This allows us to keep explicit control of the constants appearing in the aforementioned estimates in order to obtain estimates for the Courant-sharp Neumann eigenvalues.

In Proposition 7.1, we obtain an upper bound for the Courant-sharp Neumann eigenvalues of Ω in terms of some of its geometric quantities. More specifically, it depends on $|\Omega|$ the area of Ω , $\rho(\Omega)$ the square-root of the isoperimetric ratio, $t_+(\Omega)$ the smallest radius of curvature of the boundary, and the cut distance to the boundary (see Section 3 for precise definitions of the latter quantities).

A simpler presentation of this upper bound is possible in the case where Ω is convex, since one of the additional conditions in the general case is no longer required (see Section 8). In addition, we obtain an upper bound for the number of such eigenvalues by using the upper bound for the Neumann counting function which is proved in Appendix A. In particular, we have the following proposition.

PROPOSITION 1.1. — *Let Ω be an open, bounded, convex set in \mathbb{R}^2 with a C^2 boundary. There exist constants $C > 0$ and $C' > 0$, that do not depend on Ω , such that for any Courant-sharp eigenvalue $\mu_k(\Omega)$,*

$$(1) \quad \mu_k(\Omega) \leq C \left(\frac{|\Omega|}{t_+(\Omega)^4} + \frac{\rho(\Omega)^8}{|\Omega|} \right)$$

and

$$(2) \quad k \leq C' \left(\frac{|\Omega|^2}{t_+(\Omega)^4} + \rho(\Omega)^8 \right).$$

We note that the left-hand side and right-hand side of Inequality (1) have the same homogeneity with respect to scaling, and that Inequality (2) is scaling invariant. In addition, in Section 8, we obtain an upper bound for the Courant-sharp Neumann eigenvalues which also depends upon the diameter of Ω .

By Proposition 1.1, we then observe that if Ω is a sufficiently regular convex set with a large number of Courant-sharp eigenvalues, it has a large isoperimetric ratio or a large curvature at some point of its boundary (or both). If we additionally assume that $\mu_k(\Omega)$ is large compared with $|\Omega|t_+(\Omega)^{-4}$, we can conclude that the isoperimetric ratio is large. We note that a large isoperimetric ratio is enough to generate a large number of Courant-sharp Neumann eigenvalues. Indeed, this is the case for a rectangle $(0, 1) \times (0, L)$ with large L . By contrast, to the best of the authors' knowledge, it is not known whether a boundary point with large curvature alone can generate many Courant-sharp eigenvalues. It could be interesting to investigate this further.

By $-\Delta_\Omega^\beta$, we denote the Laplacian on Ω with the following Robin boundary condition

$$\frac{\partial u}{\partial \nu} + \beta u = 0 \text{ on } \partial\Omega,$$

where $\frac{\partial u}{\partial \nu}$ is the exterior normal derivative and $\beta : \partial\Omega \rightarrow \mathbb{R}$ is a non-negative, Lipschitz continuous function. We denote the corresponding eigenvalues by $(\mu_k(\Omega, \beta))_{k \geq 1}$. It was shown in [10] that there are finitely many Courant-sharp eigenvalues of $-\Delta_\Omega^\beta$. By monotonicity of the Robin eigenvalues with respect to β , we obtain the same results for the Courant-sharp Robin eigenvalues (see Subsection 2.2).

In addition, we obtain analogous results to those mentioned above for any dimension $n \geq 3$, namely Propositions 9.2, 9.3 and 9.4 in Section 9.

1.3. Organisation of the article. — In Section 2, we show that in order to obtain upper bounds for the largest Courant-sharp eigenvalue μ , it is sufficient to obtain upper bounds for the number of nodal domains and the remainder of the Dirichlet counting function. Estimates for the latter are obtained in Section 6. To deal with the former, we first consider the 2-dimensional case and set up tubular coordinates in Section 3. Following [10], we then define cut-off functions in Section 4 that allow us to distinguish between bulk and boundary domains. In Subsection 5.1 we perform the straightening of the boundary procedure and obtain the desired estimates. We then use these estimates in Subsection 5.2 to obtain an explicit upper bound for the number of nodal domains. In Subsection 5.3 and Subsection 5.4, by taking the geometry of the domain into account, we improve the estimates from Subsection 5.1 in special cases. We then combine all of the preceding results in Section 7 to obtain an upper bound for the largest Courant-sharp eigenvalue. In Section 8, we obtain

explicit upper bounds for the largest Courant-sharp eigenvalue and the number of Courant-sharp eigenvalues of an open and convex planar domain with a C^2 boundary that involve some of its geometric quantities. In particular, we prove Proposition 1.1. In Section 9, we obtain analogous results in arbitrary dimension $n \geq 3$. In Appendix A, we prove an upper bound for the Neumann counting function of a convex set, which is used in the two preceding sections to control the number of Courant-sharp eigenvalues.

2. Preliminaries

2.1. Strategy for Courant-sharp Neumann eigenvalues. — For $\mu > 0$, we define the Neumann counting function as follows:

$$N_{\Omega}^N(\mu) := \#\{k \in \mathbb{N}^* : \mu_k(\Omega) < \mu\}.$$

Let $(\lambda_k(\Omega))_{k \geq 1}$ denote the Dirichlet eigenvalues of the Laplacian on Ω . By the min-max characterisations of the Neumann and Dirichlet eigenvalues, we have, for $k \in \mathbb{N}^*$, that

$$(3) \quad \mu_k(\Omega) \leq \lambda_k(\Omega).$$

For $\mu > 0$, we define the Dirichlet counting function:

$$N_{\Omega}^D(\mu) := \#\{k \in \mathbb{N}^* : \lambda_k(\Omega) < \mu\},$$

and the corresponding *remainder* $R_{\Omega}^D(\mu)$ such that

$$(4) \quad N_{\Omega}^D(\mu) = \frac{\omega_n |\Omega|}{(2\pi)^n} \mu^{n/2} - R_{\Omega}^D(\mu),$$

where ω_n denotes the Lebesgue measure of a ball of radius 1 in \mathbb{R}^n , and the first term in the right-hand side of Equation (4) corresponds to Weyl's law. By (3), we have

$$N_{\Omega}^N(\mu) \geq N_{\Omega}^D(\mu),$$

and therefore

$$N_{\Omega}^N(\mu) \geq \frac{\omega_n |\Omega|}{(2\pi)^n} \mu^{n/2} - R_{\Omega}^D(\mu).$$

Consider an eigenpair (μ, u) for the Neumann Laplacian, and denote by $\nu(u)$ the number of its nodal domains. If u is a Courant-sharp eigenfunction associated with $\mu > 0$, $\mu = \mu_k(\Omega)$ with $\nu(u) = k$. On the other hand, Courant's Nodal Domain theorem implies that $\mu_{k-1}(\Omega) < \mu_k(\Omega)$, so that $N_{\Omega}^N(\mu) = k - 1$. We therefore have

$$(5) \quad N_{\Omega}^N(\mu) - \nu(u) < 0.$$

Hence, in order to obtain upper bounds for μ , we require upper bounds for $\nu(u)$ and $R_{\Omega}^D(\mu)$. These will be obtained in Sections 5, 6 respectively.

To obtain an upper bound for $\nu(u)$, we follow the strategy of [10] in which an important step is to straighten the boundary locally. By restricting our attention to domains with a C^2 boundary, we can make use of tubular coordinates to straighten the boundary which allow us to obtain the desired explicit estimates in Subsection 5.1.

2.2. Application to Courant-sharp Robin eigenvalues. — Analogous arguments to the above hold for the Robin eigenvalues $(\mu_k(\Omega, \beta))_{k \geq 1}$. For $\mu > 0$, we define the Robin counting function:

$$N_{\Omega}^{\beta}(\mu) := \#\{k \in \mathbb{N}^* : \mu_k(\Omega, \beta) < \mu\}.$$

For $k \in \mathbb{N}^*$, the Robin eigenvalues satisfy the following monotonicity property with respect to $\beta \geq 0$:

$$\mu_k(\Omega) \leq \mu_k(\Omega, \beta) \leq \lambda_k(\Omega),$$

where $\beta \equiv 0$ gives rise to the Neumann eigenvalues and $\beta \rightarrow +\infty$ corresponds to the Dirichlet eigenvalues. We then have

$$N_{\Omega}^{\beta}(\mu) \geq N_{\Omega}^D(\mu),$$

and hence

$$N_{\Omega}^{\beta}(\mu) \geq \frac{\omega_n |\Omega|}{(2\pi)^n} \mu^{n/2} - R_{\Omega}^D(\mu).$$

(See, for example, [10, Section 4]).

3. Tubular coordinates in 2D

We assume that $n = 2$, that is $\Omega \subset \mathbb{R}^2$. For any $r > 0$, we define the inner tubular neighbourhood of $\partial\Omega$ with radius r :

$$\partial\Omega_r^+ := \{x \in \Omega; \text{dist}(x, \partial\Omega) < r\},$$

and its volume

$$\tau(r) := |\partial\Omega_r^+|.$$

We now assume that Ω is simply connected (we consider multiply connected domains in Subsection 5.3). Let $\gamma : [0, L] \rightarrow \mathbb{R}^2$ be a closed, simple and C^2 curve parametrised by arc length, such that $\partial\Omega = \gamma([0, L])$. In particular, L is the total length of $\partial\Omega$. For each $s \in [0, L]$, we write $\mathbf{t}(s) := \gamma'(s)$, the unit tangent vector at $\gamma(s)$, and denote by $\mathbf{n}(s)$ the unit vector such that $(\mathbf{t}(s), \mathbf{n}(s))$ is a direct orthonormal basis. Up to reversing the orientation of the curve γ , we can assume that $\mathbf{n}(s)$ points towards the interior of Ω . The signed curvature of γ at the point $\gamma(s)$ (relative to our choice of orientation), denoted by $\kappa(s)$, is then defined by

$$\mathbf{t}'(s) = \kappa(s) \mathbf{n}(s).$$

We define the mapping

$$(6) \quad \begin{aligned} F: [0, L] \times \mathbb{R} &\rightarrow \mathbb{R}^2 \\ q := (s, t) &\mapsto x := \gamma(s) + t \mathbf{n}(s). \end{aligned}$$

The function F is of class C^1 , and its differential at $q := (s, t)$, expressed from the base $((1, 0), (0, 1))$ to the base $(\mathbf{t}(s), \mathbf{n}(s))$, is

$$(7) \quad J(q) = \begin{pmatrix} 1 - t \kappa(s) & 0 \\ 0 & 1 \end{pmatrix}.$$

Following Subsection 3.1 of [5], we define

$$(8) \quad t_+(\Omega) := \left(\sup_{s \in [0, L]} |\kappa(s)| \right)^{-1},$$

and, for $s \in [0, L]$, the (internal) *cut-distance* to $\partial\Omega$ at $\gamma(s)$:

$$(9) \quad \delta_+(s) := \sup\{\delta > 0; \text{dist}(F(s, t), \partial\Omega) = t \text{ for all } t \in [0, \delta]\}.$$

We set

$$(10) \quad \underline{\delta}_+(\Omega) := \inf_{s \in [0, L]} \delta_+(s)$$

and

$$(11) \quad \delta_0(\Omega) := \min\{t_+(\Omega), \underline{\delta}_+(\Omega)\}.$$

By construction, F is a diffeomorphism of class C^1 from $(0, L) \times (0, \delta_0(\Omega))$ to $\partial\Omega_{\delta_0(\Omega)}^+ \setminus \ell_0$, where ℓ_0 is the segment $F(\{0\} \times (0, \delta_0(\Omega)))$.

Given $f: \mathbb{R} \rightarrow \mathbb{R}$ continuous and piecewise C^1 and $\delta \leq \delta_0(\Omega)$, we define the function $\varphi: \partial\Omega_\delta^+ \setminus \ell_0 \rightarrow \mathbb{R}$ by $\varphi(x) = f(t)$, where $x = F(s, t)$. By the definition of $\delta_0(\Omega)$, this can alternatively be written as $\varphi(x) = f(\text{dist}(x, \partial\Omega))$.

PROPOSITION 3.1. — *The function φ is continuous, and is of class C^1 except on the regular arcs*

$$\Gamma_i = \{x \in \Omega; \text{dist}(x, \partial\Omega) = t_i\},$$

where $\{t_i; 1 \leq i \leq N\}$ are the points of discontinuity of f' in $(0, \delta_0(\Omega)]$. Furthermore,

$$(12) \quad \nabla\varphi(x) = f'(t)\mathbf{n}(s),$$

with $x = F(s, t)$, if $x \notin \bigcup_{i=1}^N \Gamma_i$.

Proof. — This follows from the chain rule, using the expression for the Jacobian matrix given in Equation (7). □

4. Cut-off functions

The purpose of this section is to define the cut-off functions $\varphi_0^\delta, \varphi_1^\delta$ in order to characterise the nodal domains as bulk domains or boundary domains, as in Subsection 2.2 of [10]. The key point here is to obtain explicit estimates.

Step 1: We construct two functions $\chi_0, \chi_1 : \mathbb{R} \rightarrow \mathbb{R}$ which are continuous, piecewise C^1 , and satisfy

- i. for $t \leq \frac{1}{4}$, $\chi_0(t) = 0$ and $\chi_1(t) = 1$;
- ii. for $t \in [\frac{1}{4}, \frac{3}{4}]$, $0 \leq \chi_0(t), \chi_1(t) \leq 1$;
- iii. for $t \geq \frac{3}{4}$, $\chi_0(t) = 1$ and $\chi_1(t) = 0$;
- iv. $\chi_0^2 + \chi_1^2 = 1$;

We write $B := \max(\|\chi_0'\|_{L^\infty}, \|\chi_1'\|_{L^\infty})$.

We first construct $\psi : \mathbb{R} \rightarrow \mathbb{R}$, continuous, piecewise C^1 and non-decreasing, such that $\psi(0) = 0$ and $\psi(1) = 1$ and satisfying $0 \leq \psi \leq 1$. We impose the additional condition

$$\psi(t)^2 + \psi(1-t)^2 = 1$$

for all $t \in \mathbb{R}$. One possible choice for ψ is given at the end of this section.

We then set

$$\begin{aligned}\chi_0(t) &:= \psi\left(2\left(t - \frac{1}{4}\right)\right); \\ \chi_1(t) &:= \psi\left(2\left(\frac{3}{4} - t\right)\right).\end{aligned}$$

The functions χ_0 and χ_1 have the desired properties with $B = 2\|\psi'\|_{L^\infty}$.

Step 2: For each $\delta \in (0, \delta_0(\Omega)]$, we construct two functions $\varphi_0^\delta, \varphi_1^\delta : \Omega \rightarrow \mathbb{R}$ which are continuous and piecewise C^1 (the gradient is continuous except for finite jumps on regular arcs), and satisfy, for some positive constant C independent of δ ,

- i. for $\text{dist}(x, \partial\Omega) \leq \frac{\delta}{4}$, $\varphi_0^\delta(x) = 0$ and $\varphi_1^\delta(x) = 1$;
- ii. for $\text{dist}(x, \partial\Omega) \in [\frac{\delta}{4}, \frac{3\delta}{4}]$, $0 \leq \varphi_0^\delta(x), \varphi_1^\delta(x) \leq 1$;
- iii. for $\text{dist}(x, \partial\Omega) \geq \frac{3\delta}{4}$, $\varphi_0^\delta(x) = 1$ and $\varphi_1^\delta(x) = 0$;
- iv. $(\varphi_0^\delta)^2 + (\varphi_1^\delta)^2 = 1$;
- v. $\|\nabla\varphi_0^\delta\|_{L^\infty}, \|\nabla\varphi_1^\delta\|_{L^\infty} \leq C\delta^{-1}$.

The functions χ_0 and χ_1 of Step 1 being given, we define, for $x \in \partial\Omega_\delta^+$,

$$\begin{aligned}\varphi_0^\delta(x) &:= \chi_0\left(\frac{t}{\delta}\right); \\ \varphi_1^\delta(x) &:= \chi_1\left(\frac{t}{\delta}\right).\end{aligned}$$

We make the obvious extensions of φ_0^δ and φ_1^δ so that both functions are continuous. From the enumerated properties of χ_0 and χ_1 and Equation (12), it follows that φ_0^δ and φ_1^δ have the desired properties with $C = B$.

Explicit constants: In order to obtain explicit estimates in what follows, it is necessary to specify ψ . One possible choice for ψ is given by

$$\psi(t)^2 := \frac{1}{\int_0^1 s(1-s) ds} \int_0^t s(1-s) ds,$$

for $t \in [0, 1]$, extended by 0 for $t \leq 0$ and by 1 for $t \geq 1$. More explicitly,

$$\psi(t) = \sqrt{3t^2 - 2t^3}$$

for $t \in [0, 1]$. In that case $\|\psi'\|_{L^\infty} = \sqrt{3}$. The functions χ_0 and χ_1 that we construct satisfy the listed properties with $B = 2\sqrt{3}$, and the functions φ_0^δ and φ_1^δ with $C = 2\sqrt{3}$.

5. Estimates of the nodal count

We wish to count the number of each type of nodal domain. For the bulk domains, one considers a Pleijel-type argument via the Faber-Krahn inequality. For the boundary domains, one reflects them in the boundary (after straightening it) and then applies the Faber-Krahn inequality to the reflected domains. See Section 2 of [10] for the full details and Subsection 5.2 below.

5.1. Straightening of the boundary. — We start by giving explicit versions of some estimates in Subsections 2.4 and 2.5 of Reference [10], in order to control some quantities of interest when we straighten the boundary using tubular coordinates.

LEMMA 5.1. — *Let V be an open set $V \subset (0, L) \times (0, 3\delta_0(\Omega)/4)$ and let $U = F(V)$. There exist constants $0 < m_- \leq m_+$ such that, for any measurable and non-negative function $g : V \rightarrow \mathbb{R}$,*

$$m_- \int_V g dq \leq \int_U f dx \leq m_+ \int_V g dq,$$

where $f = g \circ F^{-1}$. Furthermore, we can choose $m_- = 1/4$ and $m_+ = 7/4$.

Proof. — This follows directly from the change of variable formula

$$\int_U f(x) dx = \int_V g(q) (1 - t\kappa(s)) dq,$$

and the fact that $V \subset (0, L) \times (0, 3\delta_0(\Omega)/4)$, so that $t|\kappa(s)| \leq 3/4$. □

By taking $g = 1$, we deduce the following from Lemma 5.1.

COROLLARY 5.2. — *If V is an open set $V \subset (0, L) \times (0, 3\delta_0(\Omega)/4)$ and $U = F(V)$, we have*

$$(13) \quad m_-|V| \leq |U| \leq m_+|V|,$$

where m_+ , m_- are the constants in Lemma 5.1.

COROLLARY 5.3. — *There exists a positive constant M such that, for all $r \in (0, 3\delta_0(\Omega)/4]$,*

$$(14) \quad \tau(r) \leq MLr.$$

Furthermore, we can choose $M = m_+$, where m_+ is the constant in Lemma 5.1.

PROPOSITION 5.4. — *There exists a positive constant K such that the following holds: if V is an open set $V \subset (0, L) \times (0, 3\delta_0(\Omega)/4)$, $U = F(V)$, $u \in H^1(U)$ and $v := u \circ F$, we have*

$$(15) \quad \frac{\int_V |\nabla v|^2 dq}{\int_V v^2 dq} \leq K \frac{\int_U |\nabla u|^2 dx}{\int_U u^2 dx}.$$

Furthermore, we can choose $K = 4m_+$, where m_+ is the constant in Lemma 5.1.

Proof. — From Lemma 5.1, we have directly

$$\int_U u^2 dx \leq m_+ \int_V v^2 dq.$$

On the other hand, for any $q \in V$,

$$\nabla v(q) = J(q)^T (\nabla u) \circ F(q),$$

so that

$$|\nabla u|^2 \circ F(q) = \left| (J(q)^T)^{-1} \nabla v(q) \right|^2.$$

From this we deduce, using the change of variable $x = F(q)$, that

$$\begin{aligned} \int_U |\nabla u|^2 dx &= \int_V \left(\frac{1}{1 - t\kappa(s)} (\partial_s v(q))^2 + (1 - t\kappa(s)) (\partial_t v(q))^2 \right) dq \\ &\geq \frac{1}{4} \int_V |\nabla v|^2 dq. \end{aligned}$$

Putting together the two previous estimates, we obtain Inequality (15). \square

5.2. Number of nodal domains. — To ease the notation in the computations that follow, we set $A := |\Omega|$ in the rest of this section. Let us fix $\varepsilon_0 \in (0, 1)$. We consider an eigenpair (μ, u) for the Neumann Laplacian (as in Subsection 2.1 of [10]) or for $-\Delta_\Omega^\beta$ (as in Section 4 of [10]). We define $u_0 := \varphi_0^\delta u$ and $u_1 := \varphi_1^\delta u$, so that

$$u^2 = u_0^2 + u_1^2.$$

Following Subsection 2.2 of [10], we say that a nodal domain D of u is a *bulk domain* when

$$\int_D u_0^2 dx \geq (1 - \varepsilon_0) \int_D u^2 dx$$

and a *boundary domain* when

$$\int_D u_1^2 dx > \varepsilon_0 \int_D u^2 dx.$$

We denote by $\nu_0(\varepsilon_0, u)$ and $\nu_1(\varepsilon_0, u)$ the number of bulk and boundary domains respectively, so that

$$\nu(u) = \nu_0(\varepsilon_0, u) + \nu_1(\varepsilon_0, u).$$

We now need to bound $\nu_0(\varepsilon_0, u)$ and $\nu_1(\varepsilon_0, u)$ from above. Reproducing the argument in Subsection 2.3 of [10] with $n = 2$, we obtain

$$(16) \quad \nu_0(\varepsilon_0, u) \leq \frac{A}{\Lambda} \left(\frac{1 + \varepsilon_0}{1 - \varepsilon_0} \mu + \left(\frac{1 + \frac{1}{\varepsilon_0}}{1 - \varepsilon_0} \right) \frac{C^2}{\delta^2} \right),$$

where Λ is the first eigenvalue of the Dirichlet Laplacian on the disc of unit area and C is the constant given in the construction of φ_0^δ and φ_1^δ in Section 4.

Let us now consider a boundary domain D_j^1 ($1 \leq j \leq \nu_1(\varepsilon_0, u)$). In a similar way to Subsection 2.4 of [10], we define

$$U := \{x \in D_j^1; u_1(x) \neq 0\} \subset D_j^1 \cap \partial\Omega_{\frac{3\delta}{4}}^+.$$

To simplify the notation, we set $\tilde{u} := u_1$. We consider $V = F^{-1}(U)$ and $\tilde{v} = \tilde{u} \circ F$. We define

$$\sigma(s, t) := (s, -t)$$

and V^R as the interior of the closure of $V \cup \sigma(V)$. We extend \tilde{v} to a function \tilde{v}^R on V^R such that $\tilde{v}^R \circ \sigma = \tilde{v}^R$ (i.e. \tilde{v}^R is even with respect to the variable t). By the Faber-Krahn inequality,

$$\frac{\Lambda}{|V^R|} \leq \frac{\int_{V^R} |\nabla \tilde{v}^R|^2 dq}{\int_{V^R} |\tilde{v}^R|^2 dq},$$

and therefore

$$\frac{\Lambda}{2|V|} \leq \frac{\int_V |\nabla \tilde{v}|^2 dq}{\int_V |\tilde{v}|^2 dq}.$$

Applying Inequalities (13) and (15), we obtain

$$(17) \quad \frac{m_- \Lambda}{2 |U|} \leq K \frac{\int_U |\nabla \tilde{u}|^2 dx}{\int_U |\tilde{u}|^2 dx}.$$

Let us note that in the computation leading to Inequality (17), we implicitly assume that the boundary of V is regular enough, so that V^R is a connected open set and \tilde{v}^R belongs to $H_0^1(V^R)$. In order to avoid such an assumption, we can proceed as in Subsection 2.5 of [10]: we perform the steps indicated above, in a super-level set $V_\alpha = \{\tilde{v}^R > \alpha\}$ for some $\alpha > 0$, and afterwards let α go to 0. By Sard's theorem, we can find a sequence of α 's tending to 0 so that V_α is regular enough (this method was introduced in [6]). Furthermore, as shown in Section 4 of [10], Inequality (17) also holds for an eigenfunction u of $-\Delta_\Omega^\beta$.

Since $\tilde{u} = u_1 = \varphi_1^\delta u$, we find, after applying the Leibniz formula and Young's inequality,

$$\int_U |\nabla \tilde{u}|^2 dx \leq 2 \int_{D_j^1} |\nabla u|^2 dx + \frac{2C^2}{\delta^2} \int_{D_j^1} u^2 dx,$$

and we have, by definition of a boundary domain,

$$\int_U \tilde{u}^2 dx = \int_{D_j^1} u_1^2 dx > \varepsilon_0 \int_{D_j^1} u^2 dx,$$

for our choice of ε_0 . Substituting this into Inequality (17), we get

$$1 \leq \frac{4K |U|}{m_- \Lambda \varepsilon_0} \left(\mu + \frac{C^2}{\delta^2} \right).$$

Summing over all boundary domains, we find

$$\nu_1(\varepsilon_0, u) \leq \frac{4K}{m_- \Lambda \varepsilon_0} \left| \partial \Omega_{\frac{3\delta}{4}}^+ \right| \left(\mu + \frac{C^2}{\delta^2} \right).$$

By Inequality (14) and since $\delta \in (0, \delta_0(\Omega)]$, we have

$$\left| \partial \Omega_{\frac{3\delta}{4}}^+ \right| \leq ML \frac{3\delta}{4},$$

and therefore

$$(18) \quad \nu_1(\varepsilon_0, u) \leq \frac{3KML\delta}{m_- \Lambda \varepsilon_0} \left(\mu + \frac{C^2}{\delta^2} \right).$$

In order to deduce from Inequalities (16) and (18) a bound on $\nu(u)$ which is invariant by scaling, we set

$$\delta := \left(\frac{A}{\mu} \right)^{1/4}.$$

We note that this choice of δ assumes that μ is large enough. That is, $\mu \geq A\delta_0(\Omega)^{-4}$.

For our choice of $\varepsilon_0 \in (0, 1)$, we therefore obtain that

$$(19) \quad \nu(u) \leq \frac{1}{\Lambda} \left(\frac{1 + \varepsilon_0}{1 - \varepsilon_0} (A\mu) + \frac{3KM}{\varepsilon_0 m_-} \rho^2 (A\mu)^{3/4} \right. \\ \left. + C^2 \frac{1 + 1/\varepsilon_0}{1 - \varepsilon_0} (A\mu)^{1/2} + \frac{3KMC^2}{\varepsilon_0 m_-} \rho^2 (A\mu)^{1/4} \right),$$

where the parameter ρ is the square-root of the isoperimetric ratio for the domain Ω :

$$\rho := \frac{L^{1/2}}{A^{1/4}}.$$

5.3. Geometry of the domain. — Let us first sketch how to extend the above results to a multiply connected domain. We now assume only that Ω is an open, bounded and connected set in \mathbb{R}^2 , with a C^2 boundary $\partial\Omega$. The set $\mathbb{R}^2 \setminus \overline{\Omega}$ has an unbounded connected component, which we denote by D_0 . We define $b \in \mathbb{N}$ as the number of bounded connected components of $\mathbb{R}^2 \setminus \overline{\Omega}$, which we denote by D_1, \dots, D_b . For $h \in \{0, 1, \dots, b\}$, ∂D_h is a C^2 -regular closed curve, and we denote its length by L_h . We set

$$s_h := \sum_{i=0}^h L_i$$

for $h \in \{0, 1, \dots, b\}$ and $s_{-1} := 0$. As in Section 3, we choose an arc-length parametrisation $\gamma_h : [s_{h-1}, s_h] \rightarrow \mathbb{R}^2$ of ∂D_h , oriented in such a way that for all $s \in [s_{h-1}, s_h]$, the normal vector $\mathbf{n}_h(s)$ to γ_h at $\gamma_h(s)$ points towards the inside of Ω . Furthermore, we denote by $\kappa_h(s)$ the signed curvature of γ_h at $\gamma_h(s)$. We keep the notation L for the total length of the boundary $\partial\Omega$, so that

$$L = L_0 + L_1 + \dots + L_b.$$

We define

$$E = \bigcup_{h=0}^b (s_{h-1}, s_h).$$

We give a natural generalisation of the definition of the function F in Equation (6): $F : E \times \mathbb{R} \rightarrow \mathbb{R}^2$ is now defined by

$$F(s, t) := \gamma_h(s) + t\mathbf{n}_h(s)$$

if $s \in (s_{h-1}, s_h)$. Extending the definition given in Equation (8), we write

$$t_+(\Omega) := \left(\max_{h \in \{0, \dots, b\}} \sup_{s \in [s_{h-1}, s_h]} |\kappa_h(s)| \right)^{-1},$$

while we continue to define $\underline{\delta}_+(s)$, $\underline{\delta}_+(\Omega)$ and $\delta_0(\Omega)$ by Equations (9), (10) and (11). By reasoning separately on each of the connected components of $\partial\Omega$, we can

show that F is a C^1 diffeomorphism from $E \times (0, \delta_0(\Omega))$ to $\partial\Omega_{\delta_0(\Omega)}^+ \setminus \bigcup_{h=0}^b \ell_h$, where the segment ℓ_h is defined by

$$\ell_h := \gamma_h(s_{h-1}) + (0, \delta_0(\Omega))\mathbf{n}_h(s_{h-1}).$$

Furthermore, the constructions of Section 4 and the estimates of Section 5 still hold.

Let us now discuss how to improve the estimates of Subsection 5.1 for domains with particular geometric properties. We have the following improvement of Corollary 5.3.

PROPOSITION 5.5. — *If Ω is either simply connected or homeomorphic to an annulus, then, for all $\delta \in (0, \delta_0(\Omega)]$,*

$$\tau(\delta) \leq L\delta.$$

Proof. — Let us first consider the simply connected case. From the change of variable $x = F(q)$, we obtain

$$\tau(\delta) = \int_0^L \int_0^\delta (1 - t\kappa(s)) \, dt \, ds = L\delta - \frac{\delta^2}{2} \int_0^L \kappa(s) \, ds.$$

Since γ is a simple, closed, positively-oriented curve,

$$\int_0^L \kappa(s) \, ds = 2\pi,$$

giving the desired inequality (see for instance Corollary 9.5.2 of [3]).

In the case where Ω is homeomorphic to an annulus, $\partial\Omega$ has two connected components. After a similar computation for each of them, we find

$$\tau(\delta) = L\delta - \frac{\delta^2}{2} \left(\int_0^{s_0} \kappa_0(s) \, ds + \int_{s_0}^{s_1} \kappa_1(s) \, ds \right).$$

Since the normal vector has been chosen to point inwards, γ_0 and γ_1 have opposite orientations and the second term in the right-hand side vanishes. The desired inequality is an equality in this case. \square

REMARK 5.6. — Suppose Ω is a multiply-connected domain that is homeomorphic to a disc with $b \in \mathbb{N} \cup \{0\}$ discs removed. By the Gauss-Bonnet theorem, we then have

$$\int_0^L \kappa(s) \, ds = \int_{\partial\Omega} k_g(s) \, ds = 2\pi\chi(\Omega) = 2\pi(1 - b),$$

where $\chi(\Omega)$ is the Euler Characteristic of Ω and k_g denotes the geodesic curvature. If the width of the interior tubular neighbourhood is sufficiently small, that is $\delta \leq \delta_0(\Omega)$, then we obtain

$$\tau(\delta) = L\delta - \pi(1 - b)\delta^2.$$

We recover the cases $b = 0, 1$ as discussed above. If $b \geq 2$, then we have that

$$\tau(\delta) = L\delta + \pi(b-1)\delta^2 > L\delta,$$

for all $\delta \in (0, \delta_0(\Omega)]$. The quantity $L\delta$ is no longer an upper bound of $\tau(\delta)$, but is the leading order term for δ small.

A similar upper bound was obtained in [18], but there the width of the interior tubular neighbourhood need only be smaller than the in-radius of the domain (see also [1]).

5.4. Application to convex domains. — In this subsection, we assume that Ω is a convex domain. We show that Lemma 5.1 and Proposition 5.4 hold for convex domains, and give explicit constants in the corresponding bounds. We first make the following remark.

REMARK 5.7. — Let us assume that Ω is convex. Then $\delta_0(\Omega) = t_+(\Omega)$. Indeed, this is equivalent to saying that a ball of radius smaller than $t_+(\Omega)$ can roll freely inside the set Ω . This follows from a more general result that was established in [4], where this particular case is discussed on page 53, in answer to a question of J.A. Delgado [7]. Furthermore, Inequality (14) holds with $M = 1$ for any $r > 0$. Indeed, let us define the inner parallel set at distance $r > 0$ by $\Omega_r := \Omega \setminus \partial\Omega_r^+$. The set Ω_r is convex. We write $m(r) := |\Omega_r|$ and we have, by definition, $\tau(r) := |\Omega| - m(r)$ for all $r > 0$. According to [11], the function $r \rightarrow m(r)$ is differentiable and $m'(r) = -|\partial\Omega_r|$. Therefore, for $r > 0$,

$$\tau(r) = \int_0^r |\partial\Omega_t| dt.$$

The desired inequality then follows from the fact that the perimeter is non-decreasing with respect to inclusion among convex sets. We note that both results in the present remark hold in arbitrary dimension.

PROPOSITION 5.8. — *If $\Omega \subset \mathbb{R}^2$ is convex, then*

1. *Lemma 5.1 holds with $m_- = 1/4$ and $m_+ = 1$. That is, if V is an open set $V \subset (0, L) \times (0, 3t_+(\Omega)/4)$ and $U = F(V)$ with F as defined in (6), then for any measurable and non-negative function $g : V \rightarrow \mathbb{R}$,*

$$\frac{1}{4} \int_V g dq \leq \int_U f dx \leq \int_V g dq,$$

where $f = g \circ F^{-1}$.

2. *Proposition 5.4 holds with $K = 4$. That is, if V is an open set $V \subset (0, L) \times (0, 3t_+(\Omega)/4)$, $U = F(V)$, $u \in H^1(U)$ and $v := u \circ F$, we have*

$$\frac{\int_V |\nabla v|^2 dq}{\int_V v^2 dq} \leq 4 \frac{\int_U |\nabla u|^2 dx}{\int_U u^2 dx}.$$

Proof. — Since Ω is convex, we have $\kappa(s) \geq 0$ for all $s \in [0, L]$, so that

$$1/4 \leq 1 - t\kappa(s) \leq 1$$

for all $t \in (0, 3t_+(\Omega)/4)$. Using these inequalities in the change of variable formula, we obtain the desired results. \square

6. An upper bound for $R_\Omega^D(\mu)$

As discussed in Section 2, in order to obtain an upper bound for the Courant-sharp Neumann (or Robin) eigenvalues, we require an upper bound for the remainder of the Dirichlet counting function $R_\Omega^D(\mu)$.

In [2], the authors obtain an upper bound for $R_\Omega^D(\mu)$ by using an inner partition of Ω into squares of uniform side-length and Dirichlet bracketing. Suppose these squares have side-length ℓ . Then, in the notation of the present article, Inequality (13) of [2] becomes:

$$R_\Omega^D(\mu) \leq \frac{\tau(\sqrt{2}\ell)}{4\pi}\mu + \frac{|\Omega|}{\sqrt{2}\ell}\mu^{1/2}.$$

Corollary 5.3 gives an upper bound for the volume of the inner tubular neighbourhood provided that the width is sufficiently small. Choosing $\ell > 0$ so that $\sqrt{2}\ell \leq 3\delta_0(\Omega)/4$, and applying Corollary 5.3, we obtain

$$R_\Omega^D(\mu) \leq \frac{\sqrt{2}M\ell L}{4\pi|\Omega|}(|\Omega|\mu) + \frac{|\Omega|^{1/2}}{\sqrt{2}\ell}(|\Omega|\mu)^{1/2}.$$

We then use the freedom in the choice of ℓ to minimise the right-hand side, meaning that we set

$$\ell := \sqrt{\frac{2\pi|\Omega|}{ML}}\mu^{-1/4}$$

to get

$$(20) \quad R_\Omega^D(\mu) \leq \sqrt{\frac{ML|\Omega|}{\pi}}\mu^{3/4} = \sqrt{\frac{M}{\pi}}\rho(\Omega)(|\Omega|\mu)^{3/4},$$

as soon as μ is large enough for $\sqrt{2}\ell \leq 3\delta_0(\Omega)/4$ to be satisfied.

Since we used Corollary 5.3 in this section, Inequality (20) holds with $M = 7/4$, for all $\Omega \subset \mathbb{R}^2$ with a C^2 boundary, as soon as μ is large enough for the condition $\sqrt{2}\ell \leq 3\delta_0(\Omega)/4$ to be satisfied. If we additionally assume Ω to be simply connected or homeomorphic to an annulus, Proposition 5.5 tells us that Inequality (20) holds with $M = 1$ as soon as μ is large enough so that $\sqrt{2}\ell \leq \delta_0(\Omega)$. Finally, if we assume Ω to be convex, Inequality (20) holds with $M = 1$ for all $\mu > 0$.

7. Upper bound for Courant-sharp eigenvalues

We again set $A := |\Omega|$ to ease the notation. Let us consider an eigenpair (μ, u) such that u is a Courant-sharp eigenfunction. We assume that μ is large enough so that

$$(21) \quad \delta = \left(\frac{A}{\mu}\right)^{1/4} \leq \delta_0(\Omega)$$

and

$$(22) \quad \sqrt{2}\ell = \sqrt{\frac{4\pi A}{ML}}\mu^{-1/4} \leq \frac{3}{4}\delta_0(\Omega).$$

We recall that $\varepsilon_0 \in (0, 1)$ is fixed as in Subsection 5.2. By substituting Inequality (19), Equation (4) and Inequality (20) into Inequality (5), we obtain the necessary condition

$$(23) \quad \left(\frac{1}{4\pi} - \frac{1 + \varepsilon_0}{(1 - \varepsilon_0)\Lambda}\right)(A\mu) - \left(\frac{3KM}{\varepsilon_0 m_- \Lambda}\rho^2 + \sqrt{\frac{M}{\pi}}\rho\right)(A\mu)^{3/4} \\ - C^2 \frac{1 + 1/\varepsilon_0}{(1 - \varepsilon_0)\Lambda}(A\mu)^{1/2} - \frac{3KMC^2}{\varepsilon_0 m_- \Lambda}\rho^2(A\mu)^{1/4} < 0.$$

We can reformulate Inequality (23) by saying that if μ is Courant-sharp and satisfies (21) and (22),

$$f_\rho(\xi) := a_0\xi^3 + a_1(\rho)\xi^2 + a_2\xi + a_3(\rho) < 0,$$

with

$$a_0 := \left(\frac{1}{4\pi} - \frac{1 + \varepsilon_0}{(1 - \varepsilon_0)\Lambda}\right); \\ a_1(\rho) := -\left(\frac{3KM}{\varepsilon_0 m_- \Lambda}\rho^2 + \sqrt{\frac{M}{\pi}}\rho\right); \\ a_2 := -C^2 \frac{1 + 1/\varepsilon_0}{(1 - \varepsilon_0)\Lambda}; \\ a_3(\rho) := -\frac{3KMC^2}{\varepsilon_0 m_- \Lambda}\rho^2; \\ \xi := (A\mu)^{1/4}.$$

Therefore, we have that

$$A\mu \leq \xi^*(\rho)^4,$$

where $\xi^*(\rho)$ is the largest real zero of the function $\xi \mapsto f_\rho(\xi)$. We summarise the above discussion in the following proposition.

TABLE 7.1. Possible choices for the constants

| Nature of Ω | Λ | C | m_- | ε_0 | M | K |
|----------------------------|-----------------|-------------|-------|-----------------|-------|-----|
| General case | $\pi j_{0,1}^2$ | $2\sqrt{3}$ | $1/4$ | $1/6$ | $7/4$ | 7 |
| Simply or doubly connected | $\pi j_{0,1}^2$ | $2\sqrt{3}$ | $1/4$ | $1/6$ | 1 | 7 |
| Convex | $\pi j_{0,1}^2$ | $2\sqrt{3}$ | $1/4$ | $1/6$ | 1 | 4 |

PROPOSITION 7.1. — *If μ is a Courant-sharp eigenvalue of the Neumann Laplacian on Ω or of $-\Delta_\Omega^\beta$, we have*

$$\mu \leq \max \left(\frac{4096\pi^2|\Omega|}{81M^2\rho(\Omega)^4\delta_0(\Omega)^4}, \frac{|\Omega|}{\delta_0(\Omega)^4}, \frac{\xi^*(\rho(\Omega))^4}{|\Omega|} \right).$$

We note that the function f_ρ , and therefore also $\xi^*(\rho)$, only depend on the geometric properties of the domain Ω and the choice of ε_0 . Sections 4 and 5 allow us to specify several sets of possible choices of constants in the definition of f_ρ , according to the nature of the domain Ω , which we always assume to be open, bounded and connected with a C^2 boundary. We summarise them in Table 7.1, where $j_{0,1}$ denotes the smallest positive zero of the Bessel function of the first kind J_0 .

8. Geometric upper bounds for Courant-sharp eigenvalues of convex planar domains

8.1. Geometric upper bounds. — In this subsection, we suppose that Ω is an open, bounded, convex planar domain with a C^2 boundary. To simplify the notation, we set $A := |\Omega|$, $t_+ := t_+(\Omega)$ and $\rho := \rho(\Omega)$. Let μ be a Courant-sharp eigenvalue of the Neumann Laplacian on Ω , or of $-\Delta_\Omega^\beta$. We assume that μ is large enough so that Inequality (21) is satisfied. We recall that $\varepsilon_0 \in (0, 1)$ is fixed as in Subsection 5.2.

In order to obtain upper bounds for a Courant-sharp eigenvalue μ in terms of some of the geometric quantities of the domain Ω , we rewrite Inequality (23), with the constants from the last row of Table 7.1, as follows.

$$\begin{aligned} & \left(\frac{1}{4\pi} - \frac{1 + \varepsilon_0}{(1 - \varepsilon_0)\Lambda} \right) (A\mu) \\ & \leq \left(\frac{3KM}{\varepsilon_0 m_- \Lambda} \rho^2 + \sqrt{\frac{M}{\pi}} \rho + \frac{C^2}{(A\mu)^{1/4}} \frac{1 + 1/\varepsilon_0}{(1 - \varepsilon_0)\Lambda} + \frac{3KMC^2}{\varepsilon_0 m_- \Lambda} \frac{\rho^2}{(A\mu)^{1/2}} \right) (A\mu)^{3/4}, \end{aligned}$$

which implies that

$$(24) \quad A\mu \leq \left(\frac{1}{4\pi} - \frac{1+\varepsilon_0}{(1-\varepsilon_0)\Lambda} \right)^{-4} \times \left(\frac{3KM}{\varepsilon_0 m_- \Lambda} \rho^2 + \sqrt{\frac{M}{\pi}} \rho + \frac{C^2}{(A\mu)^{1/4}} \frac{1+1/\varepsilon_0}{(1-\varepsilon_0)\Lambda} + \frac{3KMC^2}{\varepsilon_0 m_- \Lambda} \frac{\rho^2}{(A\mu)^{1/2}} \right)^4.$$

As Ω is convex, $\delta_0(\Omega) = t_+$ (see Remark 5.7). In particular, there exists a ball of radius t_+ which is contained in Ω , that we denote by B_{t_+} . Then $A = |\Omega| \geq |B_{t_+}| = \pi t_+^2$, so $At_+^{-2} \geq \pi$. Together with Inequality (21), we then have

$$A\mu \geq \left(\frac{A}{t_+^2} \right)^2 \geq \pi^2.$$

Substituting this into Inequality (24), we obtain

$$(25) \quad A\mu \leq D_2^{-4} \left(\frac{288\rho^2}{\Lambda} + \frac{\rho}{\pi^{1/2}} + \frac{504}{5\pi^{1/2}\Lambda} + \frac{3456\rho^2}{\pi\Lambda} \right)^4,$$

which implies

$$(26) \quad \mu \leq D_2^{-4} A^{-1} \left(\frac{288\rho^2}{\Lambda} + \frac{\rho}{\pi^{1/2}} + \frac{504}{5\pi^{1/2}\Lambda} + \frac{3456\rho^2}{\pi\Lambda} \right)^4 =: L_1(\rho, A),$$

where $D_2 = \frac{1}{4\pi} - \frac{7}{5\Lambda}$.

With Proposition 7.1 and the preceding discussion in mind, we have that, if μ is Courant-sharp, then

$$(27) \quad \mu \leq \max \left(\frac{A}{t_+^4}, L_1(\rho, A) \right).$$

Note that since Ω is convex, Inequality (14) holds for all $r > 0$ (see Remark 5.7), so Inequality (22) is not necessary in the convex case.

In what follows, we obtain an upper bound for μ that involves some of the other geometric quantities of Ω . We consider Inequality (24) and use the fact that $\mu = \mu_k(\Omega) \geq \mu_2(\Omega)$ for $k \geq 2$. We can then make use of known geometric estimates for the first positive Neumann eigenvalue of Ω . This is sufficient for our purposes due to the monotonicity of the Robin eigenvalues with respect to the function β . That is, for $k \in \mathbb{N}^*$, $\mu_k(\Omega) \leq \mu_k(\Omega, \beta)$.

From Inequality (24), we have

$$(28) \quad A\mu \leq D_2^{-4} \left(\frac{288\rho^2}{\Lambda} + \frac{\rho}{\pi^{1/2}} + \frac{504}{5\Lambda(A\mu_2)^{1/4}} + \frac{3456\rho^2}{\Lambda(A\mu_2)^{1/2}} \right)^4,$$

which implies that

$$(29) \quad \mu \leq D_2^{-4} \left(\frac{288\rho^2}{\Lambda A^{1/4}} + \frac{\rho}{\pi^{1/2} A^{1/4}} + \frac{504}{5\Lambda A^{1/2} \mu_2^{1/4}} + \frac{3456\rho^2}{\Lambda A^{3/4} \mu_2^{1/2}} \right)^4.$$

Since Ω is convex, we invoke the 2-dimensional version of the classical inequality due to Payne and Weinberger, [13], for the first positive Neumann eigenvalue of Ω ,

$$(30) \quad \mu_2(\Omega) \geq \frac{\pi^2}{\text{diam}(\Omega)^2},$$

where $\text{diam}(\Omega)$ denotes the diameter of Ω , to obtain

$$(31) \quad \begin{aligned} \mu &\leq D_2^{-4} \left(\frac{288\rho^2}{\Lambda A^{1/4}} + \frac{\rho}{\pi^{1/2} A^{1/4}} + \frac{504 \text{diam}(\Omega)^{1/2}}{5\Lambda\pi^{1/2} A^{1/2}} + \frac{3456\rho^2 \text{diam}(\Omega)}{\pi\Lambda A^{3/4}} \right)^4 \\ &=: L_2(\rho, A, \text{diam}(\Omega)). \end{aligned}$$

By Proposition 7.1 and the fact that Ω is convex, we have the following inequality:

$$\mu \leq \max \left(\frac{A}{t_+^4}, L_2(\rho, A, \text{diam}(\Omega)) \right).$$

REMARK 8.1. — It is also possible to obtain an upper bound for the largest Courant-sharp eigenvalue of Ω involving only the geometric quantities A , ρ and t_+ via the above approach by making use of Lemma 5.2 of [12] with $K = \Omega$ and L a ball of radius t_+ that is contained in Ω .

We collect the preceding upper bounds in the following proposition.

PROPOSITION 8.2. — *Let Ω be an open, bounded and convex set with a C^2 boundary. If μ is a Courant-sharp eigenvalue of the Neumann Laplacian on Ω or of $-\Delta_\Omega^\beta$, then we have that*

$$\mu \leq \max \left(\frac{|\Omega|}{t_+(\Omega)^4}, L_1(\rho(\Omega), |\Omega|) \right),$$

and

$$\mu \leq \max \left(\frac{|\Omega|}{t_+(\Omega)^4}, L_2(\rho(\Omega), |\Omega|, \text{diam}(\Omega)) \right),$$

where the functions L_1 and L_2 are defined in (26) and (31) respectively.

Suppose that μ is large enough so that (21) holds. Note that since Ω is convex, $\text{diam}(\Omega) \leq \frac{L}{2}$ and so $\mu_2(\Omega) \geq \frac{4\pi^2}{L^2}$, by the Payne-Weinberger inequality. Substituting this into (28), we obtain

$$\mu \leq \frac{\rho^4}{AD_2^4} \left(\frac{288\rho}{\Lambda} + \frac{1}{\pi^{1/2}} + \frac{504}{5(2\pi)^{1/2}\Lambda} + \frac{3456\rho^3}{2\pi\Lambda} \right)^4 =: A^{-1}D_2^{-4}\rho^4G(\rho)^4,$$

where $\rho \mapsto G(\rho)$ is an increasing function for $\rho \geq 0$.

In fact, in this convex case, we can deduce a simplified upper bound for μ and obtain a geometric upper bound for the number of Courant-sharp eigenvalues.

PROPOSITION 8.3. — *Let Ω be an open, bounded and convex set with a C^2 boundary. Let $\mu = \mu_k(\Omega)$ or $\mu = \mu_k(\Omega, \beta)$ where β is a non-negative Lipschitz continuous function on $\partial\Omega$. There exist positive constants C and C' that do not depend on Ω such that, if μ is Courant-sharp, then*

$$\mu \leq C \left(\frac{|\Omega|}{t_+(\Omega)^4} + \frac{\rho(\Omega)^8}{|\Omega|} \right),$$

and

$$k \leq C' \left(\frac{|\Omega|^2}{t_+(\Omega)^4} + \rho(\Omega)^8 \right).$$

The first inequality of Proposition 8.3 follows from Inequalities (25) and (27), by using the Isoperimetric inequality $\rho \geq \sqrt{2}\pi^{\frac{1}{4}}$. The second inequality follows from the first one and Corollary A.2, together with Remark A.3, after applying Young's inequality repeatedly. Indeed, by following this procedure, we obtain

$$C = D_2^{-4} \left(\frac{288}{\Lambda} + \frac{1}{\sqrt{2}\pi^{3/4}} + \frac{35064}{10\pi\Lambda} \right)^4 \approx 8.98 \times 10^{17},$$

and $C' \leq C + 1$.

We remark that this proves Proposition 1.1 since the latter is Proposition 8.3 restricted to the Neumann case.

8.2. Application to the disc of unit area. — For the disc $D \subset \mathbb{R}^2$ of unit area, we have $\rho(D) = (4\pi)^{1/4}$ and $\delta_0(D) = t_+(D) = \pi^{-1/2}$. We wish to evaluate Proposition 7.1, so we must first calculate $\xi^*(\rho)$. We use the last line of Table 7.1. The value of $\xi^*(\rho)$ can be computed either numerically or with Cardano's formula. We find, for any Courant-sharp Neumann (or Robin) eigenvalue $\mu(D)$ of D ,

$$\mu(D) \leq 2.67 \times 10^{17}.$$

It is well-known that the first positive Neumann eigenvalue of D is $\pi(j'_{1,1})^2$ where $j'_{1,1}$ is the first positive zero of J'_1 , the derivative of the Bessel function J_1 . By substituting this into Inequality (29), we obtain that any Courant-sharp Neumann eigenvalue $\mu(D)$ of D satisfies

$$\mu(D) \leq 1.26 \times 10^{20}.$$

We also have $\text{diam}(D) = 2\pi^{-1/2}$. By Inequality (31), we have that any Courant-sharp Neumann (or Robin) eigenvalue $\mu(D)$ of D satisfies

$$\mu(D) \leq 2.09 \times 10^{20}.$$

By comparing the two preceding upper bounds, we remark that we do not lose a great deal by appealing to the bound of Payne and Weinberger, even though it is not the natural choice for our situation. Indeed, equality is achieved

in (30) for a rectangle of fixed area with one side shrinking to 0. Such shrinking behaviour is not possible in our situation since Ω contains a ball of radius $t_+(\Omega) > 0$.

We now evaluate the first inequality of Proposition 8.2 for the disc of unit area. We obtain that any Courant-sharp Neumann (or Robin) eigenvalue $\mu(D)$ of D satisfies

$$\mu(D) \leq 1.42 \times 10^{20},$$

which is better than the second inequality of Proposition 8.2 for the disc of unit area, but worse than using Inequality (29) directly (see above).

In addition, Proposition 8.3 applied to the disc of unit area gives that any Courant-sharp Neumann (or Robin) eigenvalue $\mu(D)$ of D satisfies

$$\mu(D) \leq 1.51 \times 10^{20},$$

and the number of such eigenvalues $k \leq 1.51 \times 10^{20}$.

For comparison with the above upper bounds, we recall that the third positive Neumann eigenvalue is the largest Courant-sharp Neumann eigenvalue as proved in [9]. In addition, $\mu_4(D) = \pi(j'_{2,1})^2 \approx 3.054^2\pi < 30$ (where $j'_{2,1}$ is the first positive zero of J'_2 , the derivative of the Bessel function J_2). The Courant-sharp Robin eigenvalues of D are not known (to date).

9. Generalisation to any dimension

9.1. Tubular neighbourhoods of the boundary. — We now assume that Ω is a bounded, open and connected set in \mathbb{R}^n , with $n \geq 3$, such that $\partial\Omega$ is a C^2 submanifold of \mathbb{R}^n . We extend the methods used in the previous sections to this situation.

To simplify the notation, we write Γ for the submanifold $\partial\Omega$. As before, for $r > 0$, we define the inner tubular neighbourhood with radius r :

$$\Gamma_r^+ := \{x \in \Omega; \text{dist}(x, \Gamma) < r\}.$$

For all $x' \in \Gamma$, we denote by $\mathbf{n}(x')$ the outward unit normal vector at x' . In order to parametrise the inner tubular neighbourhoods, we introduce the normal bundle of Γ . Furthermore, for all $x' \in \Gamma$, we identify the one-dimensional vector space spanned by $\mathbf{n}(x')$ with the real line \mathbb{R} . We therefore write the normal bundle as a trivial product:

$$N(\Gamma) := \Gamma \times \mathbb{R}$$

and we define a C^1 mapping $F : N(\Gamma) \rightarrow \mathbb{R}^n$ by

$$(32) \quad F(x', t) = x' - t\mathbf{n}(x').$$

We recall that the mapping $x' \mapsto \mathbf{n}(x')$ from Γ to \mathbb{S}^{n-1} is known as the *Gauss map* (see for instance Subsection 2.5 of [17]). Its differential is a sym-

metric linear endomorphism of $T_{x'}\Gamma$, the tangent space to Γ at x' , seen as a subspace of \mathbb{R}^n . It is called the *Weingarten endomorphism*, and we denote it by $W_{x'}$. Its eigenvalues are called the *principal curvatures* of Γ at x' . Let $(\mathbf{e}_1, \dots, \mathbf{e}_{n-1})$ be an orthonormal basis of $T_{x'}\Gamma$ consisting of eigenvectors of $W_{x'}$ and let $(\kappa_1(x'), \dots, \kappa_{n-1}(x'))$ be the associated principal curvatures. Then $(\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, 1)$ and $(\mathbf{e}_1, \dots, \mathbf{e}_{n-1}, \mathbf{n}(x'))$ are orthonormal bases of $T_{(x',t)}N(\Gamma)$, identified with $T_{x'}\Gamma \oplus \mathbb{R}$, and \mathbb{R}^n respectively, and the matrix of the differential $D_{(x',t)}F$ in those bases is

$$\begin{pmatrix} 1 - t\kappa_1(x') & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 - t\kappa_{n-1}(x') & 0 \\ 0 & \cdots & 0 & -1 \end{pmatrix}.$$

We now define

$$t_+(\Omega) := \left(\sup_{x' \in \Gamma} \max_{i \in \{1, \dots, n-1\}} |\kappa_i(x')| \right)^{-1},$$

and, for any $x' \in \Gamma$, the (internal) *cut-distance* to Γ at x' :

$$\delta_+(x') := \sup\{\delta > 0; \text{dist}(F(x', t), \Gamma) = t \text{ for all } t \in [0, \delta]\}.$$

We set

$$\underline{\delta}_+(\Omega) := \inf_{x' \in \Gamma} \delta_+(x')$$

and

$$\delta_0(\Omega) := \min\{t_+(\Omega), \underline{\delta}_+(\Omega)\}.$$

By construction, F is a diffeomorphism of class C^1 from $\Gamma \times (0, \delta_0(\Omega))$ to $\Gamma_{\delta_0(\Omega)}^+$.

LEMMA 9.1. — *Let V be an open set in $\Gamma \times (0, \delta_0(\Omega))$ and let $U = F(V)$. For any measurable, non-negative function $g : V \rightarrow \mathbb{R}$,*

$$\int_U f(x) dx = \int_\Gamma \left(\int_0^{+\infty} \mathbf{1}_V(x', t) g(x', t) \prod_{i=1}^{n-1} (1 - t\kappa_i(x')) dt \right) dx'$$

where $f = g \circ F^{-1}$, $\mathbf{1}_V$ is the characteristic function of V and dx' is the surface measure in Γ induced by the Lebesgue measure in \mathbb{R}^n .

Proof. — The proof is similar to the proof of Weyl's formula for the volume of tubes given in Chapter 6 of [3]. First, we use Proposition 3.3.16 of [3] to get

$$\int_U f(x) dx = \int_V g F^*(dx),$$

where dx denotes the Lebesgue measure in \mathbb{R}^n and $F^*(dx)$ the pullback of dx by the diffeomorphism F .

TABLE 9.1. Possible choices for the constants in \mathbb{R}^n

| Nature of Ω | C | m_- | M | K |
|--------------------|-------------|-------------|---------------|-----------|
| General case | $2\sqrt{3}$ | $1/4^{n-1}$ | $(7/4)^{n-1}$ | 7^{n-1} |
| Convex | $2\sqrt{3}$ | $1/4^{n-1}$ | 1 | 4^{n-1} |

Then, we note that, up to a change of sign, the mapping F defined in Equation (32) is the canonical map defined in Equation 2.7.5 of [3]. According to Equation 6.8.2 of [3], there exists a function G defined on $\Gamma \times (0, \delta_0(\Omega))$, to be specified below, such that

$$F^*(dx) = G(x', t) dx' \otimes dt,$$

where $dx' \otimes dt$ denotes the product measure in $N(\Gamma) = \Gamma \times \mathbb{R}$, with dt the Lebesgue measure in \mathbb{R} . From Equation 6.7.16 of [3], we obtain

$$\begin{aligned} \int_V g F^*(dx) &= \int_{\Gamma \times \mathbb{R}} \mathbf{1}_V(x', t) g(x', t) G(x', t) dx' \otimes dt \\ &= \int_{\Gamma} \left(\int_{\mathbb{R}} \mathbf{1}_V(x', t) g(x', t) G(x', t) dt \right) dx'. \end{aligned}$$

Finally, we give an explicit formula for $G(x', t)$. We fix $u \mapsto \mathbf{x}(u)$ a local parametrisation of Γ such that $\mathbf{x}(0) = x'$ and, for $i \in \{1, \dots, n-1\}$,

$$\frac{\partial \mathbf{x}}{\partial u_i}(0) = \mathbf{e}_i.$$

From Equation 6.8.9 and Proposition 6.6.2 of [3], we obtain

$$G(x', t) = \frac{\det(\Pi(\frac{\partial \mathbf{x}}{\partial u_1}(0) - t \frac{\partial \mathbf{n} \circ \mathbf{x}}{\partial u_1}(0)), \dots, \Pi(\frac{\partial \mathbf{x}}{\partial u_{n-1}}(0) - t \frac{\partial \mathbf{n} \circ \mathbf{x}}{\partial u_{n-1}}(0)))}{\det(\Pi(\frac{\partial \mathbf{x}}{\partial u_1}(0)), \dots, \Pi(\frac{\partial \mathbf{x}}{\partial u_{n-1}}(0)))},$$

where Π is the orthogonal projection from \mathbb{R}^n to $T_{x'}\Gamma$. It follows from the chain rule that, for all $i \in \{1, \dots, n-1\}$,

$$\frac{\partial \mathbf{n} \circ \mathbf{x}}{\partial u_i}(0) = W_{x'}[\mathbf{e}_i] = \kappa_i(x') \mathbf{e}_i.$$

This implies that

$$G(x', t) = \prod_{i=1}^{n-1} (1 - t \kappa_i(x')). \quad \square$$

Lemma 9.1 allows us to extend the estimates of Section 5 to higher dimensions in a straightforward manner. Indeed, Lemma 5.1, Corollaries 5.2 and 5.3, and Proposition 5.4 hold with the constants given in Table 9.1.

9.2. Upper bounds for Courant-sharp eigenvalues. — To simplify the formulas, we write $V := |\Omega|$ and $S := |\partial\Omega|$ (the total $(n-1)$ -dimensional surface measure of $\partial\Omega$) in the rest of this section. We denote by N_Ω either N_Ω^N or N_Ω^β , since the lower bound that we derive holds for both types of counting function.

Following [9], we define

$$\gamma(n) := \frac{(2\pi)^n}{\omega_n \Lambda(n)^{\frac{n}{2}}},$$

and we recall that $\gamma(n) < 1$ for all integers $n \geq 2$. For the rest of this section, we set

$$(33) \quad \varepsilon_0 := \frac{1}{2} \cdot \frac{1 - \gamma(n)^{\frac{2}{n}}}{1 + \gamma(n)^{\frac{2}{n}}}.$$

This choice of ε_0 will be explained below.

We consider (μ, u) , an eigenpair of the Neumann Laplacian on Ω or of $-\Delta_\Omega^\beta$, with μ large enough, in a sense to be made precise.

Following the steps of Subsection 5.2, we obtain an upper bound for the number of bulk domains

$$\nu_0(\varepsilon_0, u) \leq \frac{V}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1 + \varepsilon_0}{1 - \varepsilon_0} \mu + \left(\frac{1 + \frac{1}{\varepsilon_0}}{1 - \varepsilon_0} \right) \frac{C^2}{\delta^2} \right)^{\frac{n}{2}},$$

where $\Lambda(n)$ is the first eigenvalue of the Dirichlet Laplacian on a ball of volume 1 in \mathbb{R}^n . More explicitly,

$$\Lambda(n) := \omega_n^{\frac{2}{n}} j_{\frac{n}{2}-1,1}^2,$$

where ω_n denotes the volume of a ball of radius 1 in \mathbb{R}^n and $j_{\frac{n}{2}-1,1}$ the smallest positive zero of the Bessel function of the first kind $J_{\frac{n}{2}-1}$. We also obtain the following upper bound for the number of boundary domains

$$\nu_1(\varepsilon_0, u) \leq \frac{3 \cdot 2^{\frac{n}{2}-1} K^{\frac{n}{2}} S M \delta}{m_- \Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} \left(\mu + \frac{C^2}{\delta^2} \right)^{\frac{n}{2}}.$$

Both bounds hold assuming $\delta \in (0, \delta_0(\Omega)]$. Let us now set

$$\delta := \left(\frac{V^{\frac{2}{n}}}{\mu} \right)^{\frac{1}{4}}.$$

It follows from the two previous upper bounds that, if μ is large enough so that

$$(34) \quad \left(\frac{V^{\frac{2}{n}}}{\mu} \right)^{\frac{1}{4}} \leq \delta_0(\Omega),$$

then we have the following upper bound for the nodal count

$$(35) \quad \nu(u) \leq \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \xi + \frac{1+\frac{1}{\varepsilon_0}}{1-\varepsilon_0} C^2 \xi^{1/2} \right)^{\frac{n}{2}} + \frac{3 \cdot 2^{\frac{n}{2}-1} M K^{\frac{n}{2}} \rho^2}{m_{-\varepsilon_0}^{\frac{n}{2}}} \xi^{-\frac{1}{4}} \left(\xi + C^2 \xi^{1/2} \right)^{\frac{n}{2}} \right),$$

where $\xi := V^{2/n} \mu$ and $\rho := S^{1/2} / V^{1/2-1/2n}$.

An upper bound for the remainder $R_\Omega^D(\mu)$ is given in Section 2 of [2] for arbitrary dimension. Repeating the argument in Section 6, we obtain, for any $\ell \in (0, 3(4\sqrt{n})^{-1}\delta_0(\Omega))$,

$$R_\Omega^D(\mu) \leq \frac{\omega_n M S \sqrt{n} \ell}{(2\pi)^n} \mu^{\frac{n}{2}} + \frac{\pi n^{\frac{3}{2}} \omega_n V}{(2\pi)^n \ell} \mu^{\frac{n-1}{2}}.$$

Setting

$$\ell = \sqrt{\frac{\pi n V}{M S}} \mu^{-\frac{1}{4}},$$

we find that, if μ is large enough so that

$$(36) \quad \sqrt{\frac{\pi n V}{M S}} \mu^{-\frac{1}{4}} \leq \frac{3\delta_0(\Omega)}{4\sqrt{n}},$$

we have the following upper bound for the remainder

$$(37) \quad R_\Omega^D(\mu) \leq \frac{2n\omega_n}{(2\pi)^n} \sqrt{\pi M} \rho \xi^{\frac{n}{2}-\frac{1}{4}}.$$

As before, if the eigenfunction u is Courant-sharp,

$$\nu(u) = N_\Omega(\mu) + 1,$$

and therefore, if Inequalities (34) and (36) are satisfied, from Inequalities (35) and (37), we obtain

$$(38) \quad \begin{aligned} f_\rho(\xi) := & \frac{\omega_n}{(2\pi)^n} \xi^{\frac{n}{2}} - \frac{2n\omega_n \sqrt{\pi M}}{(2\pi)^n} \rho \xi^{\frac{n}{2}-\frac{1}{4}} \\ & - \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \xi + \frac{1+\frac{1}{\varepsilon_0}}{1-\varepsilon_0} C^2 \xi^{\frac{1}{2}} \right)^{\frac{n}{2}} \\ & - \frac{3 \cdot 2^{\frac{n}{2}-1} M K^{\frac{n}{2}}}{\Lambda(n)^{\frac{n}{2}} m_{-\varepsilon_0}^{\frac{n}{2}}} \rho^2 \xi^{-\frac{1}{4}} \left(\xi + C^2 \xi^{\frac{1}{2}} \right)^{\frac{n}{2}} < 0. \end{aligned}$$

A straightforward computation shows that, since

$$0 < \varepsilon_0 < \frac{1 - \gamma(n)^{\frac{2}{n}}}{1 + \gamma(n)^{\frac{2}{n}}},$$

we have

$$\lim_{\xi \rightarrow +\infty} f_\rho(\xi) = +\infty.$$

Taking into account the conditions (34), (36) and (38), we can summarise our result in the following proposition.

PROPOSITION 9.2. — *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be an open, bounded and connected set with a C^2 boundary. If μ is a Courant-sharp eigenvalue of the Neumann Laplacian on Ω or of $-\Delta_\Omega^\beta$, we have*

$$\mu \leq \max \left(\left(\frac{|\Omega|^{\frac{1}{2n}}}{\delta_0(\Omega)} \right)^4, \left(\sqrt{\frac{\pi}{M}} \frac{4n|\Omega|^{\frac{1}{2n}}}{3\rho(\Omega)\delta_0(\Omega)} \right)^4, \frac{\xi^*(\rho(\Omega))}{|\Omega|^{\frac{2}{n}}} \right),$$

with

$$\xi^*(\rho) := \sup\{\xi > 0; f_\rho(\xi) < 0\},$$

the function $t \mapsto f_\rho(t)$ being defined in Formula (38).

We note that the function f_ρ , and therefore also $\xi^*(\rho)$, only depend on the dimension n , the geometric properties of the domain Ω and the choice of ε_0 .

9.3. Geometric upper bounds for Courant-sharp eigenvalues of convex bodies.

— We now assume that the set Ω is convex. In that case, $t_+(\Omega) = \underline{\delta}_+(\Omega)$, as discussed in Remark 5.7, and therefore $\delta_0(\Omega) = t_+(\Omega)$. To simplify the formulas, we write $t_+ := t_+(\Omega)$. Again ε_0 is fixed as in (33). Additionally, the constants appearing on the left-hand side of Inequality (38) can be chosen as in the second row of Table 9.1, which implies a smaller value of $\xi^*(\rho)$. Furthermore, again from Remark 5.7, Corollary 5.3 holds, with $M = 1$ and for any $r > 0$. The estimate (37) on the remainder therefore holds without Condition (36). We conclude that, if μ is Courant-sharp, then

$$\mu \leq \max \left(\left(\frac{V^{\frac{1}{2n}}}{t_+} \right)^4, \frac{\xi^*(\rho)}{V^{\frac{2}{n}}} \right).$$

We now obtain an explicit upper bound for $\xi^*(\rho)$.

From Inequality (38) and the constants from the second row of Table 9.1, we have

$$\begin{aligned} (39) \quad & \frac{\omega_n}{(2\pi)^n} \xi^{\frac{n}{2}} - \frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho \xi^{\frac{n}{2}-\frac{1}{4}} - \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \xi + 12 \frac{1+\frac{1}{\varepsilon_0}}{1-\varepsilon_0} \xi^{\frac{1}{2}} \right)^{\frac{n}{2}} \\ & - \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} \rho^2 \xi^{-\frac{1}{4}} \left(\xi + 12 \xi^{\frac{1}{2}} \right)^{\frac{n}{2}} < 0. \end{aligned}$$

In what follows, we use the fact that, by (34),

$$\xi = V^{2/n} \mu \geq \left(\frac{V^{1/n}}{t_+} \right)^4 \geq \omega_n^{4/n},$$

since $V = |\Omega| \geq |B_{t_+}| = \omega_n t_+^n$ as $B_{t_+} \subset \Omega$.

We also invoke the Mean Value theorem which implies that for $N \geq 1$, there exists $z \in (0, x)$ such that

$$(1+x)^N - 1 = N(1+z)^{N-1}x.$$

We have that

$$\begin{aligned} & \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \xi + 12 \frac{1+\frac{1}{\varepsilon_0}}{1-\varepsilon_0} \xi^{\frac{1}{2}} \right)^{\frac{n}{2}} \\ &= \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \xi^{\frac{n}{2}} \left(1 + 12 \frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \xi^{-\frac{1}{2}} \right)^{\frac{n}{2}} \\ &\leq \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \xi^{\frac{n}{2}} \left(1 + 6n \left(1 + 12 \frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \xi^{-\frac{1}{2}} \right)^{\frac{n}{2}-1} \frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \xi^{-\frac{1}{2}} \right) \\ &\leq \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \xi^{\frac{n}{2}} \left(1 + 6n \left(1 + 12 \frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \xi^{-\frac{1}{2}} \right) \\ &= \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \xi^{\frac{n}{2}} \\ &\quad + \frac{6n}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \xi^{\frac{n-1}{2}}, \end{aligned}$$

and

$$(\xi + 12\xi^{\frac{1}{2}})^{\frac{n}{2}} = \xi^{\frac{n}{2}} (1 + 12\xi^{-\frac{1}{2}})^{\frac{n}{2}} \leq \xi^{\frac{n}{2}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}}.$$

Substituting these into (39), we obtain

$$\begin{aligned} (40) \quad & \left(\frac{\omega_n}{(2\pi)^n} - \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \right) \xi^{\frac{n}{2}} \\ &< \left(\frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho + \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}} \rho^2 \right) \xi^{\frac{n}{2}-\frac{1}{4}} \\ &\quad + \frac{6n}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \xi^{\frac{n-1}{2}}. \end{aligned}$$

We note that the coefficients of $\xi^{\frac{n}{2}}, \xi^{\frac{n-1}{2}}$ only depend on the dimension, while the coefficient of $\xi^{\frac{n}{2}-\frac{1}{4}}$ depends on the dimension and the isoperimetric ratio.

Inequality (40) implies that

$$(41) \quad \xi < D_n^{-1} \left(\frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho + \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}} \rho^2 \right) \xi^{\frac{3}{4}} \\ + \frac{6n}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \xi^{\frac{3}{4}} \xi^{-\frac{1}{4}},$$

where $D_n := \frac{\omega_n}{(2\pi)^n} - \frac{1}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}}$. Substituting $\xi^{-\frac{1}{4}} \leq \omega_n^{-\frac{1}{n}}$ into Inequality (41), we obtain

$$(42) \quad \xi < D_n^{-4} \left(\frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho + \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}} \rho^2 \right. \\ \left. + \frac{6n\omega_n^{-\frac{1}{n}}}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \right)^4,$$

which implies that

$$(43) \quad \mu < D_n^{-4} V^{-2/n} \left(\frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho + \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}} \rho^2 \right. \\ \left. + \frac{6n\omega_n^{-\frac{1}{n}}}{\Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \right)^4 \\ =: M_1(n, \rho, V).$$

If instead we consider the bound due to Payne and Weinberger, [13],

$$(44) \quad \mu_2(\Omega) \geq \frac{\pi^2}{\text{diam}(\Omega)^2},$$

then

$$\xi^{-\frac{1}{4}} \leq \frac{\text{diam}(\Omega)^{\frac{1}{2}}}{\sqrt{\pi} V^{\frac{1}{2n}}}.$$

The following bound due to Gritzmann, Wills and Wrase, [8],

$$\text{diam}(\Omega) < \frac{S^{n-1}}{\omega_{n-1}(nV)^{n-2}},$$

gives that

$$\xi^{-\frac{1}{4}} \leq \frac{\rho^{n-1}}{\sqrt{\pi} \omega_{n-1}^{\frac{1}{2}} n^{\frac{n}{2}-1}}.$$

By substituting the latter inequality into Inequality (41), we have

$$\begin{aligned}
 (45) \quad \mu &< D_n^{-4} V^{-\frac{2}{n}} \left(\frac{2n\omega_n\sqrt{\pi}}{(2\pi)^n} \rho + \frac{3 \cdot 2^{\frac{n}{2}-1} (4^{n-1})^{\frac{n}{2}+1}}{\Lambda(n)^{\frac{n}{2}} \varepsilon_0^{\frac{n}{2}}} (1 + 12\omega_n^{-\frac{2}{n}})^{\frac{n}{2}} \rho^2 \right. \\
 &\quad + \frac{6\rho^{n-1}}{\sqrt{\pi}\omega_{n-1}^{\frac{1}{2}} n^{\frac{n}{2}-1} \Lambda(n)^{\frac{n}{2}}} \left(\frac{1+\varepsilon_0}{1-\varepsilon_0} \right)^{\frac{n}{2}} \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \\
 &\quad \left. \times \left(1 + 12 \left(\frac{1+\frac{1}{\varepsilon_0}}{1+\varepsilon_0} \right) \omega_n^{-\frac{2}{n}} \right)^{\frac{n}{2}-1} \right)^4 \\
 &=: M_2(n, \rho, V).
 \end{aligned}$$

We can summarise both versions of the upper bound in the following proposition.

PROPOSITION 9.3. — *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be an open, bounded and convex set with a C^2 boundary. If μ is a Courant-sharp eigenvalue of the Neumann Laplacian on Ω or of $-\Delta_\Omega^\beta$, we have*

$$(46) \quad \mu \leq \max \left(\frac{|\Omega|^{\frac{2}{n}}}{t_+(\Omega)^4}, M_1(n, \rho(\Omega), |\Omega|) \right)$$

and

$$\mu \leq \max \left(\frac{|\Omega|^{\frac{2}{n}}}{t_+(\Omega)^4}, M_2(n, \rho(\Omega), |\Omega|) \right),$$

with M_1 and M_2 defined in Formulas (43) and (45) respectively.

Similarly to the end of Subsection 8.1, we can find simpler, although less explicit, upper bounds.

PROPOSITION 9.4. — *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be an open, bounded and convex set with a C^2 boundary. Let $\mu = \mu_k(\Omega)$ or $\mu = \mu_k(\Omega, \beta)$, with β a non-negative Lipschitz continuous function on $\partial\Omega$. There exist positive constants C_n and C'_n that do not depend on Ω such that, if μ is Courant-sharp,*

$$\mu \leq C_n \left(\frac{|\Omega|^{\frac{2}{n}}}{t_+(\Omega)^4} + \frac{\rho(\Omega)^8}{|\Omega|^{\frac{2}{n}}} \right)$$

and

$$k \leq C'_n \left(\left(\frac{|\Omega|^{\frac{1}{n}}}{t_+(\Omega)} \right)^{2n} + \rho(\Omega)^{4n} \right).$$

The first inequality in Proposition 9.4 follows from Inequalities (42) and (46), taking into account $\rho(\Omega) \geq \sqrt{n}\omega_n^{\frac{1}{2n}}$. The second follows from the first and Corollary A.2, together with Remark A.3, after applying Young's inequality repeatedly. Both C_n and C'_n could be computed explicitly.

As described in the introduction for the case of dimension 2, the previous inequality implies that a sufficiently regular convex set with a large number of Courant-sharp eigenvalues has a large isoperimetric ratio or a point in its boundary where the curvature is large (or both). Additionally, if there exists a Courant-sharp eigenvalue which is large with respect to $|\Omega|^{\frac{2}{n}}t_+(\Omega)^{-4}$, the set has a large isoperimetric ratio.

Acknowledgments: KG was supported by the Swiss National Science Foundation grant no. 200021_163228 entitled *Geometric Spectral Theory* during the academic year 2016/2017. KG also acknowledges support from the Max Planck Institute for Mathematics, Bonn, from 1 October 2017 to 31 July 2018. CL acknowledges support from the ERC (Project COMPAT, ERC-2013-ADG no. 339958), the Portuguese FCT (Project OPTFORMA, IF/00177/2013) and the Swedish Research Council (Grant D0497301). Both authors wish to thank the referee for their insightful comments and suggestions.

Appendix A. An upper bound for the Neumann counting function of a convex domain

PROPOSITION A.1. — *Let Ω be an open, bounded and convex set in \mathbb{R}^n with a C^2 boundary. Then, for all $\mu > 0$, we have*

$$(47) \quad N_{\Omega}^N(\mu) \leq \frac{n^{\frac{n}{2}}}{\pi^n} |\Omega| \mu^{\frac{n}{2}} + \frac{n^{\frac{n}{2}}}{\pi^n} |\partial\Omega| \sum_{i=0}^{n-1} \frac{\binom{n-1}{i} \pi^{i+1}}{(i+1)t_+(\Omega)^i} \mu^{\frac{n-i-1}{2}},$$

where $t_+(\Omega)$ is the smallest radius of curvature, as defined by Equation (8).

Proof. — We combine the Payne-Weinberger lower bound (see Inequality (44)) with Neumann bracketing. Reference [2] uses a similar approach with Dirichlet bracketing.

For $\alpha \in \mathbb{Z}^n$, we define the open hypercube $C_{\alpha} := \alpha + (0, 1)^n$. Let us fix $a > 0$. We define

$$\mathcal{I}_a := \{\alpha \in \mathbb{Z}^n ; aC_{\alpha} \cap \Omega \neq \emptyset\}$$

and denote by $K(a)$ the cardinality of \mathcal{I}_a . For all $\alpha \in \mathcal{I}_a$, we define $\Omega_{\alpha,a} := \Omega \cap aC_{\alpha}$. By construction, $\Omega_{\alpha,a}$ is a convex, open set with diameter bounded from above by $\sqrt{n}a$. We define

$$\tilde{\Omega}_a := \bigcup_{\alpha \in \mathcal{I}_a} \Omega_{\alpha,a},$$

which is open and bounded, but not connected.

We consider the spectrum of the Neumann Laplacian on $\tilde{\Omega}_a$, which we denote by $(\mu_k(a))_{k \geq 1}$. Since $\tilde{\Omega}_a$ has $K(a)$ connected components, 0 is an eigenvalue of multiplicity $K(a)$, that is

$$0 = \mu_1(a) = \mu_2(a) = \cdots = \mu_{K(a)}(a).$$

The eigenvalue $\mu_{K(a)+1}(a)$ is the smallest among the non-trivial eigenvalues of the sets $\Omega_{\alpha,a}$ with $\alpha \in \mathcal{I}_a$, and therefore, according to the Payne-Weinberger inequality,

$$\frac{\pi^2}{na^2} \leq \mu_{K(a)+1}(a).$$

It follows from the variational characterisation of the eigenvalues that $\mu_k(a) \leq \mu_k(\Omega)$ for any integer k (see, for example, Proposition 4(c) in Section XIII.15 of [16]), so that

$$\frac{\pi^2}{na^2} \leq \mu_{K(a)+1}(\Omega).$$

Equivalently, the counting function $N_{\Omega}^N(\cdot)$ satisfies

$$(48) \quad N_{\Omega}^N\left(\frac{\pi^2}{na^2}\right) \leq K(a).$$

To finish the proof, let us give an upper bound for $K(a)$. We define D_a as the interior of

$$\overline{\bigcup_{\alpha \in \mathcal{I}_a} aC_{\alpha}},$$

and note that $|D_a| = K(a)a^n$. Furthermore, we have

$$D_a \subset \Omega + \sqrt{na}B,$$

where B is the ball of radius 1 in \mathbb{R}^n and the right-hand side is understood as a Minkowski sum. We obtain

$$(49) \quad K(a) \leq a^{-n} |\Omega + \sqrt{na}B|.$$

Using normal coordinates in the exterior of Ω (similarly to Subsection 9.1 but for the outer parallel sets), we find, for any $\delta > 0$,

$$|\Omega + \delta B| = |\Omega| + \int_0^{\delta} \left(\int_{\partial\Omega} \prod_{i=1}^{n-1} (1 + t\kappa_i(x')) dx' \right) dt.$$

Expanding and using $\kappa_i(x') \leq t_+(\Omega)^{-1}$ for all $x' \in \partial\Omega$ and $i \in \{1, \dots, n-1\}$, we find

$$|\Omega + \delta B| \leq |\Omega| + |\partial\Omega| \sum_{i=0}^{n-1} \frac{\binom{n-1}{i} \delta^{i+1}}{(i+1)t_+^i(\Omega)}.$$

We now choose $\delta := \sqrt{na} := \pi/\sqrt{\mu}$. We obtain the desired result by substituting this value into the previous inequality and by using Inequalities (49) and (48) successively. \square

Let us denote by $F_n(|\Omega|, |\partial\Omega|, t_+(\Omega), \mu)$ the right-hand side of Inequality (47). It is a continuous and increasing function of μ .

COROLLARY A.2. — *Let Ω be an open, bounded and convex set in \mathbb{R}^n with a C^2 boundary. For any integer $k \geq 1$,*

$$(50) \quad k \leq F_n(|\Omega|, |\partial\Omega|, t_+(\Omega), \mu_k(\Omega)).$$

Proof. — Let us fix $\varepsilon > 0$. We have $N_\Omega^N((1 + \varepsilon)\mu_k(\Omega)) \geq k$. We obtain Inequality (50) by applying Inequality (47) and letting ε go to 0. \square

REMARK A.3. — We recall that $\mu_k(\Omega) \leq \mu_k(\Omega, \beta)$ for every non-negative Lipschitz continuous function β on $\partial\Omega$, and every $k \geq 1$. Since $F_n(|\Omega|, |\partial\Omega|, t_+(\Omega), \mu)$ is increasing in μ , Inequalities (47) and (50) hold when substituting $N_\Omega^\beta(\mu)$ for $N_\Omega^N(\mu)$ and $\mu_k(\Omega, \beta)$ for $\mu_k(\Omega)$, respectively.

BIBLIOGRAPHY

- [1] C. BUNDLE – “Convexity and its applications”, (P. M. Gruber & J. M. Wills, éd.), Birkhäuser, Basel, 1983.
- [2] M. VAN DEN BERG & K. GITTINS – “On the number of Courant-sharp Dirichlet eigenvalues”, *J. Spec. Theory* **6** (2016), p. 735–745.
- [3] M. BERGER & B. GOSTIAUX – *Differential geometry: manifolds, curves, and surfaces*, Graduate Texts in Mathematics, no. 115, Springer-Verlag, New York, 1988, Translated from French by Silvio Levy.
- [4] J. N. BROOKS & J. B. STRANTZEN – “Blaschke’s rolling theorem in \mathbf{R}^n ”, *Memoirs of the American Mathematical Society* **80** (1989), no. 405.
- [5] P. BÉRARD & B. HELFFER – “The weak Pleijel theorem with geometric control”, *J. Spec. Theory* **6** (2016), p. 717–733.
- [6] P. BÉRARD & D. MEYER – “Inégalités isopérimétriques et applications”, *Ann. Sci. École Norm. Sup. (4)* **15** (1982), no. 3, p. 513–541.
- [7] J. A. DELGADO – “Blaschke’s theorem for convex hypersurfaces”, *J. Differential Geom.* **14** (1979), no. 4, p. 489–496.
- [8] P. GRITZMANN, J. M. WILLS & D. WRASE – “A new isoperimetric inequality”, *J. reine angew. Math.* **379** (1987), p. 22–30.
- [9] B. HELFFER & M. PERSSON SUNDQVIST – “On nodal domains in Euclidean balls”, *Proc. Amer. Math. Soc.* **144** (2016), p. 4777–4791.
- [10] C. LÉNA – “Pleijel’s nodal domain theorem for Neumann and Robin eigenfunctions”, *Ann. Inst. Fourier* **69** (2019), no. 1, p. 283–301.

- [11] G. MATHERON – “La formule de Steiner pour les érosions”, *J. Appl. Probability* **15** (1978), no. 1, p. 126–135.
- [12] E. MILMAN – “On the role of convexity in isoperimetry, spectral gap and concentration”, *Inventiones mathematicae* **177** (2009), no. 1, p. 1–43.
- [13] L. E. PAYNE & H. F. WEINBERGER – “An optimal Poincaré inequality for convex domains”, *Archive for Rational Mechanics and Analysis* **5** (1960), no. 1, p. 286–292.
- [14] J. PEETRE – “A generalization of Courant’s nodal domain theorem”, *Math. Scand.* **5** (1957), p. 15–20.
- [15] Å. PLEIJEL – “Remarks on Courant’s nodal line theorem”, *Comm. Pure Appl. Math.* **9** (1956), p. 543–550.
- [16] M. REED & B. SIMON – *Methods of modern mathematical physics IV, Analysis of Operators*, Academic Press, New York, 1978.
- [17] R. SCHNEIDER – *Convex bodies: The Brunn-Minkowski theory*, Encyclopedia of Mathematics and Its Applications, no. 151, Cambridge University Press, 2013.
- [18] B. SZ.-NAGY – “Über Parallelmengen nichtkonvexer ebener Bereiche”, *Acta Sci. Math. Szeged* **20** (1959), no. 36, p. 47.