

Astérisque

HENK BROER

Coupled Hopf-bifurcations: Persistent examples of n -quasiperiodicity determined by families of 3-jets

Astérisque, tome 286 (2003), p. 223-229

[<http://www.numdam.org/item?id=AST_2003__286__223_0>](http://www.numdam.org/item?id=AST_2003__286__223_0)

© Société mathématique de France, 2003, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**COUPLED HOPF-BIFURCATIONS:
PERSISTENT EXAMPLES OF n -QUASIPERIODICITY
DETERMINED BY FAMILIES OF 3-JETS**

by

Henk Broer

Abstract. — In this note examples are presented of vector fields depending on parameters and determined by the 3-jet, which display persistent occurrence of n -quasiperiodicity. In the parameter space this occurrence has relatively large measure. A leading example consists of weakly coupled Hopf bifurcations. This example, however, is extended to full generality in the space of all 3-jets.

1. Introduction

In the theory of coupled reaction diffusion equations the following is of interest, see Poláčik *et al.* [12, 19]. The problem is whether persistent examples exist of (parameter dependent) dynamical systems with the following properties:

- (1) Occurrence of n -quasiperiodicity in a measure theoretically significant way.
- (2) The system is local, and the property is determined by a low order jet.
- (3) Preferably parameters are only needed in the linear part.

Below we present a solution to this problem by means of coupled Hopf families. To fix thoughts, we start with an example.

Example 1 (Weakly coupled Hopf bifurcations). — Consider a C^∞ -system of n weakly coupled Hopf bifurcations, near the origin of $\mathbb{R}^{2n}$ given by

$$(1) \quad \begin{pmatrix} \dot{x}_j \\ \dot{y}_j \end{pmatrix} = \begin{pmatrix} \alpha_j - \beta_j & \\ \beta_j & \alpha_j \end{pmatrix} \begin{pmatrix} x_j \\ y_j \end{pmatrix} - (x_j^2 + y_j^2) \begin{pmatrix} x_j \\ y_j \end{pmatrix} + O(r^4),$$

$1 \leq j \leq n$, where $r^2 = \sum_{j=1}^n (x_j^2 + y_j^2)$. The lower order part (the 3-jet) consists of n completely decoupled Hopf bifurcations, as already considered in [15]. Presently, however, we include the coupling term $O(r^4)$. Moreover, we include $2n$ parameters

2000 Mathematics Subject Classification. — 37L10, 70K43.

Key words and phrases. — Coupled Hopf bifurcations, center manifold reduction, quasiperiodic tori, quasiperiodic bifurcation.

$(\alpha, \beta) = (\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n)$, where $\alpha \in \mathbb{R}_+^n$ is small and where β varies over any compact disc $L \subset \mathbb{R}_+^n$. In multi-polar coordinates $x_j = r_j \cos \varphi_j$, $y_j = r_j \sin \varphi_j$ the decoupled lower order part reads

$$(2) \quad \begin{aligned} \dot{\varphi}_j &= \beta_j \\ \dot{r}_j &= r_j(\alpha_j - r_j^2), \end{aligned}$$

$1 \leq j \leq n$. Clearly for $\alpha_j > 0$, system (2) has an n -torus attractor $r_j = \sqrt{\alpha_j}$, where the dynamics is parallel given by $\dot{\varphi}_j = \beta_j$, $1 \leq j \leq n$. Our interest is with the fate of this dynamical phenomenon upon addition of the higher order perturbation $O(r^4)$.

Below we generalize the setting of Example 1, raising a similar problem. To solve this we apply both Center Manifold Theory [14] and KAM Theory in the dissipative setting [17, 2, 6, 3, 7]. We summarize the results of our investigation. First, for small $|\alpha|$, the family of n -tori is C^ℓ -persistent for such perturbations, where the bound on $|\alpha|$ depends on ℓ . Second, the continuum of parallel dynamics persists as a Whitney smooth family of quasiperiodic attractors, foliated over a Cantor set. Projected to the (α, β) -parameter space, this Cantor foliation has positive measure, expressed in terms of a Lebesgue density point of quasiperiodicity corresponding to $\alpha = 0$. Notably, the dynamics in between generically break up due to internal resonance: upon variation of parameters the dynamics can be asymptotically periodic (e.g., phase locked) or chaotic ($n \geq 3$), [20, 18, 21, 23].

Let us briefly outline the contents of this paper. We start by generalizing the setting of Example 1, and developing an appropriate perturbation model for the application of Center Manifold and KAM Theory. We end by a general discussion, pointing towards some interesting problems regarding quasiperiodic Hopf bifurcation that occur in a subordinate way.

2. Coupled Hopf-bifurcations

2.1. Setting of the problem. — Instead of weakly coupled case (1) we here consider the more general system

$$(3) \quad \begin{pmatrix} \dot{x}_j \\ \dot{y}_j \end{pmatrix} = \begin{pmatrix} \alpha_j - \beta_j & \\ \beta_j & \alpha_j \end{pmatrix} \begin{pmatrix} x_j \\ y_j \end{pmatrix} + O(r^2),$$

which will be subject to suitable C^3 -open conditions. As in Example 1 we include dependence on the parameter vector $(\alpha, \beta) \in \mathbb{R}_+^n \times \mathbb{R}_+^n$. We apply a standard normal form procedure to system (3), e.g. compare [22, 4, 24, 11]. Note that the linear part of (3) has a $\mathbb{T}^n$ -symmetry. Strong resonances are excluded by requiring that

$$(4) \quad \sum_{j=1}^n \beta_j k_j \neq 0 \text{ whenever } 0 < \sum_{j=1}^n |k_j| \leq 4,$$

which amounts to the first open condition. Granted (4), by near identity, polynomial changes of variables this $\mathbb{T}^n$ -symmetry can be pushed over the whole 3-jet, which then in appropriate multi-polar coordinates reads

$$(5) \quad \begin{aligned} \dot{\varphi}_j &= \beta_j - f_j(r_1^2, \dots, r_n^2) + O(r^4) \\ \dot{r}_j &= r_j(\alpha_j - g_j(r_1^2, \dots, r_n^2)) + O(r^4), \end{aligned}$$

$1 \leq j \leq n$. As in Example 1, we truncate the $O(r^4)$ term, so arriving at the present generalization of (2), which we explore for invariant n -tori. Therefore we expand

$$\begin{aligned} f_j &= f_{j1}r_1^2 + \dots + f_{jn}r_n^2 \\ g_j &= g_{j1}r_1^2 + \dots + g_{jn}r_n^2, \end{aligned}$$

with f_{ji}, g_{ji} constants, $i, j = 1, 2, \dots, n$. Invariant n -tori then are determined by n equations

$$(6) \quad g_{j1}r_1^2 + \dots + g_{jn}r_n^2 = \alpha_j,$$

$1 \leq j \leq n$. Consider the ‘action-space’ $\mathbb{R}_+^n = \{r_1^2, \dots, r_n^2\}$, where the equations (6) determine n hyperplanes. Considering the $n \times n$ -matrix

$$G = \left((g_{j,i})_{j,i=1}^n \right),$$

we impose further C^3 -open conditions

$$(7) \quad \det G \neq 0 \text{ while } G^{-1}(\alpha) \in \mathbb{R}_+^n.$$

By $c^2 := G^{-1}(\alpha)$ denote the unique (transversal) intersection point of the hyperplanes. Then the equations (6) have the unique solution

$$(8) \quad r_1^2 = c_1^2, \dots, r_n^2 = c_n^2, \text{ with } c_j^2 = c_j^2(\alpha).$$

$1 \leq j \leq n$, which determines our invariant n -torus, carrying parallel dynamics.

Remark. — The open conditions (7) are trivially satisfied in Example 1, where $G = \text{Id}_n$. One easily detects other concrete examples that are C^3 -nearby (1).

As announced in Example 1, the problem is to study the effect of the higher order perturbation $O(r^4)$ on this family of tori. As said before, to answer this we shall apply both Center Manifold Theory [14] and KAM Theory in the dissipative setting [6, 3, 7].

2.2. An appropriate perturbative setting. — We formulate a perturbation problem suitable for our purposes. Introducing the small parameter ε and putting $I_j = r_j - c_j$, we scale

$$\begin{aligned} \alpha_j &= \varepsilon^2 \bar{\alpha}_j & \beta_j &= \bar{\beta}_j \\ r_j &= \varepsilon \bar{r}_j & \varphi_j &= \bar{\varphi}_j, \end{aligned}$$

also writing $I_j = \varepsilon \bar{I}_j$, $1 \leq j \leq n$. This gives the estimates

$$\begin{aligned}\dot{\bar{\varphi}}_j &= \bar{\beta}_j + \varepsilon O(\bar{I}) \\ \dot{\bar{I}}_j &= -2\varepsilon^2 \bar{\alpha}_j \bar{I}_j + \varepsilon^2 \bar{I}_j O(\sqrt{\bar{\alpha} \bar{I}}),\end{aligned}$$

$1 \leq j \leq n$, which are uniform for $(\bar{\alpha}, \bar{\beta}) \in K \times L$, for any given compact subsets $K, L \subseteq \mathbb{R}_+^n$. A further scaling

$$\bar{I} = \varepsilon^q \bar{\bar{I}},$$

for a fixed $q > 1$, leads to the following perturbation problem:

$$(9) \quad \begin{aligned}\dot{\bar{\varphi}}_j &= \bar{\beta}_j + O(\varepsilon^{1+q}) \\ \dot{\bar{\bar{I}}}_j &= -2\varepsilon^2 \bar{\alpha}_j \bar{\bar{I}}_j + O(\varepsilon^{2+q}),\end{aligned}$$

$1 \leq j \leq n$, again with uniform estimates. Since $q > 1$, the form (9) is suitable for application of the Center Manifold Theorem [14], Thm. 4.1, implying the C^ℓ -persistence of the invariant n -torus for small values of ε .

To further investigate persistence of the quasiperiodic dynamics we apply dissipative KAM Theory as developed in [6] §§ 4 and 8. In the unperturbed case

$$\begin{aligned}\dot{\bar{\varphi}}_j &= \bar{\beta}_j \\ \dot{\bar{\bar{I}}}_j &= -2\varepsilon^2 \bar{\alpha}_j \bar{\bar{I}}_j,\end{aligned}$$

$1 \leq j \leq n$, we single out parameter vectors $\beta \in L$ such that for all $k \in \mathbb{Z}^n \setminus \{0\}$ Diophantine conditions

$$(10) \quad |\langle k, \beta \rangle| \geq \frac{\gamma}{|k|^\tau}$$

hold. Here $\tau > n - 1$ is a constant, while we choose $\gamma = c\varepsilon^q$, for an appropriate (sufficiently small) constant c , depending on K and L . These conditions define a Cantor foliation $\mathcal{C}_{\varepsilon, c} \subseteq K \times L$ for the unperturbed system. The complement $(K \times L) \setminus \mathcal{C}_{\varepsilon, c}$ has measure $O(\varepsilon^q)$ as $\varepsilon \downarrow 0$. The main result of the present paper is:

Theorem 2 (Perturbation Theorem). — *Consider system (9), with parameter vectors $(\bar{\alpha}, \bar{\beta}) \in K \times L$, for given compact subsets $K, L \subseteq \mathbb{R}_+^n$. Also let $\ell \in \mathbb{N}$ be sufficiently large. Then, for $\varepsilon > 0$ and sufficiently small the (9) has the following properties:*

- (1) *The unperturbed n -torus family $I = 0$ persists as a unique C^ℓ -family of hyperbolic n -torus attractors $\mathcal{T}_\varepsilon$, also depending C^ℓ on ε .*
- (2) *For parameter values $(\bar{\alpha}, \bar{\beta}) \in \mathcal{C}_{\varepsilon, c}$ as described above, with c sufficiently small, the unperturbed tori persist as quasiperiodic tori inside $\mathcal{T}_\varepsilon$.*
- (3) *The union of tori inside $\mathcal{T}_\varepsilon$ with non-quasiperiodic dynamics has Lebesgue measure $O(\varepsilon^q)$, as $\varepsilon \downarrow 0$, $1 \leq j \leq n$, uniformly in $(\bar{\alpha}, \bar{\beta}) \in K \times L$.*

Proof (sketch). — For simplicity we perform the KAM Theory inside the center manifold, see [6], the Appendix. This means that for some $\ell' < \ell$, there exists a $C^{\ell'}$ -reparametrization $\Phi_\varepsilon : K \times L \rightarrow \mathbb{R}_+^n \times \mathbb{R}_+^n$, extending to a $C^{\ell'}$ -diffeomorphism

$$\Psi_\varepsilon : \mathbb{T}^n \times \{0\} \times K \times L \longrightarrow \mathbb{R}^{2n} \times \mathbb{R}_+^n \times \mathbb{R}^n$$

such that for $(\bar{\alpha}, \bar{\beta}) \in \mathcal{C}_\varepsilon$, the map Ψ_ε is a conjugacy from the unperturbed system to the full (perturbed) system (9).⁽¹⁾ The dependence of the map Ψ_ε on ε is smooth.

This implies that (9) has a subsystem consisting of a Whitney smooth foliation of quasiperiodic n -tori. Moreover, in the $C^{\ell'}$ -topology $|\Psi_\varepsilon - \text{Id}| = o(\varepsilon)$ as $\varepsilon \downarrow 0$, which implies that the image $\Phi(\mathcal{C}_\varepsilon)$ has positive (almost full) measure as $\varepsilon \downarrow 0$. $\square$

Returning to the context of system (3) we conclude

Corollary 3. — *Consider system (3), with parameter vectors $\bar{\alpha} \in K$, and $\bar{\beta} \in L$, for any given compact sets $K, L \subset \mathbb{R}_+^n$ with L not containing any strong resonances (4). Also let $\ell \in \mathbb{N}$ be sufficiently large. Then, up to condition (7), for $\varepsilon > 0$, sufficiently small, the system (3) has the properties 1., 2. and 3. of Theorem 2. Moreover, the behaviour described above is persistent under sufficiently C^3 -small perturbations.*

The $\mathcal{T}_\varepsilon$ -dynamics in between the Cantor foliation generically break up due to internal resonance. Finally notice that the weakly coupled case of Example 1 also is fully covered by the Corollary.

2.3. Conclusive remarks, towards quasiperiodic Hopf bifurcation. — Concerning the asymptotics of the measure estimate, we note that the KAM Theorem [6, 7] may well be applied in the C^∞ -setting, keeping track of the normal linear part. This does not lead to contradictions, since the quasiperiodic tori by parallelity are of class C^∞ , since they are ℓ normally hyperbolic for any ℓ . A further (quasiperiodic) normalizing and reparametrizing (see, e.g., [3]) leads to sharper estimates on the complement of quasiperiodicity, which become of arbitrary large order in ε , as $\varepsilon \downarrow 0$, compare [7], §5.2. Assuming real analyticity, even exponentially small estimates can be obtained for the complement of quasiperiodicity. Compare [7, 16] and further references given there.

Next we point out a relationship with quasiperiodic bifurcation theory. Until now we restricted $\bar{\alpha}$ to compact subsets K of the open cone $\mathbb{R}_+^n$, thereby only covering interior points. Consider the $\bar{\alpha}$ -regime near the boundary of this cone. In the unperturbed system (2) of Example 1, this gives normally elliptic subtori, and clearly the perturbative cases involve subordinate quasiperiodic Hopf bifurcations, compare [3, 7, 8, 5, 25, 9, 10, 23]. In the present, general case the theory will be even more interesting, compare [13, 26], for general reference also see [1]. Investigation of

⁽¹⁾To be precise, $K \times L$ has to be shrunk at its boundary by a layer of thickness $O(\varepsilon^q)$.

this will require further research, as is also indicated by the following example, kindly provided by Florian Wagener.

Example 4 (Two weakly coupled Hopf families). — Consider the ‘integrable’ system

$$\begin{aligned}\dot{\varphi}_1 &= \beta_1 \\ \dot{\varphi}_2 &= \beta_2 \\ \dot{r}_1 &= r_1(\alpha_1 - r_1^2) - r_2^4 \\ \dot{r}_2 &= r_2(\alpha_2 - r_2^2).\end{aligned}$$

A brief calculation reveals that the non-hyperbolic tori occur for

$$\begin{aligned}27\alpha_2^4 &= 4\alpha_1^3 r_1 = \frac{1}{\sqrt{3}}\sqrt{\alpha_1} r_2 = \sqrt{\alpha_2}, \\ \alpha_2 &= 0 \quad r_1 = \sqrt{\alpha_1} \quad r_2 = 0.\end{aligned}$$

The problem is to study the effects of integrable and non-integrable higher order terms. Notice that the hyperplanes in parameter space where subordinate quasiperiodic Hopf bifurcations take place, have shifted somewhat. However, by the scaling all interior points of the cone are drawn within the regime with invariant n -tori.

Observe that this program is reminiscent to the Hamiltonian case of elliptic subtori, [6, 3, 7, 16].

Acknowledgements. — Helpful discussions are acknowledged with Peter Poláčik, Florian Wagener, Carles Simó, Floris Takens and Bernold Fiedler. Also thanks is due to the Universitat de Barcelona for hospitality during part of the preparation of this paper.

References

- [1] C. Baesens, J. Guckenheimer, S. Kim and R.S. MacKay, Three coupled oscillators: Mode-locking, global bifurcation and toroidal chaos, *Physica D*, (1991), 387–475.
- [2] B.L.J. Braaksma & H.W. Broer, On a quasi-periodic Hopf bifurcation, *Ann. Institut Henri Poincaré, Analyse non linéaire*, **4**, no.2 (1987), 115–168.
- [3] B.L.J. Braaksma, H.W. Broer and G.B. Huitema, Towards a Quasiperiodic Bifurcation Theory, *Mem. AMS*, **83**(421) (1990), 83–175.
- [4] H.W. Broer, Formal normal forms for vector fields and some consequences for bifurcations in the volume preserving case. In: D. Rand and L.S. Young (eds.), *Dynamical Systems and Turbulence, Warwick 1980 LNM* **898** (1981), 75–89, Springer-Verlag.
- [5] H.W. Broer, KAM-Theory: Multi-Periodicity in conservative and dissipative systems, *Nieuw Arch. Wisk.* **14**(1), (1996), 1–15.
- [6] H.W. Broer, G.B. Huitema and F. Takens, Unfoldings of Quasi-Periodic Tori, *Mem. AMS*, **83**(421) (1990), 1–82.
- [7] H.W. Broer, G.B. Huitema and M.B. Sevryuk, Quasi-periodic tori in families of dynamical systems: order amidst chaos, *LNM* **1645**, Springer Verlag 1996.
- [8] H.W. Broer & F. Takens, Mixed spectrum and rotational symmetry, *Arch. Rational Mech. An.* **124** (1993), 13–42.

- [9] H.W. Broer, F. Takens and F.O.O. Wagener, Integrable and non-integrable deformations of the skew Hopf bifurcation, *Reg. and Chaot. Dyn.* **4** (1999), 17–43.
- [10] H.W. Broer and F.O.O. Wagener, Quasi-periodic stability of subfamilies of an unfolded skew-Hopf bifurcation, *Arch. Rational Mech. An.* **152** (2000), 283–326.
- [11] S-N. Chow, C. Li and D. Wang, *Normal Forms and Bifurcation of Planar Vector Fields*. Cambridge University Press 1994.
- [12] B. Fiedler and P. Poláčik, Complicated dynamics of scalar reaction diffusion equations with a nonlocal term, *Proc. Roy. Soc. Edinburgh Sect. A, (Mathematical and Physical Sciences)* **115**(1-2) (1990), 167–192.
- [13] J. Guckenheimer and P. Holmes, Nonlinear Oscillations, Dynamical Systems and Bifurcations of Vector Fields, *Appl. Math. Sciences* **42**, Springer Verlag 1983.
- [14] M.W. Hirsch, C.C. Pugh and M. Shub, Invariant Manifolds, *LNM* **583**, Springer Verlag 1977.
- [15] E. Hopf, A mathematical example displaying features of turbulence. *Commun. Appl. Math.* **1** (1948), 303–322.
- [16] A. Jorba and J. Villanueva, On the Normal Behaviour of Partially Elliptic Lower-Dimensional Tori of Hamiltonian Systems, *Nonlinearity* **10** (1997), 783–822.
- [17] J.K. Moser, Convergent series expansions for quasi-periodic motions, *Math. Ann.* **169** (1967), 136–176.
- [18] S.E. Newhouse, D. Ruelle and F. Takens, Occurrence of strange Axiom A attractors near quasiperiodic flows on $\mathbb{T}^m$, $m \leq 3$, *Commun. Math. Phys.* **64** (1978), 35–40.
- [19] P. Poláčik and V. Šošovička, Stable periodic solutions of a spatially homogeneous non-local reaction-diffusion equation, *Proc. Roy. Soc. Edinburgh Sect. A, (Mathematical and Physical Sciences)* **126**(4) (1996), 867–884.
- [20] D. Ruelle and F. Takens, On the nature of turbulence, *Commun. Math. Phys.* **20** (1971), 167–192; **23** (1971), 343–4.
- [21] C. Simó, Perturbations of translations in the two-dimensional torus: the case near resonance, *Proceedings VI CEDYA*, Universidad de Zaragoza, 1984.
- [22] F. Takens, Singularities of Vector Fields, *Commun. Math. Phys.* **20**, 1970, 167–192, **23**, (1971), 343–344. *Publ. I.H.E.S.* **43** (1974), 47–100.
- [23] F. Takens and F.O.O. Wagener, Resonances in skew and reducible quasi-periodic Hopf bifurcations, *Nonlinearity* **13** (2000), 377–396.
- [24] A. Vanderbauwhede, Centre Manifolds, Normal Forms and Elementary Bifurcations, *Dynamics Reported* **2** (1989), 89–170.
- [25] F.O.O. Wagener, On the skew Hopf bifurcation. *PhD thesis*, University of Groningen, 1998.
- [26] H. Zoladek, Bifurcation of a certain family of planar vector fields tangent to axes, *Journ. Diff. Eqns.* **67** (1987), 1–55.

H. BROER, Department of Mathematics and Computer Science, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands