BULLETIN DE LA S. M. F.

JAN KUBARSKI About Stefan's definition of a foliation with singularities : a reduction of the axioms

Bulletin de la S. M. F., tome 118, nº 4 (1990), p. 391-394

http://www.numdam.org/item?id=BSMF_1990__118_4_391_0

© Bulletin de la S. M. F., 1990, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (http://smf. emath.fr/Publications/Bulletin/Presentation.html) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/legal.php). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

\mathcal{N} umdam

Article numérisé dans le cadre du programme Numérisation de documents anciens mathématiques http://www.numdam.org/ Bull. Soc. math. France, 118, 1990, p. 391-394.

ABOUT STEFAN'S DEFINITION OF A FOLIATION WITH SINGULARITIES : A REDUCTION OF THE AXIOMS

ΒY

JAN KUBARSKI (*)

RÉSUMÉ. — L'article présent concerne la définition d'un feuilletage de Stefan. Le résultat principal de cet article est le fait qu'un axiome de la définition d'un feuilletage de Stefan [4, chap. 1] est une conséquence des autres.

ABSTRACT. — The aim of this paper is to give an accurate proof of the fact formulated in [3, p. 45] that one of the axioms of Stefan's foliations [4, chap. 1] follows from the remaining ones.

The following definitions of a *foliation with singularities* comes from the work by P. STEFAN [4].

Suppose V is a connected Hausdorff C^{∞} and paracompact (equivalently and with a countable basis) manifold of dimension n. By a *foliation of* V with singularities we mean a partition \mathcal{F} of V into sets such that :

(1) for each element $L \in \mathcal{F}$, there exists a structure of differentiable manifold σ on L such that

- (i) (L, σ) is a connected immersed submanifold of V,
- (ii) (L, σ) is a *leaf* of V with respect to all locally connected topological spaces, i.e. if X is an arbitrary locally connected topological space and $f: X \to V$ is a continuous function such that $f[X] \subset L$, then $f: X \to (L, \sigma)$ is continuous;

(2) for each $x \in V$, there exists a local chart φ on V around x with the following properties :

- (a) φ is a surjection $D_{\varphi} \to U_{\varphi} \times W_{\varphi}$ where U_{φ} , W_{φ} are open neighbourhoods of 0 in \mathbb{R}^k and \mathbb{R}^{n-k} , respectively, and k is the dimension of the leaf through x (denoted by L_x);
- (b) $\varphi(x) = (0,0);$

^(*) Texte reçu le 19 mai 1988, révisé le 26 septembre 1990.

J. KUBARSKI, Institute of Mathematics, Technical University of Lodz, Al. Politechniki 11, 90-924 Lodz, Pologne.

BULLETIN DE LA SOCIÉTÉ MATHÉMATIQUE DE FRANCE 0037-9484/1990/391/\$ 5.00 © Société mathématique de France

(c) if $L \in \mathcal{F}$, then $\varphi[L \cap D_{\varphi}] = U_{\varphi} \times \ell_{\varphi,L}$ where

$$\ell_{\varphi,L} = \left\{ w \in W_{\varphi} : \varphi^{-1}(0,w) \in L \right\}.$$

A chart φ which fulfils the above condition is called *distinguished* around x.

THEOREM. — Let \mathcal{F} be a partition of V into connected immersed submanifolds of V, fulfilling (2). Then \mathcal{F} is a foliation with singularities.

REMARK. — This theorem is formulated in [3, p. 45] without an accurate proof. The author say that it easily follows in the same way as in the case without singularities, indicating [1]. It turns out that this theorem needs a subtler proof. The reasoning as in [1] gives the proof provided some added assumption

(1)
$$\frac{\partial}{\partial \varphi^i}\Big|_y \in T_y(L_x) \text{ for } i \leq k \text{ and all } y \in L_x \cap D_{\varphi}, \ k = \dim L_x,$$

is satisfied, which is exactly the body of Stefan's lemma [4, Lemma 3.1]. That this added condition follows from the remaining ones is the aim of our paper.

Proof of the Theorem : according to STEFAN ([4, Lemma 3.1]), it is sufficient to show that each distinguished chart $\varphi = (\varphi^1, \ldots, \varphi^n)$ around x has the property (1).

Assume to the contrary that, for a distinguished chart φ around x, this property does not hold at a point $y_0 \in L_x \cap D_{\varphi}$. Then, of course, there exists a vector $v \in T_{y_0}(L_x)$ such that

(2)
$$\varphi_{*y_0}(v) \notin T_{(\hat{y}_0, c_0)} \left(U_{\varphi} \times \{c_0\} \right)$$

where $(\hat{y}_0, c_0) = \varphi(y_0), \, \hat{y}_0 \in U_{\varphi}, \, c_0 \in W_{\varphi}.$

Take any smooth curve

$$c: (-\varepsilon, \varepsilon) \longrightarrow L_x, \quad \varepsilon > 0,$$

such that $c(0) = y_0$, $\dot{c}(0) = v$ and $\operatorname{Im} c \subset D_{\varphi}$. Consider the curve $\varphi \circ c : (-\varepsilon, \varepsilon) \to \mathbb{R}^n$. Let $pr_2 : U_{\varphi} \times W_{\varphi} \to W_{\varphi}$ denotes the projection onto the second factor. By (2) :

$$(pr_2)_*((\varphi \circ c)(0)) \neq 0.$$

томе 118 — 1990 — N° 4

392

Diminishing $\varepsilon > 0$, if necessary, we may assume that

$$pr_2 \circ \varphi \circ c : (-\varepsilon, \varepsilon) \longrightarrow W_{\varphi}$$

is an embedding. Denote the set $\text{Im}(pr_2 \circ \varphi \circ c)$ by *I*. Of course,

$$U_{\varphi} \times I \subset U_{\varphi} \times \ell_{\varphi, L_x}$$

(because $I \subset pr_2 \circ \varphi[D_{\varphi} \cap L_x] = \ell_{\varphi,L_x}$) and $U_{\varphi} \times I$ is a k + 1-dimensional hypersurface of \mathbb{R}^n , thus a locally compact space. Put — for brevity —

$$M := L_x \cap D_x$$

understanding it as an open submanifold of L_x , and consider the injective immersion

$$ar{arphi}: M \longrightarrow \mathbb{R}^n, \quad x \longmapsto \varphi(x).$$

By the above $\bar{\varphi}[M] \supset U_{\varphi} \times I$.

For each point $x \in M$, we choose a neighbourhood $U(x) \subset M$ of x such that

$$\bar{\varphi}|_{U(x)}: U(x) \longrightarrow \mathbb{R}^n$$

is an embedding. By the assumption of the second axiom of countability of V, each connected immersed submanifold of V fulfils this axiom (see Appendix). Then M, as an open submanifold of the manifold L_x , has a countable basis. Choose a countable open covering $\{U_i; i \in \mathbb{N}\}$ of M such that each \overline{U}_i is compact and contained in some $U(x_i)$. We prove that

$$\bar{\varphi}[\bar{U}_i] \cap (U_{\varphi} \times I)$$

— as a subset of the space $U_{\varphi} \times I$ — has no interior. We have a little more, namely that the set $\bar{\varphi}[U(x_i)] \cap (U_{\varphi} \times I)$ has no interior. If it were not, then by taking an nonempty and open subset $U \subset U_{\varphi} \times I$ such that $X \subset \bar{\varphi}[U(x_i)]$, we would obtain the mapping

$$\left(\tilde{\varphi}|_{U(x_i)}\right)^{-1}|_X: X \to U(x_i)$$

from a (k + 1)-dimensional manifold to a k-dimensional one, being an immersion, which is not possible. Thus $U_{\varphi} \times I$ is an union of a countable sequence of nowhere dense sets

$$\Big\{\bar{\varphi}\big[\bar{U}_i\big]\cap (U_\varphi\times I)\,;\,\,i\in\mathbb{N}\Big\},$$

which leads to a contradiction with Baire's theorem for locally compact spaces. The theorem is proved. \Box

BULLETIN DE LA SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Appendix : The following theorem is well known; here we give a simple proof of it.

THEOREM. — Each connected immersed submanifold L of a C^{∞} Hausdorff paracompact manifold V has a countable basis.

Proof. — Let $f: L \to V$ be an immersion. The assumptions imply the existence of a Riemann tensor G on V. Its pullback f^*G is a Riemann tensor on L. A connected manifold which possesses a Riemann tensor is separable [2], therefore it has a countable basis. \Box

BIBLIOGRAPHIE

- [1] CHEVALLEY (C.). Theory of Lie groups. Princeton University Press, 1946.
- [2] HELGASON (S.). Differential Geometry and symmetric spaces. New York and London, Academic Press, 1962.
- [3] PRADINES (J.). Graph and holonomy of singular foliations. Preprint, Santiago de Compostela, September 1984.
- [4] STEFAN (P.). Accessible sets, orbits, and foliations with singularities, Proc. London Math. Soc., t. 29, 1974, p. 699–713.

томе 118 — 1990 — №° 4