

quatrième série - tome 44 fascicule 6 novembre-décembre 2011

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Gaëtan CHENEVIER

On the infinite fern of Galois representations of unitary type

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ON THE INFINITE FERN OF GALOIS REPRESENTATIONS OF UNITARY TYPE

BY GAËTAN CHENEVIER

To Julia and Valeria

ABSTRACT. – Let E be a CM number field, p an odd prime totally split in E , and let X be the p -adic analytic space parameterizing the isomorphism classes of 3-dimensional semisimple p -adic representations of $\mathrm{Gal}(\overline{E}/E)$ satisfying a selfduality condition “of type $\mathrm{U}(3)$ ”. We study an analogue of the infinite fern of Gouvêa-Mazur in this context and show that each irreducible component of the Zariski-closure of the modular points in X has dimension at least $3[E : \mathbb{Q}]$. As important steps, and in any rank, we prove that any first order deformation of a generic enough crystalline representation of $\mathrm{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ is a linear combination of trianguline deformations, and that unitary eigenvarieties are étale over weight space at the non-critical classical points. As another application, we give a surjectivity criterion for the localization at p of the *adjoint*^{*} Selmer group of a p -adic Galois representation attached to a cuspidal cohomological automorphic representation of $\mathrm{GL}_n(\mathbb{A}_E)$ of type $\mathrm{U}(n)$ (for any n).

RÉSUMÉ. – Soient E un corps de nombres CM, p un nombre premier impair totalement décomposé dans E , et soit X l’espace analytique p -adique paramétrant les classes d’isomorphie de représentations p -adiques semisimples de dimension 3 de $\mathrm{Gal}(\overline{E}/E)$ satisfaisant une condition d’auto dualité « de type $\mathrm{U}(3)$ ». Nous étudions un analogue de la fougère infinie de Gouvêa-Mazur dans ce contexte et démontrons que l’adhérence Zariski des points modulaires de X a toutes ses composantes irréductibles de dimension au moins $3[E : \mathbb{Q}]$. Au passage, nous prouvons en toute dimension que toute déformation à l’ordre 1 d’une représentation cristalline suffisamment générique de $\mathrm{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ est une combinaison linéaire de déformations triangulines, et que les variétés de Hecke unitaires sont étales sur l’espace des poids aux points classiques non critiques. Enfin, nous obtenons un critère de surjectivité de l’application de localisation en p du groupe de Selmer *adjoint*^{*} d’une représentation galoisienne p -adique attachée à une représentation automorphe cuspidale cohomologique de $\mathrm{GL}_n(\mathbb{A}_E)$ qui est de type $\mathrm{U}(n)$ (pour tout n).

The author is supported by the C.N.R.S.

* Pronounce “adjoint primed Selmer group.”

Introduction

Let E be a number field, p a prime number, S a finite set of places of E containing the places above p and ∞ , $G_{E,S}$ the Galois group of a maximal algebraic extension of E unramified outside S , and let $n \geq 1$ be an integer. We are interested in the set of n -dimensional, semisimple, continuous representations

$$\rho : G_{E,S} \longrightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$$

taken up to isomorphism. This set turns out to be the $\overline{\mathbb{Q}}_p$ -points of a rigid analytic space \mathfrak{X} (or \mathfrak{X}_n) over \mathbb{Q}_p in a natural way⁽¹⁾. An interesting subset

$$\mathfrak{X}^g \subset \mathfrak{X}(\overline{\mathbb{Q}}_p)$$

is the subset of representations which are *geometric*, in the sense they occur as a subquotient of $H_{\mathrm{et}}^i(X_{\overline{E}}, \overline{\mathbb{Q}}_p)(m)$ for some proper smooth algebraic variety X over E and some integers $i \geq 0$ and $m \in \mathbb{Z}$. There are several basic open questions that we can ask about \mathfrak{X} and its locus \mathfrak{X}^g ; here are some of them:

Questions. – Does \mathfrak{X}^g possess some specific structure inside \mathfrak{X} ? What can we say about its various closures in \mathfrak{X} (for example, for the Zariski or the p -adic topologies)? What if we replace \mathfrak{X}^g by its subset \mathfrak{X}^c of ρ 's which are crystalline at the places of E above p ?

Regarding the first question, a trivial observation is that \mathfrak{X}^g is countable, as the set of isomorphism classes of algebraic varieties over E is countable, thus it certainly contains no analytic subset of \mathfrak{X} of positive dimension. In the simplest case $E = \mathbb{Q}$ and $S = \{\infty, p\}$ then \mathfrak{X}_1 is the space of p -adic continuous characters of \mathbb{Z}_p^* (a finite union of 1-dimensional open balls) and \mathfrak{X}^c is the subset of characters of the form $x \mapsto x^m$ for $m \in \mathbb{Z}$, which is Zariski-dense in \mathfrak{X}_1 . For a general E and S , we leave as an exercise to the reader to check that class field theory and the theory of complex multiplication show that \mathfrak{X}^c is also Zariski-dense in \mathfrak{X}_1 assuming Leopoldt's conjecture for E at p .

As a second and much more interesting example, let us recall the discovery of Gouvêa and Mazur [33]. They assume that $d = 2$, $E = \mathbb{Q}$, and say $S = \{\infty, p\}$ to simplify. Let q be a power of an odd prime p and let

$$\bar{\rho} : G_{\mathbb{Q},S} \longrightarrow \mathrm{GL}_2(\mathbb{F}_q)$$

be an absolutely irreducible odd Galois representation. Let $R(\bar{\rho})$ denote the universal odd $G_{\mathbb{Q},S}$ -deformation ring of $\bar{\rho}$ in the sense of Mazur and let $\mathfrak{X}(\bar{\rho})$ be its analytic generic fiber: it is the connected component of \mathfrak{X}_2 parameterizing the ρ with residual representation $\bar{\rho}$. In general $\mathfrak{X}(\bar{\rho})$ is a rather complicated space, and Mazur first studied it in the *unobstructed case*⁽²⁾ $H^2(G_{\mathbb{Q},S}, \mathrm{ad}(\bar{\rho})) = 0$, for which class field theory shows that $R(\bar{\rho}) \simeq \mathbb{Z}_q[[x, y, z]]$, hence $\mathfrak{X}(\bar{\rho})$ is the open unit ball of dimension 3 over \mathbb{Q}_q . In this case, Gouvêa and Mazur showed that \mathfrak{X}^c is Zariski-dense in $\mathfrak{X}(\bar{\rho})$. They actually show that the subset

$$\mathfrak{X}^{\mathrm{mod}} \subset \mathfrak{X}(\bar{\rho})$$

⁽¹⁾ We will not use this space in the sequel, but for its definition see [13, Thm D].

⁽²⁾ The philosophy of special values of L -functions suggests that this unobstructed case is the generic situation. It is now known for instance that for $\bar{\rho} = \bar{\rho}_\Delta$ attached to Ramanujan's $\Delta = \sum_{n>0} \tau(n)q^n$, $p > 13$ and $p \neq 691$, the deformation problem of $\bar{\rho}$ is unobstructed (Mazur, Weston [39]).

of p -adic Galois representations $\rho_f(m)$ attached to an eigenform $f \in S_k(\mathrm{SL}_2(\mathbb{Z}))$ for some weight k , and some $m \in \mathbb{Z}$, is Zariski-dense in $\mathfrak{X}(\bar{\rho})$. This subset is non empty as $\bar{\rho}$ is modular (Khare-Wintenberger).

Their proof relies heavily on the theory of p -adic families of modular eigenforms due to Coleman, extending pioneering works of Hida, that we briefly recall. Let $f = q + a_2q^2 + a_3q^3 + \cdots \in \overline{\mathbb{Q}}_p[[q]]$ be a classical modular eigenform of level 1, of some weight k , and such that $\bar{\rho}_f = \bar{\rho}$; the representation ρ_f corresponds to some $x_f \in \mathfrak{X}(\bar{\rho})(\overline{\mathbb{Q}}_p)$. Attached to f we have two p -Weil numbers of weight $k - 1$ which are the roots of the polynomial $X^2 - a_pX + p^{k-1}$. The main result of Coleman asserts that if φ is one of them, we can attach to (f, φ) an affinoid (equidimensional) subcurve

$$C_{(f, \varphi)} \subset \mathfrak{X}(\bar{\rho})$$

containing x_f as well as a Zariski-dense subset of modular points, and such that for each modular point $x_{f'} \in C_{(f, \varphi)}$ then f' has a p -Weil number with the same p -adic valuation as φ . This curve $C_{(f, \varphi)}$ is not quite canonical but its germ at x_f is. The infinite fern of Gouvêa-Mazur is by definition the union of all the $C_{(f, \varphi)}$ for all f and choice of φ . A simple but important observation made by Gouvêa and Mazur is that in some neighborhood of x_f in $\mathfrak{X}(\bar{\rho})$ the two curves $C_{(f, \varphi)}$ and $C_{(f, \varphi')}$ only intersect at the point x_f if φ and φ' have different valuations: this essentially follows from the previous properties and the fact that the “weight” varies analytically in Coleman’s families. From this “fractal” picture it follows at once that the Zariski-closure of the modular points, or which is the same the Zariski-closure of the fern, has dimension at least 2 inside $\mathfrak{X}(\bar{\rho})$, and a simple argument of Tate-twists using Sen’s theory gives then the result. The story does not quite end here as some years later, Coleman and Mazur defined a wonderful object, *the eigencurve*, which sheds new light on the infinite fern. They define a *refined modular point* as a pair

$$(x, \varphi) \in \mathfrak{X}(\bar{\rho}) \times \mathbb{G}_m$$

where $x = x_f$ is modular and φ is a p -Weil number of f . The interesting fact is that the Zariski-closure of the refined modular points in $\mathfrak{X}(\bar{\rho}) \times \mathbb{G}_m$ is an *equidimensional curve*, the so-called $\bar{\rho}$ -*eigencurve*. Its image in $\mathfrak{X}(\bar{\rho})$ is the *complete infinite fern*, which simultaneously analytically continues each leaf of the infinite fern itself. This picture for $\mathfrak{X}(\bar{\rho})$ provides a rather satisfactory answer towards the first of the main questions above, even though very little is known about the geometry of the eigencurve at present. It is believed that $\mathcal{E}(\bar{\rho})$ has only finitely many irreducible components. An amazing consequence of this conjecture would be that for a well chosen modular point $x \in \mathfrak{X}(\bar{\rho})$ the analytic continuation of a well chosen leaf at x would be Zariski-dense in $\mathfrak{X}(\bar{\rho})$! However, as far as we know there is no non-trivial case in which this conjecture or its variants in other dimensions > 1 is known.

Let us mention that when S is general and $\bar{\rho}$ is possibly obstructed, the approach above of Gouvêa-Mazur still shows that each irreducible component of the Zariski-closure of the modular points in $\mathfrak{X}(\bar{\rho})$ has dimension at least 3. In a somehow opposite direction, it is conjectured that in all cases $\mathfrak{X}(\bar{\rho})$ is equidimensionnal of dimension 3, and that each of its irreducible components contains a smooth modular point. This conjecture, combined with the result of Gouvêa-Mazur, implies the Zariski-density of the modular points in $\mathfrak{X}(\bar{\rho})$ for each odd $\bar{\rho}$ (say absolutely irreducible). Relying on $R = T$ theorems of Taylor-Wiles,

Diamond et al., Boeckle was able to show that conjecture under some rather mild assumption on $\bar{\rho}$, hence the Zariski-density in most cases: we refer to [6] for the precise statements.

Our main aim in this paper is to study a generalization of this picture to the higher dimensional case. Our most complete results will concern some pieces of \mathfrak{X}_3 satisfying some sort of self-duality condition. Let E be a CM field⁽³⁾, and let

$$\bar{\rho} : G_{E,S} \longrightarrow \mathrm{GL}_3(\mathbb{F}_q)$$

be an absolutely irreducible Galois representation such that $\bar{\rho}^* \simeq \bar{\rho}^c$, where c is a generator of $\mathrm{Gal}(E/F)$ and F the maximal totally real subfield of E . Let $\mathfrak{X}(\bar{\rho}) \subset \mathfrak{X}_3$ denote the closed subspace of $x \in \mathfrak{X}_3$ such that $\rho_x^* \simeq \rho_x^c$ and $\bar{\rho}_x \simeq \bar{\rho}$. This $\mathfrak{X}(\bar{\rho})$ has conjectural equidimension $6[F : \mathbb{Q}]$, and under an unobstructedness assumption similar to Mazur's one it is actually an open unit ball over \mathbb{Q}_q in that number of variables. There is a natural notion of modular points in $\mathfrak{X}(\bar{\rho})$: they are the x such that ρ_x is a p -adic Galois representation attached to a cuspidal automorphic representation Π of $\mathrm{GL}_3(\mathbb{A}_E)$ such that Π_∞ is cohomological, $\Pi^\vee \simeq \Pi^c$, and such that for v finite dividing p or outside S , Π_v is unramified. Those Galois representations have been constructed by Rogawski; they are cut out from the étale cohomology of (some abelian varieties over) the Picard modular surfaces and they are related to automorphic forms on unitary groups in 3 variables associated to E/F .

THEOREM A. – *Assume that p is totally split in E . Then each irreducible component of the Zariski-closure of the modular points in $\mathfrak{X}(\bar{\rho})$ has dimension at least $6[F : \mathbb{Q}]$.*

In particular, in the unobstructed case the set of modular points of $\mathfrak{X}(\bar{\rho})$ is Zariski-dense if it is non-empty.

In the appendix, we give several examples of elliptic curves A over \mathbb{Q} such that the deformation problem of type $U(3, E/\mathbb{Q})$ of $\bar{\rho} := \mathrm{Sym}^2 A[p]_{|\mathrm{G}_E}(-1)$ is unobstructed for $p = 5$ and $E = \mathbb{Q}(i)$. As in the work of [6] in the $\mathrm{GL}(2, \mathbb{Q})$ case, we expect that combining Theorem A with suitable $R = T$ theorems (as in [16] and subsequent work), one should be able to remove the unobstructedness assumption under suitable assumptions on $\bar{\rho}$. However, we postpone this to a subsequent study.


As in the work of Gouvêa-Mazur, a very important “constructive” ingredient of our proof is the theory of families of modular forms (for $U(3, E/F)$ here), or better, the related eigenvarieties. They can be quickly defined as follows, from the notion of *refined modular points*. Assume $F = \mathbb{Q}$ for simplicity and fix some prime v of E above p , so $E_v = \mathbb{Q}_p$ by assumption. For each modular point $x \in \mathfrak{X}(\bar{\rho})$, it is known that $\rho_{x,p} := \rho_x|_{\mathrm{G}_{E_v}}$ is crystalline with distinct Hodge-Tate weights, say $k_1(x) < k_2(x) < k_3(x)$. Define a refined modular point as a pair $(x, (\tilde{\varphi}_i(x))) \in \mathfrak{X}(\bar{\rho}) \times \mathbb{G}_m^3$ such that x is a modular point and such that

$$(p^{k_1(x)} \tilde{\varphi}_1(x), p^{k_2(x)} \tilde{\varphi}_2(x), p^{k_3(x)} \tilde{\varphi}_3(x))$$

is an ordering of the eigenvalues of the crystalline Frobenius of $D_{\mathrm{crys}}(\rho_{x,p})$; there are up to 6 ways to refine a given modular point. We define the $\bar{\rho}$ -eigenvariety

$$\mathcal{E}(\bar{\rho}) \subset \mathfrak{X}(\bar{\rho}) \times \mathbb{G}_m^3$$

⁽³⁾ Throughout this paper, a CM field is assumed to be imaginary.



as the Zariski-closure of the refined modular points⁽⁴⁾; the *complete infinite fern of type* $U(3, E/\mathbb{Q})$ is the subset $\mathcal{F}(\bar{\rho}) \subset \mathcal{X}(\bar{\rho})$ image of $\mathcal{E}(\bar{\rho})$ under the first projection. By a former result of the author, this eigenvariety turns out to be equidimensional of dimension 3, and has some additional properties. The analogues of Coleman's arcs through a modular point $x \in \mathcal{X}(\bar{\rho})$ are now 3-dimensional locally closed subspaces $C_{x,\varphi}$ (the “leaves of the fern at x ”) indexed by each refinements (x, φ) of x . The germ of $C_{x,\varphi}$ at x is canonical, and the modular points are actually Zariski-dense in $C_{x,\varphi}$. The $\bar{\rho}$ -infinite fern of type $U(3, E/\mathbb{Q})$ is the union of all the leaves $C_{x,\varphi}$ constructed this way when (x, φ) runs over all the refined modular points: see the figure above.

For dimension reasons, this situation is subtler than the one of Gouvêa-Mazur. Indeed, assume that $\bar{\rho}$ is unobstructed to simplify, so $\mathcal{X}(\bar{\rho})$ is the open unit ball of dimension 6. Remark that there is a one dimensional family of twists by Galois characters of type $U(1)$, and $\mathcal{E}(\bar{\rho})$ is stable by this family of twists as well, so we should rather think of $\mathcal{X}(\bar{\rho})$ as a 5-dimensional space in which evolves the 2-dimensional fern $\mathcal{F}(\bar{\rho})$ (with a Zariski-dense accumulating subset of six times multiple points). The problem is that even if we knew that any two of the six leaves are transversal at each modular point, this would not be enough to exclude the possibility of a 4-dimensional Zariski-closure. The situation is even worse when $[F : \mathbb{Q}] > 1$ as in this case we have to pass from dimension $3[F : \mathbb{Q}]$ to $6[F : \mathbb{Q}]$.

⁽⁴⁾ Let us warn the reader that it is actually not quite the right definition (for instance Theorem D below may not hold with this one), although we shall content ourselves with it in this introduction and refer to Def. 2.2 for the right one.

The idea of our proof is to study the relative positions of the tangent spaces of the local leaves $C_{x,\varphi}$, when φ varies, at a given modular point $x \in \mathfrak{X}(\bar{\rho})$. A key intermediate result is the following:

THEOREM B. – *Assume $\bar{\rho}$ modular. There exist modular points $x \in \mathfrak{X}(\bar{\rho})$ such that $\rho_{x,p}$ is absolutely irreducible and has distinct crystalline Frobenius eigenvalues in $k(x)$. If x is such a point, then the image of the natural map on tangent spaces*

$$\bigoplus_{y \mapsto x, y \in \mathcal{E}(\bar{\rho})} T_y(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho}))$$

has dimension ≥ 6 .

It is not difficult to show that Theorem B actually implies Theorem A (see §2.8). The first part of Theorem B is a simple application of eigenvarieties, but its second part is rather deep. It relies on two general results of independent interest whose proofs form the technical heart of this paper. The idea is to study the image of the map in Theorem B in the tangent space of the deformation space of the local representation $\rho_{x,p}$, on which we will have a sufficiently efficient control as we shall see. As an indication of this, recall that since the works of Kisin, Colmez and Bellaïche-Chenevier, it is known that the restriction at $G_{\mathbb{Q}_p}$ of the family of Galois representations over $\mathcal{E}(\bar{\rho})$ has some very specific properties: they are *trianguline* in the neighborhood of the “non-critical and regular” modular points.

The first important result we prove is a purely local theorem about the deformation space of a given crystalline representation of $G_{\mathbb{Q}_p} = \text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$. Let L be a finite extension of \mathbb{Q}_p and let V be a crystalline representation of $G_{\mathbb{Q}_p}$ of any L -dimension n . Assume that $\text{End}_{G_{\mathbb{Q}_p}}(V) = L$, that V has distinct Hodge-Tate numbers, and that the eigenvalues φ_i of the crystalline Frobenius on $D_{\text{crys}}(V)$ belong to L and satisfy $\varphi_i \varphi_j^{-1} \neq 1, p$ for all $i \neq j$. Let \mathfrak{X}_V be the deformation functor of V to the category of local Artinian L -algebras with residue field L . It is pro-representable and formally smooth of dimension $n^2 + 1$. For each of the $n!$ orderings \mathcal{F} of the φ_i (such an ordering is called again a *refinement* of V , for some obvious reasons), we defined in [1, §2] the \mathcal{F} -trianguline deformation subfunctor $\mathfrak{X}_{V,\mathcal{F}} \subset \mathfrak{X}_V$, whose dimension is $\frac{n(n+1)}{2} + 1$. Roughly speaking, the choice of \mathcal{F} corresponds to a choice of a triangulation of the (φ, Γ) -module of V over the Robba ring, and $\mathfrak{X}_{V,\mathcal{F}}$ parameterizes the deformations such that this triangulation lifts. When the φ -stable complete flag of $D_{\text{crys}}(V)$ defined by \mathcal{F} is in general position compared to the Hodge filtration, we say that \mathcal{F} is *non-critical*.

THEOREM C. – *Assume that n “well-chosen” refinements of V are non-critical (e.g. all of them), or that $n \leq 3$. Then on tangent spaces we have an equality*

$$\mathfrak{X}_V(L[\varepsilon]) = \sum_{\mathcal{F}} \mathfrak{X}_{V,\mathcal{F}}(L[\varepsilon]).$$

In other words, *any first order deformation of a generic crystalline representation is a linear combination of trianguline deformations*. Maybe surprisingly, our proof of this result is by induction on the dimension of V , which requires first to extend the statement to the world of non-étale (φ, Γ) -modules over the Robba ring and work in this general setting (as in [1]). We also have to extend and complete several results of [1] to

general paraboline deformation functors, and even in the trianguline case the proofs given here are actually slightly different from the ones there. Following the ideas of this paper, Theorem C has another purely local application to the Zariski-density of the crystalline points in some components of the representation space of $G_{\mathbb{Q}_p}$ (for instance in the residually irreducible ones), that will however be given elsewhere. Last but not least, let us remark that the codimension of the crystalline locus in \mathfrak{X}_V , namely $\frac{n(n+1)}{2}$, actually coincides with the (conjectural) dimension of $\mathfrak{X}(\bar{\rho})$ if $\bar{\rho}$ is of type $U(n)$ (see below). This numerical coincidence reminds of course the ones discussed in [16] and is fundamental to the application to the Zariski-density of modular points (e.g. in the unobstructed case of Thm. A).

The second key ingredient for the proof of Theorem B is an important theorem about eigenvarieties that we simply state as a slogan in this introduction:

THEOREM D. – *Eigenvarieties are étale over the weight space at non-critically refined, regular, classical points.*

See Theorems 4.8 and 4.10 for precise statements concerning unitary eigenvarieties. In terms of families of p -adic automorphic forms, this theorem means that the natural p -adic family passing through a non-critically refined (and regular) automorphic representation may be parameterized by a polydisc with weight maps for parameters (either automorphic, or Hodge-Tate-Sen weights), which is maybe the most natural statement we may expect in the theory of p -adic families. For ordinary refinements several instances of this slogan have been proved by Hida. Our proof relies in particular on a number of properties of automorphic representations for unitary groups (including some multiplicity one results of Rogawski for $U(3)$ and of Labesse for $U(n)$), on the generalized theta-maps studied by Jones, and on a result of Kisin.

Let us go back to Theorem A. We have dealt with cases of absolutely irreducible $\bar{\rho}$ to fix ideas, but it holds in any residually reducible case, with the same proof, if we work with pseudocharacters instead of representations (using [13] for $p = 3!$). More interestingly, we expect that our approach will lead to a proof of the Zariski-density of the infinite fern of type $U(n)$ for any n . In this case $\mathfrak{X}(\bar{\rho})$ has conjectural dimension $[F : \mathbb{Q}] \frac{n(n+1)}{2}$, $\mathcal{E}(\bar{\rho})$ has dimension $n[F : \mathbb{Q}]$, and there are generically $n!^{[F:\mathbb{Q}]}$ local leaves in $\mathcal{F}(\bar{\rho})$ through each modular point. (When $n = 3$, the coincidence $3! = 3(3 + 1)/2 = 6$ actually plays no specific role.) Although Theorems C and D apply in this generality, we were faced with a new difficulty concerning the existence of sufficiently many global automorphic representations which are *generic enough* at p : we refer to §5.5 for a discussion about it. On the other hand, the degree of the base field is rather insensitive to our method, which allowed us for instance to extend the result of Gouvêa-Mazur to the Hilbert modular case (see Theorem 5.9):

THEOREM E. – *Let F be a totally real field of even degree, p an odd prime totally split in F , \mathfrak{d} the Leopoldt defect of F at p , and let $\bar{\rho} : G_{F,S} \rightarrow \mathrm{GL}_2(\mathbb{F}_q)$ be an absolutely irreducible modular Galois representation.*

Then the irreducible components of the Zariski-closure of the essentially modular points in $\mathfrak{X}(\bar{\rho})$ all have dimension at least $1 + \mathfrak{d} + 2[F : \mathbb{Q}]$. If $H^2(G_{F,S}, \mathrm{ad}(\bar{\rho})) = 0$, then the modular points are Zariski-dense in $\mathfrak{X}(\bar{\rho})$.

We end this introduction by stating an interesting general result about adjoint' Selmer groups that follows from the theory of eigenvarieties and Theorems *C* and *D* above. Let $n \geq 1$ be any integer, let E/F be a CM field, and let Π be a cohomological cuspidal automorphic representation of $\mathrm{GL}_n(\mathbb{A}_E)$ such that $\Pi^\vee \simeq \Pi^c$. Assume that p is totally split in E and that Π is unramified at each place of E above p . Let

$$\rho_\Pi : G_E \longrightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$$

be a p -adic Galois representation attached to Π . It turns out that the adjoint representation $\mathrm{ad}(\rho_\Pi)$ has a natural extension to G_F that we denote by $\mathrm{Ad}'(\rho_\Pi)$, and whose H^1 has a natural interpretation as the tangent space of the deformation functor of ρ_Π of type $U(n)$. This interpretation allows us to define subspaces $H_{\mathcal{G}}^1(F, \mathrm{Ad}'(\rho_\Pi))$ for any choice of refinements of the $\rho_{\Pi|G_v}$ for each place v of F dividing p . In the following statement, we use standard notations from the theory of Selmer groups: f is the Bloch-Kato condition (automatic outside p for $\mathrm{Ad}'(\rho_\Pi)$) and $H_s^1 = H^1/H_f^1$. See also §3.24 for the precise definition of *weakly generic and regular*, let us simply say that a crystalline representation of $G_{\mathbb{Q}_p}$ with sufficiently generic crystalline Frobenius eigenvalues and whose refinements are all non-critical has this property (in particular it is absolutely indecomposable).

THEOREM F. – *Assume that $\rho_{\Pi|G_v}$ is weakly generic and regular for each $v|p$. Then*

$$H^1(F, \mathrm{Ad}'(\rho_\Pi)) \longrightarrow \prod_{v|p} H_s^1(F_v, \mathrm{Ad}'(\rho_\Pi))$$

is surjective. Moreover, $H^1(F, \mathrm{Ad}'(\rho_\Pi)) = \sum_{\mathcal{G}} H_{\mathcal{G}}^1(F, \mathrm{Ad}'(\rho_\Pi))$ and for S big enough we have $\dim H^2(G_{F,S}, \mathrm{Ad}'(\rho_\Pi)) = \dim H_f^1(F, \mathrm{Ad}'(\rho_\Pi))$.

The second part of this theorem is a variant for Selmer groups of Theorem *C*. It is the infinitesimal counterpart of the infinite fern of type $U(n, E/F)$ in the adjoint' Selmer group. It says in particular that $H^1(F, \mathrm{Ad}'(\rho_\Pi))$ is spanned by elements which are not too far from being geometric. As far as we know, this is the first example of such a statement. The first part of the theorem is actually deduced from the second one, and the assertion on H^2 follows from the computation of the sign of ρ_Π in [2]. We believe that a variant of Theorem *F* should hold for a huge class of Selmer groups. Let us mention here that the proof of Theorem *F* also uses some base-change trick to reduce to a situation where the multiplicity one results of Labesse (and Theorem *D*) apply.

The theorem above has a corollary concerning the Galois representations associated to essentially-selfdual cuspidal, cohomological, automorphic representations of GL_n over a totally real field: see Thm. 6.11. Here is a special case. Let Π be a selfdual, cohomological, and cuspidal, automorphic representation of $\mathrm{GL}_n(\mathbb{A}_{\mathbb{Q}})$, and let $\rho_\Pi : G_{\mathbb{Q}} \longrightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$ be an associated p -adic Galois representation, normalized so that $\rho_\Pi^* \simeq \rho_\Pi(n-1)$. Recall that ρ_Π is symplectic if n is even and orthogonal otherwise by [2]. Assume again that Π_p is unramified.

THEOREM G. – *Assume that $\rho_{\Pi|G_{\mathbb{Q}_p}}$ is weakly generic and regular, and set $V = \mathrm{Sym}^2(\rho_\Pi)(n-1)$ if n is even, $V = \Lambda^2(\rho_\Pi)(n-1)$ otherwise. Then the restriction map $H^1(\mathbb{Q}, V) \longrightarrow H_s^1(\mathbb{Q}_p, V)$ is surjective.*

Let $f = q + a_2q^2 + \cdots$ be a classical modular eigenform of weight $k > 1$, level prime to p , and let $\rho_f : G_{\mathbb{Q},S} \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}}_p)$ be an associated p -adic Galois representation.

THEOREM H. – Assume that $\rho_f|_{G_{\mathbb{Q}_p}}$ is absolutely indecomposable and that the ratio of the two roots of the Hecke polynomial $X^2 - a_pX + \varepsilon(p)p^{k-1}$ of f at p is not a root of unity. Set $V_n = \mathrm{Sym}^n(\rho_f) \otimes \det(\rho_f)^{-n/2}$.

For $n = 2$ and $n = 6$, the restriction $H^1(\mathbb{Q}, V_n) \rightarrow H_s^1(\mathbb{Q}_p, V_n)$ is surjective and $\dim H_f^1(\mathbb{Q}, V_n) = \dim H^2(G_{\mathbb{Q},S}, V_n)$.

As the reader may have noticed, all the global statements of this paper concern number fields which are totally split at p . It is clear from our method that this assumption could be removed but we simply did not have the energy to do so. One reason for this is our reliance on the local results of [29] and [1], which are only written at the moment for the base field \mathbb{Q}_p . Another slightly annoying assumption we have is the regularity condition in Theorems D and F. Again, we expect that it could be removed if the technical problems discussed in [1, §4] and occurring in studying “refined families” were solved. For instance, these two improvements together, combined with the recent developments on the Sato-Tate conjecture, would lead to an extension of Theorem H to any $n \equiv 2 \pmod{4}$.

Aknowledgements. – We are grateful to Gebhard Böckle for some stimulating discussions at the origin of this paper. Part of our proof relies on my book with Joël Bellaïche, it is a pleasure to thank him here for the many fruitful exchanges we had the past years. We also thank Laurent Berger and Kevin Buzzard for some useful discussions, Owen Jones for having made his preprint available, Karim Belabas for some discussions about the use of PARI GP, and the referee for his remarks. Some results of this paper were explained by the author in a Cours Peccot at the Collège de France in March-April 2008 [11], it is a pleasure to thank this institution as well as the Fondation Peccot for their support.

1. The universal Galois representation of type $U(n)$

1.1. Galois representations of type $U(n)$ with n odd and their deformations

Let E be a CM field, F its maximal totally real subfield and fix $c \in G_F$ a complex conjugation. Let p be an odd prime number. Let S be a finite set of primes of E containing the primes ramified over F and the primes above p . We assume that $c(S) = S$ and we shall sometimes view S as a set of places of F containing the subset S_∞ of all the Archimedean primes. We fix an odd integer n ; eventually we shall take $n = 3$.

Let A be a commutative ring and let $\rho : G_{E,S} \rightarrow \mathrm{GL}_n(A)$ be an absolutely irreducible representation, in the sense that $A[\rho(G_{E,S})] = M_n(A)$. We say that ρ is of type $U(n)$ if

$$\rho^* \simeq \rho^c$$

as $A[G_{E,S}]$ -modules, or which is the same if $\mathrm{trace} \rho(g^{-1}) = \mathrm{trace} \rho(cgc)$ for all $g \in G_E$. Let $\mathrm{Gal}(E/F)$ act on GL_n by $c(g) := {}^t g^{-1}$ and view $\mathcal{G}_n := \mathrm{GL}_n \rtimes \mathrm{Gal}(E/F)$ as a group scheme over \mathbb{Z} . The terminology ρ is of type $U(n)$ comes from the fact that \mathcal{G}_n may be viewed as the reduced Langlands dual group of the unitary groups in n variables associated to E/F (recall n is odd) and the following lemma, which is a variant of [16, §1].

LEMMA 1.2. – ρ is of type $U(n)$ if, and only if, there exists a representation

$$\tilde{\rho} : G_{F,S} \longrightarrow GL_n(A) \rtimes \text{Gal}(E/F)$$

where the induced map $G_{F,S} \rightarrow \text{Gal}(E/F)$ is the natural map and $\tilde{\rho}|_{G_{E,S}} = \rho$.

Indeed, $\tilde{\rho}(c) := Pc$, with $P \in GL_n(A)$, defines an extension of ρ as in the statement if, and only if, ${}^tP = P$ and ${}^t\rho(g)^{-1} = P^{-1}\rho(cgc)P$ for all $g \in G_{E,S}$. This last condition actually implies ${}^tP = \mu P$ for some $\mu \in A^*$ as ρ is absolutely irreducible, and $\mu = 1$ as n is odd. Note that the extension $\tilde{\rho}$ is not unique, but $\tilde{\rho}'$ is another one if, and only if, $\tilde{\rho}'(c) = P'c$ with $P' = \lambda P$ and $\lambda \in A^*$.

Let q be a power of p and let us fix from now on

$$\bar{\rho} : G_{E,S} \longrightarrow GL_n(\mathbb{F}_q)$$

an absolutely irreducible, continuous, representation of type $U(n)$. Let \mathbb{Z}_q be the Witt vectors of \mathbb{F}_q and let \mathcal{C} be the category of Artinian local \mathbb{Z}_q -algebras with residue field $A/m_A = \mathbb{F}_q$. Consider the deformation functor

$$\mathcal{D} : \mathcal{C} \rightarrow \text{Sets}$$

defined as follows: for any object A of \mathcal{C} , $\mathcal{D}(A)$ is the set of A -isomorphism classes of continuous representations $\rho_A : G_{E,S} \longrightarrow GL_n(A)$ of type $U(n)$ such that $\rho_A \otimes_A \mathbb{F}_q$ is isomorphic to $\bar{\rho}$. As $\bar{\rho}$ is absolutely irreducible, this functor is pro-representable by a complete local noetherian \mathbb{Z}_q -algebra $R(\bar{\rho})$ with residue field \mathbb{F}_q . Indeed, it is the quotient of Mazur's universal $G_{E,S}$ -deformation ring $R'(\bar{\rho})$ by the ideal generated by the elements $\text{trace } \rho^u(g^{-1}) - \text{trace } \rho^u(cgc)$, for all $g \in G_{E,S}$, where ρ^u is Mazur's universal $R'(\bar{\rho})$ -valued deformation. (Note that the argument so far did not use the assumption p odd.)

Using class field theory and arguments of Mazur we will see below that the Krull dimension of $R(\bar{\rho})/(p)$ is always $\geq [F : \mathbb{Q}] \frac{n(n+1)}{2}$. The precise structure of $R(\bar{\rho})$ is presumably extremely complicated in general, however we shall not be interested in these kinds of complications in this paper.

Let $\mathfrak{g}_n \simeq M_n(\mathbb{F}_q)$ be the Lie algebra of \mathcal{G}_n over \mathbb{F}_q viewed as a representation of $\mathcal{G}_n(\mathbb{F}_q)$. Fix once and for all some $\tilde{\rho}$ lifting $\bar{\rho}$ as in Lemma 1.2. Thanks to this choice we may view \mathfrak{g}_n as a $G_{F,S}$ -module, that we shall denote by $\text{Ad}'(\bar{\rho})$. This latter module does not depend on the choice of $\tilde{\rho}$. By definition, $\text{Ad}'(\bar{\rho})|_{G_E}$ is the usual adjoint representation of $\bar{\rho}$. Moreover, if $\tilde{\rho}(c) = Pc$, then c acts on $\text{Ad}'(\bar{\rho}) = M_n(\mathbb{F}_q)$ as⁽⁵⁾ $X \mapsto -P^tXP^{-1}$. In particular, $H^0(G_{F,S}, \text{Ad}'(\bar{\rho})) = 0$ as $\bar{\rho}$ is absolutely irreducible and p is odd.

DEFINITION 1.3. – We say that the $U(n)$ -deformation theory of $\bar{\rho}$ is unobstructed, or for short that $\bar{\rho}$ is unobstructed, if $H^2(G_{F,S}, \text{Ad}'(\bar{\rho})) = 0$.

By Tate's global and local duality theorems, $\bar{\rho}$ is unobstructed if, and only if,

$$\text{III}_S^1(G_{F,S}, \text{Ad}'(\bar{\rho})(1)) = 0 \quad \text{and} \quad H^0(F_v, \text{Ad}'(\bar{\rho})(1)) = 0 \quad \text{for all finite prime } v \in S$$

(note that \mathfrak{g}_n is a selfdual $\mathbb{F}_q[\mathcal{G}_n(\mathbb{F}_q)]$ -module and see [34, Rem. 5.2.(c)]).

⁽⁵⁾ As an exercise, the reader can check that up to conjugating $\bar{\rho}$ if necessary, we may choose a $\tilde{\rho}$ such that $\tilde{\rho}(c) = c$, i.e. $P = \text{id}$ (use that n and p are odd).

PROPOSITION 1.4. – *If $\bar{\rho}$ is unobstructed, then $R(\bar{\rho})$ is formally smooth over \mathbb{Z}_q of relative dimension*

$$[F : \mathbb{Q}] \frac{n(n+1)}{2}.$$

Set $V = \text{Ad}'(\bar{\rho})$. Tate's global Euler characteristic formula shows that

$$\dim H^1(G_{F,S}, V) = [F : \mathbb{Q}] \dim V^{\tilde{\rho}(c)=-1} + \dim H^0(G_{F,S}, V) + \dim H^2(G_{F,S}, V)$$

(dimensions are over \mathbb{F}_q and recall that p is odd). But $X \mapsto XP$ identifies $M_n(\mathbb{F}_q)^{\tilde{\rho}(c)=-1}$ with the subspace of symmetric matrices in $M_n(\mathbb{F}_q)$, which has dimension $\frac{n(n+1)}{2}$, so $\dim H^1(G_{F,S}, \text{Ad}'(\bar{\rho})) = [F : \mathbb{Q}] \frac{n(n+1)}{2}$ if $\bar{\rho}$ is unobstructed.

The proposition follows from this computation and from the following general lemma. If A is an object of \mathcal{C} , two group homomorphisms with target $\mathcal{G}_n(A)$ are said *equivalent* if they are conjugate by an element in $\text{Ker}(\mathcal{G}_n(A) \rightarrow \mathcal{G}_n(\mathbb{F}_q))$. Define another functor $\mathcal{D}' : \mathcal{C} \rightarrow \text{Sets}$ as follows: for an object A of \mathcal{C} , $\mathcal{D}'(A)$ is the set of equivalence classes of continuous homomorphisms $r_A : G_{F,S} \rightarrow \mathcal{G}_n(A)$ such that $r_A \otimes_A \mathbb{F}_q = \tilde{\rho}$ (which has been fixed above).

LEMMA 1.5. – *The map $r_A \mapsto r_{A|G_E}$ induces an isomorphism $\mathcal{D}' \xrightarrow{\sim} \mathcal{D}$. Moreover, $\mathcal{D}(\mathbb{F}_q[\varepsilon]) = H^1(G_{F,S}, \text{Ad}'(\bar{\rho}))$ and if $H^2(G_{F,S}, \text{Ad}'(\bar{\rho})) = 0$, then \mathcal{D} is formally smooth over \mathbb{Z}_q .*

Proof. – For each A , $\mathcal{D}'(A) \rightarrow \mathcal{D}(A)$ is surjective by the proof of Lemma 1.2. Let $r_1, r_2 \in \mathcal{D}'(A)$ be such that $r_1|_{G_E} \simeq r_2|_{G_E}$; up to equivalence we may assume that $r_1|_{G_E} = r_2|_{G_E}$. Set $r_i(c) = P_i c$, where $P_i \in \text{GL}_n(A)$. Then both P_1 and P_2 intertwine the conjugate and dual of $r_1|_{G_E}$, so $P_1 = \lambda P_2$ for some $\lambda \in A^*$. As each r_i lifts $\tilde{\rho}$, $\lambda \in 1 + m_A$. As p is odd, $\lambda = \mu^2$ for some μ in $1 + m_A$, so $\mu^{-1} r_1 \mu$ is equivalent to r_1 and coincides with r_2 . This concludes the proof of the first part of the statement, the other assertions follow from this and standard facts from group cohomology. \square

Set $\mathbb{Q}_q = \mathbb{Z}_q[1/p]$. If R is a local complete noetherian \mathbb{Z}_q -algebra, the functor $\text{Hom}_{\mathbb{Z}_q}(R, -)$ (continuous \mathbb{Z}_q -algebra morphisms) from the category of affinoid \mathbb{Q}_q -algebras to Sets is representable by a rigid analytic space⁽⁶⁾ over \mathbb{Q}_q called Berthelot's generic fiber of R (see [27, §7] for the basics on this construction). If $R = \mathbb{Z}_q[[T_1, \dots, T_r]]/I$, this space is simply the closed subspace defined by $I = 0$ in the r -dimensional open unit ball $|T_i| < 1$ ($i = 1, \dots, r$) over \mathbb{Q}_q . In the sequel, we will be interested in Berthelot's generic fiber of $R(\bar{\rho})$, that we shall denote by

$$\mathfrak{X}(\bar{\rho}).$$

For example, in the unobstructed case $\mathfrak{X}(\bar{\rho})$ is isomorphic to the open unit ball over \mathbb{Q}_q of dimension $[F : \mathbb{Q}] \frac{n(n+1)}{2}$. Before stating the universal property of $\mathfrak{X}(\bar{\rho})$ we need to set some notations and review some facts concerning families of representations.

Let Y be an affinoid over \mathbb{Q}_q and $\rho : G_{E,S} \rightarrow \text{GL}_n(\mathcal{O}(Y))$ a continuous representation. For $y \in Y$ a point, with residue field $k(y)$, we denote by $\rho_y : G_{E,S} \rightarrow \text{GL}_n(k(y))$ the evaluation of ρ at y and by $\bar{\rho}_y : G_{E,S} \rightarrow \text{GL}_n(k_y)$ the semi-simplification of its residual

⁽⁶⁾ We use the foundations of rigid spaces in [7].

representation. Here k_y denotes the residue field of $k(y)$; it comes with a natural morphism $\mathbb{F}_q \rightarrow k_y$. We say that ρ is a lift of $\bar{\rho}$ if $\bar{\rho}_y \simeq \bar{\rho} \otimes_{\mathbb{F}_q} k_y$ for all $y \in Y$.

Let \mathcal{D}^{an} be the (contravariant) functor from the category of \mathbb{Q}_q -affinoids to Sets where $\mathcal{D}^{\text{an}}(Y)$ is the set of $\mathcal{O}(Y)$ -isomorphism classes of continuous homomorphisms $\rho_Y : G_{E,S} \rightarrow \text{GL}_n(\mathcal{O}(Y))$ which are of type $\text{U}(n)$ and lift $\bar{\rho}$. As $\bar{\rho}$ is absolutely irreducible, $\rho_Y \mapsto \text{trace } \rho_Y$ induces a bijection between $\mathcal{D}^{\text{an}}(Y)$ and the set of continuous pseudocharacters $T_Y : G_{E,S} \rightarrow \mathcal{O}(Y)$ lifting trace $\bar{\rho}$ (for the obvious definition) and of type $\text{U}(n)$, that is such that $T_Y(g^{-1}) = T_Y(cgc)$ for all $g \in G_{E,S}$. The following fact is probably well-known, it follows for instance from [13, Thm. 3.16], but this is much simpler here as $\bar{\rho}$ is absolutely irreducible.

LEMMA 1.6. – $\mathfrak{X}(\bar{\rho})$ represents \mathcal{D}^{an} .

This space $\mathfrak{X}(\bar{\rho})$ is actually a connected component of the $\text{U}(n)$ -type locus of the space \mathfrak{X}_n mentioned in the introduction.

1.7. Modular Galois representations of type $\text{U}(n)$ and examples

We keep the assumptions of §1.1. The main source of Galois representations of type $\text{U}(n)$ is the degree $n - 1$ étale cohomology of the quotients of the complex open unit ball $\sum_{i=1}^{n-1} |z_i|^2 < 1$ in \mathbb{C}^{n-1} by the arithmetic congruence subgroups of $\text{U}(n - 1, 1)(\mathbb{R})$ attached to E/F . When $n = 3$ these are also called Picard modular surfaces. Thanks to the advances in the theory of automorphic forms we may describe them using $\text{GL}(n)$.

Let Π be a cuspidal automorphic representation of $\text{GL}_n(\mathbb{A}_E)$ such that:

- (P1) $\Pi^\vee \simeq \Pi^c$,
- (P2) Π_v is cohomological for each Archimedean place v of E ,
- (P3) Π_v is unramified whenever $v \notin S$ or v divides p .

Fix once and for all a pair $\iota = (\iota_p, \iota_\infty)$ of field embeddings $\iota_p : \bar{\mathbb{Q}} \rightarrow \bar{\mathbb{Q}}_p$ and $\iota_\infty : \bar{\mathbb{Q}} \rightarrow \mathbb{C}$. Class-field theory for $n = 1$, work of Rogawski for $n = 3$, and recent works of Shin [38] and of the participants of the book project of the Paris GRFA seminar for any n [15, 24], attach to such a Π and ι a continuous semi-simple representation

$$\rho_\Pi : G_{E,S} \rightarrow \text{GL}_n(\bar{\mathbb{Q}}_p)$$

of type $\text{U}(n)$ which is compatible with the Frobenius semi-simplified local Langlands correspondence at all primes v not dividing p and which is crystalline at each prime v of E above p . In particular, this representation is geometric in the sense of Fontaine-Mazur. Actually, infinitely many suitable real quadratic base changes of it are even geometric in the sense of the introduction by construction (all when $[F : \mathbb{Q}] > 1$). It is known that ρ_Π is irreducible when $n = 3$, and conjectured in general. Fix a ring homomorphism $\bar{\mathbb{Z}}_p \rightarrow \bar{\mathbb{F}}_p$. Denote by $\bar{\rho}_\Pi : G_{E,S} \rightarrow \text{GL}_n(\bar{\mathbb{F}}_p)$ the isomorphism class of the semi-simplification of ρ_Π .

DEFINITION 1.8. – A representation $\rho : G_{E,S} \rightarrow \text{GL}_n(\bar{\mathbb{Q}}_p)$ is modular of type $\text{U}(n)$ if there exists a Π satisfying (P1) to (P3) as above such that $\rho \simeq \rho_\Pi$. A representation $\bar{\rho} : G_{E,S} \rightarrow \text{GL}_n(\bar{\mathbb{F}}_p)$ is modular of type $\text{U}(n)$ if there exists a Π as above such that $\bar{\rho} \simeq \bar{\rho}_\Pi$.

For short, we shall often say *modular* for *modular of type $\text{U}(n)$* . It is tempting to believe that a variant of Serre's conjecture holds in this context:

Is any continuous absolutely irreducible $\bar{\rho}$ of type $U(n)$ modular?

When $n = 3$, by Rogawski's construction, any modular $\bar{\rho}$ occurs in $H_{\text{et}}^2(S_{\bar{E}}, \bar{\mathbb{F}}_p(1))$ where S is a suitable Picard modular surface over E . A delicate aspect of this variant of Serre's conjecture is that such an H^2 might contain some classes⁽⁷⁾ that do not lift to cohomology classes in characteristic 0: this happens for instance for Mumford's fake projective planes. Nevertheless, the recent advances on $R = T$ theorems and potential modularity in this context (e.g. by Clozel, Harris and Taylor and their co-workers) suggest a positive answer.

EXAMPLE 1.9. – If f is a classical modular eigenform then $\text{Symm}^2(\rho_f) \otimes \det(\rho_f)|_{G_{E,S}}^{-1}$ is of type $U(3)$. When F/\mathbb{Q} is Galois and solvable, it is modular by results of Arthur-Clozel (if $F \neq \mathbb{Q}$), Gelbart-Jacquet and Rogawski. By the modularity theorem, this applies in particular to $\text{Symm}^2 A[p]^*(1)$ for any elliptic curve A over \mathbb{Q} .

1.10. Statement of the theorem

Let $\bar{\rho}$ be a Galois representation of type $U(n)$ and $\mathfrak{X}(\bar{\rho})$ the generic fiber of its universal deformation ring of type $U(n)$. A point $x \in \mathfrak{X}(\bar{\rho})$ is called *modular* if the Galois representation $\rho_x : G_{E,S} \rightarrow \text{GL}_n(k(x))$ attached to x is modular. Let

$$\mathfrak{X}^{\text{mod}} \subset \mathfrak{X}(\bar{\rho})$$

be the (countable) subset of modular points. By definition, $\bar{\rho}$ is modular if, and only if, $\mathfrak{X}^{\text{mod}} \neq \emptyset$. We can now state the main result of this paper.

THEOREM 1.11. – *Assume that $\bar{\rho}$ is modular of type $U(3)$, unobstructed, and that p splits in E . Then $\mathfrak{X}^{\text{mod}}$ is Zariski-dense in $\mathfrak{X}(\bar{\rho})$.*

A subset of points Z of a rigid space Y is called *Zariski-dense* if the only reduced closed subspace of Y containing Z is Y_{red} . When Y is quasi-Stein (which is the case of the $\mathfrak{X}(\bar{\rho})$), it is equivalent to ask that any global function $f \in \mathcal{O}(Y)$ vanishing at all points of Z is locally nilpotent. In the context of the theorem, $\mathfrak{X}(\bar{\rho})$ is the open unit ball of dimension $d = [F : \mathbb{Q}] \frac{n(n+1)}{2}$ over \mathbb{Q}_q , so the statement simply means that any power series $f \in \mathbb{C}_p[[t_1, \dots, t_d]]$ converging on the whole open ball $|t_i| < 1$ and vanishing on $\mathfrak{X}^{\text{mod}}$ is identically 0.

EXAMPLE 1.12. – Set $E = \mathbb{Q}(i)$, $p = 5$, $S = \{\infty, 2, 5, 17\}$ and let A be the elliptic curve $y^2 + xy + y = x^3 - x^2 - x$ over \mathbb{Q} , which is isogenous to the Jacobian of $X_0(17)$. Then $\bar{\rho} = \text{Symm}^2(A[5])(-1)|_{G_E}$ is modular of type $U(3)$ and unobstructed (see the appendix).

COROLLARY 1.13. – *Under the same assumptions, for any object A of \mathcal{C} and any lift $\rho_A \in \mathcal{D}(A)$ of $\bar{\rho}$, there is a finite extension L of \mathbb{Q}_q and a set $\rho_1, \rho_2, \dots, \rho_r$ of modular Galois representations $G_{E,S} \rightarrow \text{GL}_3(\mathcal{O}_L)$ of type $U(3)$ such that the $\mathbb{Z}_q[G_{E,S}]$ -module ρ_A is a subquotient of $\bigoplus_{i=1}^r \rho_i$.*

⁽⁷⁾ I do not know any example of such a class that generates an irreducible Galois representation of dimension 3.

Indeed, this follows from the theorem and the following general fact due to Chevalley: if A is a complete local noetherian ring, and if $\{a_i, i \in I\}$ is a family of proper, cofinite length, ideals of A such that the intersection of all the a_i is zero, then the topology defined by the a_i is the adic topology of the maximal ideal of A .

We will actually prove the following result, of which Theorem 1.11 follows at once by Prop. 1.4.

THEOREM 1.14. – *Assume that p splits in E . Then the irreducible components of the Zariski-closure of $\mathfrak{X}^{\text{mod}}$ inside $\mathfrak{X}(\bar{\rho})$ all have dimension at least $6[F : \mathbb{Q}]$.*

We are led to the following optimistic conjecture (for any integer n).

CONJECTURE 1.15. – *Let $\bar{\rho}$ be any Galois representation of type $U(n)$, then $\mathfrak{X}^{\text{mod}}$ is Zariski-dense in $\mathfrak{X}(\bar{\rho})$.*

2. The infinite fern of Galois representations of type $U(3)$

The main ingredient in the proof of Thm. 1.14 is the so-called infinite fern inside $\mathfrak{X}(\bar{\rho})$ and its study. We use the notations of §1.1 and we assume furthermore that $\bar{\rho}$ is a modular Galois representation of type $U(3)$ and that the odd prime p splits in E . We denote by S_p the set of places of F dividing v , $S_0 = S \setminus (S_\infty \cup S_p)$, and we fix once and for all, for each $v \in S_p$, a place \tilde{v} of E above p , so $\mathbb{Q}_p = F_v = E_{\tilde{v}}$. Set $I = \{1, 2, 3\}$.

2.1. The eigenvariety and the complete infinite fern

Let \mathcal{W} and \mathcal{W}_0 be the rigid analytic spaces over \mathbb{Q}_p parameterizing respectively the p -adic continuous characters of \mathbb{Q}_p^* and \mathbb{Z}_p^* . So \mathcal{W}_0 is a finite disjoint union of 1-dimensional open unit balls over \mathbb{Q}_p and $\mathcal{W} \simeq \mathbb{G}_m \times \mathcal{W}_0$. Consider the subset

$$\mathcal{Z} \subset \mathfrak{X}(\bar{\rho})(\overline{\mathbb{Q}}_p) \times \mathcal{W}(\overline{\mathbb{Q}}_p)^{I \times S_p}$$

of pairs (ρ_Π, δ) satisfying (i) and (ii) below:

(i) ρ_Π is the modular Galois representation of type $U(3)$ associated to Π .

In particular $\rho_{\Pi,v} := (\rho_\Pi)|_{G_{E_{\tilde{v}}}}$ is crystalline with distinct Hodge-Tate weights for each $v \in S_p$. We denote by $k_{1,v} < k_{2,v} < k_{3,v}$ these weights by increasing order, with the convention that the cyclotomic character has Hodge-Tate weight -1 .

(ii) $\delta = (\delta_{i,v})$ where for each $(i, v) \in I \times S_p$, the character $\delta_{i,v} : \mathbb{Q}_p^* \rightarrow \overline{\mathbb{Q}}_p^*$ is the product of $x \mapsto x^{-k_{i,v}}$ and of the character sending \mathbb{Z}_p^* to 1 and p to $\varphi_{i,v}$, where $(\varphi_{i,v})_{i \in I}$ is an ordering of the eigenvalues of the crystalline Frobenius of $D_{\text{crys}}(\rho_{\Pi,v})$ (recall that $E_{\tilde{v}} = \mathbb{Q}_p$).

Each such pair is called a *refined modular point*. There are up to $6^{[F:\mathbb{Q}]}$ ways to refine a given modular point, corresponding to the number of ways to choose an ordering of the eigenvalues of the crystalline Frobenius of $D_{\text{crys}}(\rho_{\Pi,v})$ for each $v \in S_p$.

DEFINITION 2.2. – *The eigenvariety of type $U(3)$ of $\bar{\rho}$ is the Zariski-closure*

$$\mathcal{E}(\bar{\rho}) \subset \mathfrak{X}(\bar{\rho}) \times \mathcal{W}^{I \times S_p}$$

of the subset \mathcal{Z} inside $\mathfrak{X}(\bar{\rho}) \times \mathcal{W}^{I \times S_p}$. The complete infinite fern of type $U(3)$ of $\bar{\rho}$

$$\mathcal{F}(\bar{\rho}) \subset \mathfrak{X}(\bar{\rho})$$

is the set theoretic image of $\mathcal{E}(\bar{\rho})$ via the first projection $\mathrm{pr}_1 : \mathfrak{X}(\bar{\rho}) \times \mathcal{W}^{I \times S_p} \rightarrow \mathfrak{X}(\bar{\rho})$.

By definition $\mathcal{E}(\bar{\rho})$ is a reduced analytic space. The *weight space* is the space $\mathcal{W}_0^{I \times S_p}$. The natural map $\mathcal{W} \rightarrow \mathcal{W}_0$ and the second projection induce a natural map

$$\kappa : \mathcal{E}(\bar{\rho}) \rightarrow \mathcal{W}_0^{I \times S_p}.$$

This map turns out to refine the Hodge-Tate-Sen map. Indeed, recall that the work of Sen defines, for each $v \in S_p$, a monic polynomial

$$P_{\mathrm{Sen},v}(t) \in \mathcal{O}(\mathfrak{X}(\bar{\rho}))[t]$$

of degree 3 whose evaluation at any $x \in \mathfrak{X}(\bar{\rho})$ is the usual Sen polynomial of $\rho_{x,v}$ (the roots of which are the generalized Hodge-Tate weights of $\rho_{x,v}$). The coefficients of this collection of polynomials give rise to a natural map $\kappa_{\mathrm{HT}} : \mathfrak{X}(\bar{\rho}) \rightarrow \mathbb{A}^{I \times S_p}$. The morphism $\beta : \mathcal{W}_0^{I \times S_p} \rightarrow \mathbb{A}^{I \times S_p}$ given by the formula $\beta((\delta_{i,v}))_v = \prod_i (t + (\frac{\partial \delta_{i,v}}{\partial x})_{|x=1})$ is the composite of a finite covering of $\mathbb{A}^{I \times S_p}$ of degree $6^{[F:\mathbb{Q}]}$, étale over the locus parametrizing polynomials with nonzero discriminant, with an étale morphism of infinite degree (essentially a p -adic logarithm). By definition, we have $\kappa_{\mathrm{HT}} \circ \mathrm{pr}_1 = \beta \circ \kappa$ on $\mathcal{E}(\bar{\rho})$.

THEOREM 2.3. – *$\mathcal{E}(\bar{\rho})$ is equidimensional of dimension $3[F:\mathbb{Q}]$. The map κ is locally finite: $\mathcal{E}(\bar{\rho})$ is admissibly covered by the open affinoids U such that $\kappa(U) \subset \mathcal{W}_0^{I \times S_p}$ is open affinoid and $\kappa|_U : U \rightarrow \kappa(U)$ is finite.*

This follows from the main theorem of the theory of p -adic families of automorphic forms for the definite unitary group $U(3)$ (see [10],[12]), and from the base change results of Rogawski [35]. This point of view on eigenvarieties is a generalization of the one of Coleman and Mazur in [18]. Concretely, the equations defining $\mathcal{E}(\bar{\rho})$ inside $\mathfrak{X}(\bar{\rho}) \times \mathcal{W}^{I \times S_p}$ are given by characteristic power series of certain compact Hecke operators acting on the flat family of Banach spaces of p -adic automorphic forms for the definite unitary group $U(3)$. Let us explain now how to deduce this statement from other ones in the literature. Our method is actually a bit different from the one in [18].

Proof. – First, consider the unitary group U/F in three variables attached to the positive definite hermitian norm $(z_1, z_2, z_3) \mapsto \sum_{i=1}^3 \mathrm{Norm}_{E/F}(z_i)$ on E^3 . This group is necessarily quasi-split at all finite places of F , as we are in odd dimension. The work of Rogawski defines by base change a bijection between automorphic representations π of $\mathrm{GL}_3(\mathbb{A}_E)$ satisfying (P1) and (P2) and stable tempered L -packets Π of U , which is compatible with a local base change (that Rogawski also defines) at all places ([35, §13]). The following lemma follows for instance from Rogawski's classification [35, p. 174, §13.1].

LEMMA 2.4. – *For any finite union of Bernstein component \mathcal{B}' of $\mathrm{GL}_3(\mathbb{A}_{E,S_0})$, there is a finite union \mathcal{B} of Bernstein components of $U(\mathbb{A}_{F,S_0})$ such that for each irreducible π' in \mathcal{B}' and any irreducible π of $U(\mathbb{A}_{F,S_0})$ whose base-change is π' , we have $\pi \in \mathcal{B}$.*

As $\mathfrak{X}(\bar{\rho})$ has finitely many irreducible components, and by [1, Lemma 7.8.17], there is a finite union of Bernstein components \mathcal{B}' of $\mathrm{GL}_3(\mathbb{A}_{E,S_0})$ such that for each Π with $\rho_\Pi \in \mathfrak{X}(\bar{\rho})$ we have $\Pi_{S_0} \in \mathcal{B}'$. Choose an associated \mathcal{B} as in the above lemma. Up to enlarging \mathcal{B} if necessary, we may find a compact open subgroup $K \subset U(\mathbb{A}_{F,S_0})$ cutting exactly the union of components \mathcal{B} , in the sense that an irreducible representation τ of $U(\mathbb{A}_{F,S_0})$ belongs to \mathcal{B} if, and only if, $\tau^K \neq 0$.

Let X/\mathbb{Q}_q be the p -adic eigenvariety of U associated to (ι_p, ι_∞) , to the tame level K (spherical outside S), to the set of all places S_p above p , and with respect to the spherical Hecke algebra \mathcal{H} outside S (see [12, Thm. 1.6]). By Rogawski's results and [1, Cor. 7.5.4] (or [10, Cor. 7.7.1]), this eigenvariety carries a continuous pseudocharacter $T : G_{E,S} \rightarrow \mathcal{O}(X)$ such that for each refined classical point x associated to some Π satisfying (P1)–(P3) the evaluation of T at x is $\mathrm{trace}(\rho_\Pi)$. In turn, for each $x \in X$ we have a natural associated semi-simple Galois representation $\rho_x : G_{E,S} \rightarrow \mathrm{GL}_3(\overline{k(x)})$, whose residual semi-simple representation will be denoted by $\bar{\rho}_x : G_{E,S} \rightarrow \mathrm{GL}_3(\bar{k}_x)$, where k_x is the residue field of $k(x)$ (a \mathbb{F}_q -algebra, as $k(x)$ is a \mathbb{Q}_q -algebra). The locus $X(\bar{\rho}) \subset X$ of x such that $\bar{\rho}_x \simeq \bar{\rho}$ is an admissible closed and open subspace of X (this follows for instance from the Brauer-Nesbitt theorem and [13, Lemma 3.9]). By definition of \mathcal{B} and the properties of $X(\bar{\rho})$, there is a natural injection $j : Z \rightarrow X(\bar{\rho})(\overline{\mathbb{Q}_p})$, and $j(Z)$ is a Zariski-dense subset of $X(\bar{\rho})$. The universal property of $\mathfrak{X}(\bar{\rho})$ defines a canonical analytic map $\phi : X(\bar{\rho}) \rightarrow \mathfrak{X}(\bar{\rho})$. The eigenvariety X is also equipped with a finite analytic map $\nu : X \rightarrow \mathcal{W}^{I \times S_p}$. By definition we have $(\phi \times \nu) \circ j = \mathrm{id}_Z$ and property (i) of [12, Thm 1.6] ensures that

$$\phi \times \nu : X(\bar{\rho}) \rightarrow \mathfrak{X}(\bar{\rho}) \times \mathcal{W}^{I \times S_p}$$

is a closed immersion (it is finite because ν is finite). As a consequence, $\phi \times \nu$ induces an isomorphism $X(\bar{\rho}) \xrightarrow{\sim} \mathcal{E}(\bar{\rho})$, and the last statement follows from the properties of $X(\bar{\rho})$. \square

REMARK 2.5. – The proof above is pseudo-character theoretic, hence extends verbatim to the case of any semi-simple $\bar{\rho}$. Moreover, this theorem would also hold for any odd n by the same argument, if we appeal instead of Rogawski's work to the recent works of Mœglin (definition of the tempered L -packets for quasi-split unitary groups) and Labesse (base change to $\mathrm{GL}(n)$) instead of the work of Rogawski, as long as $[F : \mathbb{Q}] \geq 2$ (assumption occurring in Labesse's base-change at the moment).

An important property of the eigenvariety is the so-called *classicity criterion*. Say that $(\delta_{i,v}) \in \mathcal{W}_0$ is *algebraic* if for each $(i, v) \in I \times S_p$ there exists an integer $k_{i,v}$ such that $\delta_{i,v}$ is the character $u \mapsto u^{-k_{i,v}}$. Say that $(x, \delta) \in \mathcal{E}(\bar{\rho})$ is of *algebraic weight* if $\kappa(x)$ is algebraic. In this case, we denote by $k_{i,v}(x)$ the $k_{i,v}$ above and set $\varphi_{i,v}(x) := p^{k_{i,v}(x)} \delta_{i,v}(p)$. The next proposition is [12, Thm. 1.6.(vi)].

PROPOSITION 2.6. – *Let $x \in \mathcal{E}(\bar{\rho})$ be of algebraic weight. Assume that $\forall v \in S_p$:*

- (i) $k_{1,v}(x) < k_{2,v}(x) < k_{3,v}(x)$,
- (ii) $\mathbf{v}(\varphi_{1,v}(x)) < k_{2,v}(x) < \mathbf{v}(\varphi_{3,v}(x))$,

(iii) $\forall i, j \in I, \varphi_{i,v}(x)\varphi_{j,v}(x)^{-1} \neq p$.

Then $x \in \mathcal{Z}$.

In particular, \mathcal{Z} is an accumulation subset of $\mathcal{E}(\bar{\rho})$. More precisely, if $U \subset \mathcal{E}(\bar{\rho})$ is an open affinoid as in the statement of Thm. 2.3, then \mathcal{Z} is Zariski-dense in U if and only if each connected component of $\kappa(U)$ contains an algebraic weight.

Recall that if A and B are two subsets (of points) of a rigid analytic space, we say that A *accumulates at* B if for each $b \in B$ and for each affinoid neighborhood U of b , there exists an affinoid neighborhood $V \subset U$ of b such that $A \cap V$ is Zariski-dense in V . An *accumulation subset* is a subset accumulating at itself. For instance the positive integer \mathbb{N} is an accumulation subset of the closed unit disc over \mathbb{Q}_p , and the algebraic weights form an accumulation subset of $\mathcal{W}_0^{I \times S_p}$.

2.7. The local leaves and the infinite fern

Let $x = \rho_{\Pi} \in \mathfrak{X}(\bar{\rho})$ be a modular point, and choose δ so that $(x, \delta) \in \mathcal{Z}$. By Theorem 2.3 and Prop. 2.6, we may find a basis of affinoid neighborhoods $U_{x,\delta} \subset \mathcal{E}(\bar{\rho})$ of (x, δ) such that \mathcal{Z} is Zariski-dense in $U_{x,\delta}$, such that $\kappa(U_{x,\delta})$ is an open affinoid, and such that $\kappa : U_{x,\delta} \rightarrow \kappa(U_{x,\delta})$ is finite (necessarily surjective when restricted to any irreducible components of $U_{x,\delta}$). As β is étale at $\kappa(x)$ we may even choose $U_{x,\delta}$ small enough so that the previous assertion holds with $\beta \circ \kappa = \kappa_{HT} \circ \text{pr}_1$ instead of κ . In particular, if $V = \beta(\kappa(U_{x,\delta}))$ then the induced map $\text{pr}_1 : U_{x,\delta} \rightarrow \kappa_{HT}^{-1}(V)$ is a finite map, thus $\text{pr}_1(U_{x,\delta})$ is a *locally closed subset of* $\mathfrak{X}(\bar{\rho})$, and even a closed subset of the admissible open $\kappa_{HT}^{-1}(V) \subset \mathfrak{X}(\bar{\rho})$.

We have constructed this way up to $6^{|S_p|}$ locally closed subspaces of $\mathfrak{X}(\bar{\rho})$, inside $\mathcal{F}(\bar{\rho})$ and containing x , namely the $\text{pr}_1(U_{x,\delta})$, that we will call the *leaves of the fern* at x . Each of these leaves has equidimension $3[F : \mathbb{Q}]$; with this definition, only its germ at x is canonical. By construction, each of these leaves also contains a Zariski-dense subset of modular points, a Zariski-dense subset of which even has the extra property that the eigenvalues of its crystalline Frobenius at p are distinct (see e.g. the proof below of Lemma 2.9), so that each of them will admit exactly $6^{|S_p|}$ associated refined modular points and so on... The *infinite fern* is the union of all the leaves constructed this way, from any modular point, namely

$$\bigcup_{(x,\delta) \in \mathcal{Z}} \text{pr}_1(U_{x,\delta}) \subset \mathfrak{X}(\bar{\rho}),$$

a picture of which has been given in the introduction. Our main aim from now on will be to bound below the dimension of the Zariski-closure of the infinite fern. As explained in the introduction, the idea of our proof is to study the relative positions of the local leaves $\text{pr}_1(U_{x,\delta})$, when δ varies, at a given modular point $x \in \mathfrak{X}(\bar{\rho})$.

2.8. First reduction toward Theorem A

Let

$$\mathfrak{X}^{\text{mod}'} \subset \mathfrak{X}^{\text{mod}}$$

be the subset of modular points x such that for all $v \in S_p$, $\rho_{\Pi,v}$ is irreducible and the eigenvalues of its crystalline Frobenius are distinct and in $k(x)^*$. We claim first that $\mathfrak{X}^{\text{mod}'}$ is dense in $\mathfrak{X}^{\text{mod}}$:

LEMMA 2.9. – *For any $x \in \mathfrak{X}^{\text{mod}}$ and any affinoid neighborhood U of x in $\mathfrak{X}(\bar{\rho})$, we have $\mathfrak{X}^{\text{mod}'} \cap U \neq \emptyset$. In particular, $\mathfrak{X}^{\text{mod}'} \neq \emptyset$.*

Proof. – We use an argument similar to [1, §7.7]: starting from any modular point we show that by moving in at most two leaves of the fern we may find such a modular point by an argument of Newton polygon. So let $x \in \mathfrak{X}^{\text{mod}}$ be any modular point and U as in the statement. Choose any refinement δ of x and consider an affinoid neighborhood $U_{x,\delta}$ of (x, δ) in $\mathcal{E}(\bar{\rho})$ as in §2.7, and such that $\text{pr}_1(U_{x,\delta}) \subset U$. Over this affinoid, the maps $y \mapsto |\mathbf{v}(\delta_{i,v}(p)(y))|$ are bounded by some integer M by the maximum modulus principle. We may choose a point $y \in U_{x,\delta}$ of algebraic weight such that $k_{2,v}(y) - k_{1,v}(y)$ and $k_{3,v}(y) - k_{2,v}(y)$ are both bigger than $M + 1$ for each v , and such that

$$\forall (i, v) \in S_p, \quad \mathbf{v}(\delta_{i,v}(p)(y)) = \mathbf{v}(\delta_{i,v}(p)(x)) = \mathbf{v}(\varphi_{i,v}(x)) - k_{i,v}(x).$$

In particular, such a y is modular by Prop. 2.6. Moreover, for each $(i, v) \in S_p$ we have $\mathbf{v}(\varphi_{i,v}(y)) \neq k_{i+1,v}(y)$ where the index i is taken mod 3. In particular, up to replacing x by y if necessary, we may assume that the point x we started from has this property as well, and also that we had chosen a refinement δ with the property that

$$\forall (i, v) \in I \times S_p, \quad \mathbf{v}(\delta_{i,v}(p)(x)) \neq 0.$$

But then, a modular point y as above has the property that for any $(i, j, v) \in I \times I \times S_p$, $\mathbf{v}(\varphi_{i,v}(y)) \neq k_{j,v}(y)$, so $\rho_{y,v}$ is absolutely irreducible by weak admissibility of its D_{crys} . Its crystalline Frobenius eigenvalues have distinct valuations, so they are distinct and in $k(y)$. \square

REMARK 2.10. – In the examples given in the appendix, the natural modular lift $\text{Sym}^2(V_p(A))(-1)_{|G_{E,S}}$ never belongs to $\mathfrak{X}^{\text{mod}'}$.

If Y is a rigid analytic space and $y \in Y$ a point we denote by $T_y(Y)$ the tangent space of Y at y (a $k(y)$ -vector space).

THEOREM 2.11. – *For any $x \in \mathfrak{X}^{\text{mod}'}$, the image of the natural map $\text{dpr}_1 :$*

$$\bigoplus_{y=(x,\delta) \in \mathcal{E}(\bar{\rho})} T_y(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho}))$$

has dimension at least $6[F : \mathbb{Q}]$.

(Note that for any $x \in \mathfrak{X}^{\text{mod}'}$, the residue field of any $(x, \delta) \in \mathcal{E}(\bar{\rho})$ is $k(x)$ by definition.)

Let us first show that Thm 2.11 implies Thm 1.14. Let W be the Zariski-closure of $\mathfrak{X}^{\text{mod}}$ in $\mathfrak{X}(\bar{\rho})$, equipped with its reduced structure. Let C be an irreducible component of W . Recall that affinoid algebras are excellent and Jacobson, so the regular locus of any reduced rigid analytic space is a Zariski-open and Zariski-dense subspace (see e.g. [20, §1]). Thus we may choose some modular point $x_0 \in C \cap \mathfrak{X}^{\text{mod}}$ such that x_0 is a smooth point of W (hence of C). Choose $U \subset \mathfrak{X}(\bar{\rho})$ an affinoid neighborhood of x_0 , which is small enough so that $U \cap W$ is regular, and equal to $U \cap C$. By Lemma 2.9, we may find a modular point x in U that furthermore belongs to $\mathfrak{X}^{\text{mod}'}$, by construction $x \in U \cap W = U \cap C$. We are going

to apply Theorem 2.11 to this point x . Note that for each associated refined modular point $(x, \delta) \in \mathcal{E}(\bar{\rho})$,

$$(2.1) \quad \text{Im} \left(d\text{pr}_1 : T_{(x, \delta)}(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho})) \right) \subset T_x(C).$$

Indeed, let $U_{x, \delta} \subset \mathcal{E}(\bar{\rho})$ be an affinoid neighborhood of (x, δ) in $\mathcal{E}(\bar{\rho})$ as in §2.7. As Z is Zariski-dense in $U_{x, \delta}$ we have $\text{pr}_1(U_{x, \delta}) \subset W$. Thus if $U_{x, \delta}$ is chosen small enough, then $\text{pr}_1(U_{x, \delta}) \subset U \cap W = U \cap C$, and (2.1) follows. As a consequence, Theorem 2.11 implies that $\dim T_x(C) \geq 6[F : \mathbb{Q}]$. As x is a smooth point of C , it follows that C itself has dimension at least $6[F : \mathbb{Q}]$, and we are done.

The end of the paper will be devoted to the proof of Theorem 2.11. As explained in the introduction, there are two important ingredients, treated in the next two chapters. The first one is a purely local result on the deformation space of a crystalline representation, and the other one a geometric property of eigenvarieties at non-critical classical points. As a motivation, the reader may already have a look to the end of the proof in §5.1.

3. The linear span of trianguline deformations

3.1. The setting

Let L be a finite extension of \mathbb{Q}_p , $G_{\mathbb{Q}_p} = \text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$, $n \geq 1$ an integer and let V be a continuous L -linear representation of $G_{\mathbb{Q}_p}$ of dimension n . If $m \in \mathbb{Z}$, we set $V(m) = V \otimes \chi^m$ where $\chi : G_{\mathbb{Q}_p} \rightarrow \mathbb{Z}_p^*$ is the cyclotomic character.

Denote by \mathcal{C} the category whose objects are the finite dimensional local \mathbb{Q}_p -algebras A equipped with an isomorphism $\pi : A/m_A \xrightarrow{\sim} L$ (so A has a unique structure of L -algebra such that π is L -linear), and whose morphisms are the local L -algebra homomorphisms. Let

$$\mathfrak{X}_V : \mathcal{C} \rightarrow \text{Sets}$$

be the deformation functor of V to \mathcal{C} . Recall that for an object A of \mathcal{C} , $\mathfrak{X}_V(A)$ is the set of isomorphism classes of pairs (V_A, π) where V_A is a free A -module of rank n equipped with a continuous A -linear representation of $G_{\mathbb{Q}_p}$ and $\pi : V_A \otimes_A L \xrightarrow{\sim} V$ is an $L[G_{\mathbb{Q}_p}]$ -isomorphism. Assume that $\text{End}_{G_{\mathbb{Q}_p}}(V) = L$ and that $\text{Hom}_{G_{\mathbb{Q}_p}}(V, V(-1)) = 0$, so that $H^i(G_{\mathbb{Q}_p}, \text{End}_L(V)) = 0$ if $i \geq 2$ and $\dim_L H^1(G_{\mathbb{Q}_p}, \text{End}_L(V)) = n^2 + 1$ by Tate's results on Galois cohomology of local fields. By standard results of Mazur [32], \mathfrak{X}_V is pro-representable, formally smooth over L , of dimension $n^2 + 1$:

PROPOSITION 3.2. – $\mathfrak{X}_V \simeq \text{Spf}(L[[X_0, \dots, X_{n^2}]])$.

Our main aim in this section will be the study and comparison of a collection of subfunctors of \mathfrak{X}_V when V is crystalline in the sense of Fontaine. Two of them are the subfunctors of crystalline and Hodge-Tate deformations, but we shall actually mostly be interested in an additional collection of subfunctors introduced in [1, Chap. 2] under the name of *trianguline deformation functors*. They are some sorts of analogues of the ordinary deformation functors of an ordinary representation defined by Mazur but that apply to any crystalline representation; they depend on the datum of a *refinement* of the crystalline representation. In rank 2 they are close to the h -deformation functors previously defined by Kisin in [29]. They belong

to an even more general collection of *paraboline deformation functors* that we define and study below using the theory of (φ, Γ) -modules over the Robba ring.

3.3. Paraboline deformation functors of (φ, Γ) -modules over the Robba ring

Let A be a finite dimensional \mathbb{Q}_p -algebra, equipped with the topology given by any norm of \mathbb{Q}_p -vector space on A . Recall that the Robba ring with coefficients in A is the A -algebra \mathcal{R}_A of power series

$$f = \sum_{n \in \mathbb{Z}} a_n (z - 1)^n, \quad a_n \in A,$$

converging on some annulus of \mathbb{C}_p of the form $r(f) \leq |z - 1| < 1$. It is a topological A -algebra, namely an inductive limit of projective limits of affinoid algebras. It is equipped with A -linear commuting actions of φ and $\Gamma := \mathbb{Z}_p^*$ given by the formulae $\varphi(f)(z) = f(z^p)$ and $\gamma(f)(z) = f(z^\gamma)$, the action of Γ being continuous. We set $\mathcal{R} = \mathcal{R}_{\mathbb{Q}_p}$, so $\mathcal{R}_A = \mathcal{R} \otimes_{\mathbb{Q}_p} A$.

A (φ, Γ) -module over \mathcal{R}_A is a finite free \mathcal{R}_A -module D equipped with commuting semi-linear actions of φ and Γ , such that $\mathcal{R}\varphi(D) = D$ (“non degeneracy of φ ”) and such that the action of Γ on D is continuous.⁽⁸⁾ We refer to [1, §2.2] for a summary of the basic facts concerning the theory of (φ, Γ) -modules over the Robba ring. If V is a \mathbb{Q}_p -representation of $G_{\mathbb{Q}_p}$, denote by $D_{\text{rig}}(V)$ its associated (φ, Γ) -module over \mathcal{R} . Recall that the functor $V \mapsto D_{\text{rig}}(V)$ is a tensor equivalence between the category of \mathbb{Q}_p -representations of $G_{\mathbb{Q}_p}$ and étale (φ, Γ) -modules over \mathcal{R} (Fontaine, Colmez-Cherbonnier, Kedlaya). In this equivalence, if V is a \mathbb{Q}_p -representation of $G_{\mathbb{Q}_p}$ equipped with a $\mathbb{Q}_p[G_{\mathbb{Q}_p}]$ -linear action of A , then V is free as A -module if and only if $D_{\text{rig}}(V)$ is a (φ, Γ) -module over \mathcal{R}_A by [1] Lemma 2.2.7.

Let D be a (φ, Γ) -module over \mathcal{R}_L and let $\mathcal{F} = (\text{Fil}_i(D))_{i \in \mathbb{Z}}$ be an increasing filtration of D by (φ, Γ) -submodules of D , each $\text{Fil}_i(D)$ being a direct summand as \mathcal{R}_L -submodule. As a convention, we shall always assume that such filtrations are normalized so that $\text{Fil}_i(D) = 0$ if $i \leq 0$ and, for $i \geq 1$, if $\text{Fil}_i(D) = \text{Fil}_{i-1}(D)$ then $\text{Fil}_i(D) = D$. Let us define the *paraboline deformation functor* of D associated to \mathcal{F}

$$\mathfrak{X}_{D, \mathcal{F}} : \mathcal{C} \rightarrow \text{Sets}$$

as follows. For an object A of \mathcal{C} define $\mathfrak{X}_{D, \mathcal{F}}(A)$ as the set of isomorphism classes of triples $(D_A, \text{Fil}_i(D_A), \pi)$ where D_A is a (φ, Γ) -module over \mathcal{R}_A , $\text{Fil}_i(D_A)$ is an increasing filtration of D_A by (φ, Γ) -submodules over \mathcal{R}_A of D , each $\text{Fil}_i(D_A)$ being a direct summand as \mathcal{R}_A -submodule, and where $\pi : D_A \otimes_A L \xrightarrow{\sim} D$ is a (φ, Γ) -module L -isomorphism such that for each $i \geq 0$ we have $\pi(\text{Fil}_i(D_A)) = \text{Fil}_i(D)$. To be explicit, by an isomorphism between two such triples we mean a \mathcal{R}_A -linear isomorphism $\psi : D_A \rightarrow D'_A$ commuting with φ and Γ , mapping Fil_i onto Fil_i , and such that $\pi = \pi' \cdot \psi$.

When $\text{Fil}_1(D) = D$ then $\mathfrak{X}_{D, \mathcal{F}}$ is simply the *deformation functor* of D , and we shall simply denote it by \mathfrak{X}_D . When Fil_i is a complete flag in D , i.e. $\text{rk}_{\mathcal{R}_L}(\text{Fil}_i(D)) = i$ for $i = 1, \dots, \text{rk}_{\mathcal{R}_L}(D)$, $\mathfrak{X}_{D, \mathcal{F}}$ has been studied in [1, §2.3] under the name of trianguline or

⁽⁸⁾ It means the following. Fix an \mathcal{R} -basis of D and for $\gamma \in \Gamma$ let M_γ be the matrix of γ . Then the coefficients of all the M_γ , $\gamma \in \Gamma$, have to converge on some fixed annulus $r < |z - 1| < 1$, and for any $r < r' < 1$ in $p^\mathbb{Q}$ the map $\gamma \mapsto M_\gamma$, $\Gamma \rightarrow M_n(\mathcal{O}[r, r'])$ is continuous, where $\mathcal{O}[r, r']$ is the affinoid algebra of analytic functions on the annulus $r \leq |z - 1| \leq r'$ for the sup. norm.

triangular deformation functor of D . In the case $D = D_{\text{rig}}(V)$ is étale, then $- \mapsto D_{\text{rig}}(-)$ induces an equivalence $\mathfrak{X}_V \xrightarrow{\sim} \mathfrak{X}_D$ by [1, Prop. 2.3.13], and we shall set as well

$$\mathfrak{X}_{V,\varphi} := \mathfrak{X}_V \times_{\mathfrak{X}_D} \mathfrak{X}_{D,\varphi}$$

($\xrightarrow{\sim} \mathfrak{X}_{D,\varphi}$). When furthermore each $\text{Fil}_i(D)$ is étale, hence is the D_{rig} of a unique Galois stable filtration $\text{Fil}_i(V)$, then $\mathfrak{X}_{V,\varphi}(A)$ coincides with the isomorphism classes of deformations of V to A equipped with some Galois stable filtrations over A lifting $\text{Fil}_i(V)$; however, there is no such description for a general φ .

By a well-known result of Mazur [32, Prop. 1], \mathfrak{X}_D is pro-representable if $\text{End}_{(\varphi,\Gamma)}(D) = L$ and D is étale. Let $\text{End}_{(\varphi,\Gamma),\varphi}(D)$ be the L -vector space of (φ,Γ) -module endomorphisms of D preserving $\text{Fil}_i(D)$ for each i .

PROPOSITION 3.4. – $\mathfrak{X}_{D,\varphi}$ admits a versal pro-deformation. If $\text{End}_{(\varphi,\Gamma),\varphi}(D) = L$, then it is pro-representable.

Proof. – Set $F = \mathfrak{X}_{D,\varphi}$. We obviously have $|F(L)| = 1$. Let $A' \rightarrow A$ and $A'' \rightarrow A$ be two morphisms in \mathcal{C} . By Schlessinger's criterion [36] we need to check that the natural map

$$(3.2) \quad F(A' \times_A A'') \rightarrow F(A') \times_{F(A)} F(A'')$$

is bijective whenever $A'' \rightarrow A$ is surjective.

Let $E : \mathcal{C} \rightarrow \text{Ens}$ be the following functor: for an object A of \mathcal{C} , let $E(A)$ be the set of (φ,Γ) -module structures over \mathcal{R}_A on the \mathcal{R}_A -module $D \otimes_L A$ that preserve the *constant filtration* $\text{Fil}_i(D) \otimes_L A$ and that induce the (φ,Γ) -module structure of D via the given map $A \rightarrow L$. For all A, A' and A'' , we claim that (3.2) is bijective when F is replaced by E . Indeed, $\mathcal{R}_{A' \times_A A''} = \mathcal{R}_{A'} \times_{\mathcal{R}_A} \mathcal{R}_{A''}$, and if $R(A)$ denotes the “parabolic” \mathcal{R}_A -algebra of \mathcal{R}_A -module endomorphisms of $D \otimes_L A$ preserving $\text{Fil}_i(D) \otimes_L A$, then R commutes with finite fiber products in \mathcal{C} as well. Moreover, if we fix some basis $(e_i)_{i=1}^n$ of D adapted to the filtration $\text{Fil}_i(D)$, then it is equivalent to give an element of $E(A)$ and a collection of matrices M_* , for $*$ = φ or in Γ , lifting the matrices in $M_n(\mathcal{R}_L)$ defined by the (φ,Γ) -module structure of D , and satisfying $y(M_x)M_y = x(M_y)M_x$ for all $x, y \in \{\varphi, \gamma \in \Gamma\}$. Indeed, the non degeneracy of φ and the continuity of Γ follow automatically from these properties for D .

Set $R^1(A) = \text{Ker}(R(A)^* \rightarrow R(L)^*)$, the group $R^1(A)$ acts by conjugacy on $E(A)$ and we have $F(A) = R^1(A) \backslash E(A)$ by definition. If $A'' \rightarrow A$ is any surjection in \mathcal{C} , then $R^1(A'') \rightarrow R^1(A)$ is surjective as well, as parabolic subgroups of GL_n are smooth over \mathbb{Q}_p . Moreover, R^1 obviously commutes with finite fiber products in \mathcal{C} . It follows at once that (3.2) is surjective whenever $f : A'' \rightarrow A$ is. It is even bijective if for all $x \in E(A'')$, the group homomorphism induced on stabilizers

$$(3.3) \quad R^1(A'')_x \rightarrow R^1(A)_{E(f)(x)}$$

is surjective (these observations really are Mazur's, see [32, p. 390]). This is clearly satisfied if $A = L$ as the latter stabilizer is trivial. As a consequence, conditions (H1) and (H2) of Schlessinger hold. In particular $F(L[\varepsilon])$ is an L -vector space and we shall see in Prop. 3.6 below that it is always finite dimensional (contrary to $E(L[\varepsilon])$ in this (φ,Γ) -module context), hence (H3) and the first part of the statement follow. When $\text{End}_{(\varphi,\Gamma),\varphi}(D) = L$, then $R^1(A)_x = A^*$ for any A and $x \in E(A)$ by Remark 3.5 below, hence (3.3) is surjective, and (H4) holds. \square

REMARK 3.5. – If $\text{End}_{(\varphi, \Gamma), \mathcal{P}}(D) = L$, then a simple induction on the length of A shows that for each element (D_A, Fil_i, π) of $\mathfrak{X}_{D, \mathcal{P}}(A)$, we have $\text{End}_{(\varphi, \Gamma), (\text{Fil}_i)}(D_A) = A$. In particular, $\mathfrak{X}_{D, \mathcal{P}}(A)$ is in canonical bijection with the isomorphism classes of pairs (D_A, Fil_i) lifting D with its filtration (forgetting the π).

If D_1 and D_2 are two (φ, Γ) -modules over \mathcal{R}_L , then $\text{Hom}_{\mathcal{R}_L}(D_1, D_2)$ has a natural structure of (φ, Γ) -module as follows: for any $u \in \text{Hom}_{\mathcal{R}_L}(D_1, D_2)$, set $\gamma(u)(x) := \gamma(u(\gamma^{-1}(x)))$ and $\varphi(u)(\varphi(x)) := \varphi(u(x))$. This last formula defines a unique element $\varphi(u) \in \text{Hom}_{\mathcal{R}_L}(D_1, D_2)$ as D_1 admits an \mathcal{R}_L -basis in $\varphi(D_1)$ by definition. In particular, $\text{End}_{\mathcal{R}_L}(D)$ is a (φ, Γ) -module of rank n^2 over \mathcal{R}_L and we check at once with the formulas above that the \mathcal{R}_L -submodule

$$\text{End}_{\mathcal{P}}(D) := \{u \in \text{End}_{\mathcal{R}_L}(D), u(\text{Fil}_i(D)) \subset \text{Fil}_i(D) \quad \forall i \in \mathbb{N}\}$$

is a (φ, Γ) -submodule of $\text{End}_{\mathcal{R}_L}(D)$ (which is actually a direct summand as \mathcal{R}_L -submodule). For $i \in \mathbb{Z}$ we set $\text{gr}_i(-) := \text{Fil}_i(-)/\text{Fil}_{i-1}(-)$. If $n_i = \text{rk}_{\mathcal{R}_L}(\text{gr}_i(D))$, we obviously have

$$(3.4) \quad \text{rk}_{\mathcal{R}_L}(\text{End}_{\mathcal{P}}(D)) = \sum_{i \leq j} n_i n_j.$$

We refer to [19, §2] and [31] for the main properties of the cohomology of (φ, Γ) -modules over \mathcal{R} , that we shall denote by $H_{(\varphi, \Gamma)}^i(-)$. In particular, $H_{(\varphi, \Gamma)}^0(\text{End}_{\mathcal{P}}(D)) = \text{End}_{(\varphi, \Gamma), \mathcal{P}}(D)$ and $H_{(\varphi, \Gamma)}^2(\text{End}_{\mathcal{P}}(D))$ is dual to the L -vector space of (φ, Γ) -morphisms $D \rightarrow D(-1)$ preserving \mathcal{P} .

PROPOSITION 3.6. – (i) If $\text{Hom}_{(\varphi, \Gamma)}(\text{gr}_i(D), D/\text{Fil}_i(D)) = 0$ for each i , then $\mathfrak{X}_{D, \mathcal{P}}$ is a subfunctor of \mathfrak{X}_D .

(ii) There is a natural isomorphism $\mathfrak{X}_{D, \mathcal{P}}(L[\varepsilon]) \xrightarrow{\sim} H_{(\varphi, \Gamma)}^1(\text{End}_{\mathcal{P}}(D))$ and

$$\dim_L \mathfrak{X}_{D, \mathcal{P}}(L[\varepsilon]) = \dim_L H_{(\varphi, \Gamma)}^0(\text{End}_{\mathcal{P}}(D)) + \dim_L H_{(\varphi, \Gamma)}^2(\text{End}_{\mathcal{P}}(D)) + \sum_{i \leq j} n_i n_j.$$

(iii) If $H_{(\varphi, \Gamma)}^2(\text{End}_{\mathcal{P}}(D)) = 0$ then $\mathfrak{X}_{D, \mathcal{P}}$ is formally smooth over L .

Proof. – Part (i) follows as in [1, Lemma 2.3.7]. The second part of (ii) follows from the first one, (3.4) and the Euler characteristic formula of Liu [31]. Let us check the first part of (ii). It is a semi-linear analogue of a well-known fact in the context of group representations. Using the notations of the proof of Prop. 3.4, for all A we have a natural identification $\mathfrak{X}_{D, \mathcal{P}}(A) = R^1(A) \setminus E(A)$.

Consider the $\mathcal{R}_L[\varepsilon]$ -module $D_0 := D \otimes_L L[\varepsilon] = D \oplus \varepsilon D$ equipped with its constant filtration as in the proof of Prop. 3.4. Any element of $E(L[\varepsilon])$ is given by unique elements $c_* \in \text{End}_{\mathcal{R}}(D)$, where $*$ = φ or $\gamma \in \Gamma$, satisfying the formulas

$$\tilde{\gamma}(x) := \gamma(x) + c_{\gamma}(\gamma(x))\varepsilon, \quad \tilde{\varphi}(x) := \varphi(x) + c_{\varphi}(\varphi(x))\varepsilon, \quad \forall x \in D \subset D \oplus \varepsilon D.$$

Note that the map c_{φ} (resp. c_{γ}) is well-defined as D admits an \mathcal{R}_L -basis in $\varphi(D)$ (resp. as γ is bijective on D). A straightforward computation shows that the commutation of $\tilde{\varphi}$ and $\tilde{\gamma}$ is equivalent to the relation $(\varphi - 1)c_{\gamma} = (\gamma - 1)c_{\varphi}$ in $\text{End}_{\mathcal{R}}(D)$, for the (φ, Γ) -module structure recalled above on $\text{End}_{\mathcal{R}}(D)$. The continuity of the action of Γ is automatic, as well as the non degeneracy of $\tilde{\varphi}$. Moreover, $*$ preserves each $\text{Fil}_i D \otimes_L L[\varepsilon]$ if, and only if, $c_* \in \text{End}_{\mathcal{P}}(D)$. In other words, c_* is a 1-cocycle in $\text{End}_{\mathcal{P}}(D)$ and $E(L[\varepsilon])$ coincides with

this space of cocycles. Another straightforward computation shows that two such lifts are isomorphic in $\mathfrak{X}_{D,\varphi}(L[\varepsilon]) = E(L[\varepsilon])/R^1(L[\varepsilon])$ if and only if the associated 1-cocycles c_* and c'_* differ by $(* - 1)u$ for some $u \in \text{End}_\varphi(D)$ independent of $*$, hence a coboundary, which concludes the proof of (ii) (the L -linearity part of the statement is immediate).

For part (iii), let A be an object of \mathcal{C} , m its maximal ideal, and I an ideal of A such that $mI = 0$. Fix an element of $E(A/I)$, hence a $\mathcal{R}_{A/I}$ -linear (φ, Γ) -module structure on $D \otimes_L A/I$ preserving $\text{Fil}_i(D) \otimes_L A/I$, and consider the problem of lifting this structure to an element of $E(A)$. First, we may certainly lift the actions of φ and of Γ to $D \otimes_L A$ in such a way that they preserve $\text{Fil}_i(D) \otimes_L A$. If $\tilde{\varphi}$ and $\tilde{\gamma}$ denote such a lift, consider the \mathcal{R}_L -linear map $\langle \tilde{\varphi}, \tilde{\gamma} \rangle \in \text{End}_\varphi(D) \otimes_L I$ defined by the formula

$$\forall x \in D \otimes_L A, (\tilde{\varphi}\tilde{\gamma} - \tilde{\gamma}\tilde{\varphi})(x) = \langle \tilde{\varphi}, \tilde{\gamma} \rangle(\varphi\gamma(\bar{x})),$$

where \bar{x} is the image of x in D . This map $\langle \tilde{\varphi}, \tilde{\gamma} \rangle$ is well-defined as γ is bijective on D and as D admits an \mathcal{R} -basis in $\varphi(D)$. The elements $\tilde{\varphi}$ and $\tilde{\gamma}$ are uniquely defined up to adding any elements $u(\varphi(\cdot))$ and $v(\gamma(\cdot))$ where $u, v \in \text{End}_\varphi(D) \otimes_L I$, and

$$\langle \tilde{\varphi} + u(\varphi(\cdot)), \tilde{\gamma} + v(\gamma(\cdot)) \rangle = \langle \tilde{\varphi}, \tilde{\gamma} \rangle + (\varphi - 1)v - (\gamma - 1)u.$$

This concludes the proof as $H_{(\varphi, \Gamma)}^2(\text{End}_\varphi(D)) = \text{End}_\varphi(D)/(\varphi - 1, \gamma - 1)\text{End}_\varphi(D)$. Note that as in the classical case, it is straightforward to check that the set of lifts is either empty or an affine space under $H_{(\varphi, \Gamma)}^1(\text{End}_\varphi(D)) \otimes_L I$. \square

Let $I \subset \mathbb{N}$ be the (finite) subset of jumps of the filtration Fil_i , i.e. the integers i such that $\text{gr}_i(D) \neq 0$. There is a natural functor morphism $\mathfrak{X}_{D,\varphi} \rightarrow \prod_{i \in I} \mathfrak{X}_{\text{gr}_i(D)}$ sending an object (D_A, Fil_i, π) to $(\text{gr}_i(D_A), \text{gr}_i(\pi))_{i \in I}$.

PROPOSITION 3.7. – *If $H_{(\varphi, \Gamma)}^2(\text{Hom}_{\mathcal{R}}(D/\text{Fil}_i(D), \text{gr}_i(D))) = 0$ for each i , then $\mathfrak{X}_{D,\varphi} \rightarrow \prod_{i \in I} \mathfrak{X}_{\text{gr}_i(D)}$ is formally smooth. In particular, it is surjective on points.*

Proof. – The proposition is obvious when $|I| \leq 1$ so assume that $|I| > 1$. Considering the filtration Fil' on D such that $\text{Fil}'_1(D) = \text{Fil}_j(D)$ for j the biggest integer such that $\text{Fil}_j(D) \neq D$, $\text{Fil}'_2(D) = D$, and arguing by induction on $|I|$, we may assume that $I = \{1, 2\}$. So D is an extension of $D_1 := D/\text{Fil}_1(D)$ by $D_2 := \text{Fil}_1(D)$, which defines a unique class $c \in H_{(\varphi, \Gamma)}^1(\text{Hom}_{\mathcal{R}}(D_1, D_2))$. Let A be an object of \mathcal{C} , m its maximal ideal, and I an ideal of A such that $mI = 0$. Fix $(U, \text{Fil}_i(U), \pi) \in \mathfrak{X}_{\varphi, D}(A/I)$, it defines as above an element $c_{A/I} \in H_{(\varphi, \Gamma)}^1(\text{Hom}_{\mathcal{R}}(U/\text{Fil}_1(U), \text{Fil}_1(U)))$ that maps to c modulo m . For $i = 1, 2$, choose $(D'_i, \pi_i) \in \mathfrak{X}_{D_i}(A)$ lifting respectively $U/\text{Fil}_1(U)$ and $\text{Fil}_1(U)$. We have to show that the natural map

$$H_{(\varphi, \Gamma)}^1(\text{Hom}_{\mathcal{R}}(D'_1, D'_2)) \longrightarrow H_{(\varphi, \Gamma)}^1(\text{Hom}_{\mathcal{R}}(U/\text{Fil}_1(U), \text{Fil}_1(U)))$$

is surjective. But its cokernel injects into $H_{(\varphi, \Gamma)}^2(\text{Hom}_{\mathcal{R}}(D_1, D_2))$. \square

EXAMPLE 3.8 (The rank 1 case). – Let A be an object of \mathcal{C} and let $\delta : \mathbb{Q}_p^* \rightarrow A^*$ be a continuous character. We denote by $\mathcal{R}_A(\delta)$ the (φ, Γ) -module of rank 1 over \mathcal{R}_A having a basis e such that $\varphi(e) = \delta(p)e$ and $\gamma(e) = \delta(\gamma)e$ for all $\gamma \in \Gamma$. By [1, Prop. 2.3.1], each (φ, Γ) -module of rank 1 over \mathcal{R}_A has the form $\mathcal{R}_A(\delta)$ for a unique character δ as above. In particular, if D is a (φ, Γ) -module of rank 1 over \mathcal{R}_L , then $\mathfrak{X}_D \simeq \text{Spf}(L[[X, Y]])$.

3.9. Crystalline (φ, Γ) -modules

If D is any (φ, Γ) -module over \mathcal{R} , we set $\mathcal{D}_{\text{crys}}(D) = (D[1/t])^\Gamma$, where $t = \log(z) \in \mathcal{R}$. As \mathcal{R} is a domain and $\text{Frac}(\mathcal{R})^\Gamma = \mathbb{Q}_p$, a standard argument shows that

$$(3.5) \quad \mathcal{D}_{\text{crys}}(D) \otimes_L \mathcal{R}_L[1/t] \rightarrow D[1/t]$$

is injective for any D , so we have $\dim_{\mathbb{Q}_p} \mathcal{D}_{\text{crys}}(D) \leq \text{rk}_{\mathcal{R}} D$. Mimicking Fontaine, we say that D is *crystalline* if the equality holds. By left exactness of $- \mapsto \mathcal{D}_{\text{crys}}(-)$, any (φ, Γ) -module subquotient of a crystalline D is also crystalline. Note that $\mathcal{D}_{\text{crys}}(D)$ has a \mathbb{Q}_p -linear invertible action of φ .

LEMMA 3.10. – *Let D be a (φ, Γ) -module over \mathcal{R}_L . The map*

$$D' \mapsto \mathcal{D}_{\text{crys}}(D')$$

induces a bijection between the set of crystalline (φ, Γ) -submodules of D which are direct summand as \mathcal{R}_L -module, and the set of φ -stable subspaces of $\mathcal{D}_{\text{crys}}(D)$. The inverse bijection is $W \mapsto (\mathcal{R}[1/t] \cdot W) \cap D$.

Indeed, a (φ, Γ) -submodule $D' \subset D$ is a direct summand as \mathcal{R}_L -module if, and only if $D' = D'[1/t] \cap D$ by [1, Prop. 2.2.2], so the lemma follows from (3.5). In particular, there is always a biggest crystalline (φ, Γ) -submodule of D , namely $(\mathcal{R}[1/t] \cdot D[1/t]^\Gamma) \cap D$. Furthermore, when D is crystalline then Lemma 3.10 induces a rank preserving bijection between the set of filtrations of D as in §3.3 and the set of increasing filtrations $(\text{Fil}_i)_{i \in \mathbb{Z}}$ of $\mathcal{D}_{\text{crys}}(D)$ by φ -stable \mathbb{Q}_p -vector space (normalized with the same conventions as before), we shall often identify such filtrations below. In particular, the paraboline deformation functors of D can (and will) be viewed as attached to such filtrations.

Consider the following properties of a crystalline (φ, Γ) -module of rank n over \mathcal{R}_L :

- (i) φ has n distinct eigenvalues in L^* on $\mathcal{D}_{\text{crys}}(D)$.
- (ii) For any two such eigenvalues ϕ, ϕ' , we have $\phi' \phi^{-1} \neq p$.
- (iii) $\text{End}_{(\varphi, \Gamma)}(D) = L$.

COROLLARY 3.11. – *Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L satisfying (i), (ii) and (iii). Then each paraboline deformation functor $\mathfrak{X}_{D, \varphi}$ of D is a pro-representable subfunctor of \mathfrak{X}_D , formally smooth over L of dimension $\text{rk}_{\mathcal{R}_L}(\text{End}_{\varphi}(D)) + 1$.*

Proof. – The pro-representability follows from Prop. 3.4 and (iii). As D is crystalline, remark that $\text{Hom}_{(\varphi, \Gamma), L}(D, D') \subset \text{Hom}_{L[\varphi]}(\mathcal{D}_{\text{crys}}(D), \mathcal{D}_{\text{crys}}(D'))$ for any (φ, Γ) -module D' over \mathcal{R}_L . By Prop. 3.6, the subfunctor property follows from (i) and the formal smoothness from (ii). \square

For any D , Berger defines as well a decreasing, exhaustive, filtration on $\mathcal{D}_{\text{crys}}(D)$ called the *Hodge-filtration*, that we shall denote with upper indices Fil^i for $i \in \mathbb{Z}$ (see [3], [4], as well as [1, §2.2.7]). A first important result of Berger is that when $D = D_{\text{rig}}(V)$, the filtered φ -module $\mathcal{D}_{\text{crys}}(D)$ is canonically isomorphic to the classical $D_{\text{crys}}(V)$ defined by Fontaine. In particular, V is crystalline if, and only if, D is. Another beautiful result of Berger [4] is that $D \mapsto \mathcal{D}_{\text{crys}}(D)$ induces an exact \otimes -equivalence of category between the category of crystalline (φ, Γ) -modules over \mathcal{R} and the category of filtered φ -vector space, étale corresponding to weakly admissible.

Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L . Define $\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}_D$ as the subfunctor of deformations (D_A, π) such that D_A is crystalline, viewed as (φ, Γ) -module over \mathcal{R} . As the crystalline (φ, Γ) -modules are stable by direct sums, subquotients, and tensor products, $\mathfrak{X}_{D, \text{crys}} \rightarrow \mathfrak{X}_D$ is relatively representable (Ramakrishnan's criterion).

PROPOSITION 3.12. – *Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L .*

- (a) *If D is equipped with an action of an object A of \mathcal{C} , then D is free as \mathcal{R}_A -module if and only if $\text{Fil}^i(\mathcal{D}_{\text{crys}}(D))$ is free as A -module for each $i \in \mathbb{Z}$.*
- (b) *The functor $- \mapsto \mathcal{D}_{\text{crys}}(-)$ induces a canonical bijection between $\mathfrak{X}_{D, \text{crys}}(A)$ and the set of isomorphism classes of pairs (E_A, π) where E_A is a filtered $A[\varphi]$ -module such that $\text{Fil}^i E_A$ is free and direct summand as A -module and $\pi : E_A \otimes_A L \xrightarrow{\sim} \mathcal{D}_{\text{crys}}(D)$ is an A -linear isomorphism in the category of filtered φ -modules.*
- (c) *If D satisfies properties (i) and (iii) above, then $\mathfrak{X}_{D, \text{crys}}$ is formally smooth over L of dimension*

$$1 + \sum_{i < j} n_i n_j,$$

where $n_i = \dim_L \text{Fil}^i(\mathcal{D}_{\text{crys}}(D)) / \text{Fil}^{i+1}(\mathcal{D}_{\text{crys}}(D))$ for all $i \in \mathbb{Z}$.

Proof. – Part (a) holds by the same proof as [1, Lemma 2.2.7]. Part (b) follows from (a) and Berger's equivalence. Using (b), part (c) is a simple exercise on deformations of filtered φ -modules using (i) and (iii) that is left to the reader. \square

Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L and let $W \subset \mathcal{D}_{\text{crys}}(D)$ be an L -subvector space. We say that W is *non-critical* (in $\mathcal{D}_{\text{crys}}(D)$) if there exists an integer $j \in \mathbb{Z}$ such that $\mathcal{D}_{\text{crys}}(D) = W \oplus \text{Fil}^j(\mathcal{D}_{\text{crys}}(D))$. If $\mathcal{P} = (\text{Fil}_i)$ is a φ -stable filtration of $\mathcal{D}_{\text{crys}}(D)$ as above, we say that \mathcal{P} is non-critical if Fil_i is non-critical for all $i \geq 0$.

PROPOSITION 3.13. – *Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L satisfying (i) and (iii). For any non-critical filtration \mathcal{P} , we have $\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}_{D, \mathcal{P}}$.*

Proof. – By (i), $\mathfrak{X}_{D, \mathcal{P}}$ is a subfunctor of \mathfrak{X}_D . Fix $(D_A, \pi) \in \mathfrak{X}_{D, \text{crys}}(A)$ and let $W \subset \mathcal{D}_{\text{crys}}(D)$ be an element of \mathcal{P} . By (i) again, there is a unique φ -stable A -submodule $W_A \subset \mathcal{D}_{\text{crys}}(D_A)$ which is free over A and such that $\pi(W_A) = W$. Note that if $W \subset W'$ then $W_A \subset W'_A$. Set $T_A = (\mathcal{R}[1/t]W_A) \cap D$, we will show that it is free over \mathcal{R}_A and lifts $(\mathcal{R}[1/t]W) \cap D$.

By Prop. 3.12 (a) and Berger's equivalence, it is equivalent to check that if we equip W_A with the induced Hodge-filtration of $\mathcal{D}_{\text{crys}}(D_A)$, then $\text{Fil}^i(W_A)$ is free over A for each i and lifts $\text{Fil}^i(W)$. But if j is such that $\text{Fil}^j(\mathcal{D}_{\text{crys}}(D)) \oplus W = \mathcal{D}_{\text{crys}}(D)$, then $\text{Fil}^j(\mathcal{D}_{\text{crys}}(D_A)) \oplus W_A = \mathcal{D}_{\text{crys}}(D_A)$ by Nakayama's lemma, and the result follows. \square

Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L . For the applications of this paper, the most important paraboline deformation functors of D will be the ones associated to the $L[\varphi]$ -stable filtrations $\mathcal{P} = (\text{Fil}_i)$ of $\mathcal{D}_{\text{crys}}(D)$ which are *complete flags*, i.e. $\dim_L \text{Fil}_i = i$ for $0 \leq i \leq \text{rk}_{\mathcal{R}_L}(D)$. Such a filtration is called a *refinement* of D , and we shall usually denote them by the letter \mathcal{F} . Via Lemma 3.10, refinements of D correspond to triangulations of D in the sense of Colmez [19]. Note that D admits a refinement (or a triangulation) if and only if the characteristic polynomial of φ on $\mathcal{D}_{\text{crys}}(D)$ splits in L . Before stating the main result

about the trianguline deformation functors of D , we need some preliminary remarks about the extension of Sen's theory to (φ, Γ) -modules.

For any (φ, Γ) -module D over \mathcal{R} , it makes sense to consider its evaluation at a p^n -th root of unity for any sufficiently big n , which defines a semi-linear Γ -module $\mathcal{D}_{\text{Sen}}(D)$ of rank $\text{rk}_{\mathcal{R}} D$ over $\mathbb{Q}_p(\mu_{p^\infty})$ by extending the scalars (see e.g. [1, §2.2.7]). The characteristic polynomial of the Sen operator of $\mathcal{D}_{\text{Sen}}(D)$ will be denoted by $P_{\text{Sen}}(D)$, its eigenvalues are called the (generalized) Hodge-Tate weights of D . As in the classical case, let us say that D is Hodge-Tate if its Sen operator is semi-simple with integral eigenvalues. If D is crystalline, it is Hodge-Tate and its Hodge-Tate weights are the jumps (with multiplicity) of its Hodge-filtration. When D is a (φ, Γ) -module over \mathcal{R}_A , then $\mathcal{D}_{\text{Sen}}(D)$ is a free $A \otimes_{\mathbb{Q}_p} \mathbb{Q}_p(\mu_{p^\infty})$ -module, hence we may also consider the relative characteristic polynomial $P_{\text{Sen}/A}(D)$ of the Sen operator viewed as an A -linear endomorphism.

Let D be a Hodge-Tate (φ, Γ) -module over \mathcal{R}_L . There is a relatively representable subfunctor $\mathfrak{X}_{D, \text{Sen}} \subset \mathfrak{X}_D$ parameterizing deformations D_A which are Hodge-Tate, viewed as (φ, Γ) -module over \mathcal{R} . For any D , the relative Sen polynomial induces a natural functor $P_{\text{Sen}} : \mathfrak{X}_D \rightarrow (\widehat{\mathbb{G}}_a)^d$, this latter space being identified with the completion at $P_{\text{Sen}}(D)$ of the (affine) space of monic polynomial of degree d . The following lemma is easy ([1, Lemma 2.2.11, Prop. 2.3.3]).

LEMMA 3.14. – (a) *If $0 \rightarrow D \rightarrow D' \rightarrow D'' \rightarrow 0$ is an exact sequence of (φ, Γ) -modules over \mathcal{R}_A , then $P_{\text{Sen}/A}(D') = P_{\text{Sen}/A}(D)P_{\text{Sen}/A}(D'')$.*

(b) *If $\delta : \mathbb{Q}_p^* \rightarrow A^*$ is a continuous character, $P_{\text{Sen}/A}(\mathcal{R}_A(\delta)) = T + \left(\frac{\partial \delta}{\partial \gamma}\right)_{|\gamma=1}$. In particular, if D has rank 1 over \mathcal{R}_L then $P_{\text{Sen}} : \mathfrak{X}_D \rightarrow \widehat{\mathbb{G}}_a$ is formally smooth.*

Consider the following property:

(iv) $P_{\text{Sen}/L}(D)$ has $\text{rk}_{\mathcal{R}_L} D$ distinct roots in \mathbb{Z} .

Under (iv), $\mathfrak{X}_{D, \text{Sen}}$ is simply the locus of \mathfrak{X}_D defined by $P_{\text{Sen}} = P_{\text{Sen}}(D)$. When D is crystalline, $\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}_{D, \text{Sen}}$.

PROPOSITION 3.15. – *Let D be a crystalline (φ, Γ) module of rank n over \mathcal{R}_L satisfying (i), (ii), (iii) and (iv) above. Let \mathcal{I} be a non-critical refinement of D .*

Then $\mathfrak{X}_{D, \text{crys}}$ and $\mathfrak{X}_{D, \mathcal{I}}$ are pro-representable subfunctors of \mathfrak{X}_D , and they are formally smooth over L of respective dimension $\frac{n(n-1)}{2} + 1$ and $\frac{n(n+1)}{2} + 1$. The morphism P_{Sen} is formally smooth and $\mathfrak{X}_{D, \text{crys}} = \mathfrak{X}_{D, \text{Sen}} \times_{\mathfrak{X}_D} \mathfrak{X}_{D, \mathcal{I}}$. In particular, we have an exact sequence on tangent spaces:

$$0 \rightarrow \mathfrak{X}_{D, \text{crys}}(L[\varepsilon]) \rightarrow \mathfrak{X}_{D, \mathcal{I}}(L[\varepsilon]) \xrightarrow{P_{\text{Sen}}} L^n \rightarrow 0.$$

Proof. – This is [1, Thm. 2.5.1, Thm. 2.5.10]. Here is a slightly different proof. The first assertion follows from Cor. 3.11, Prop. 3.12 (c) and Prop. 3.13, as well as the inclusion

$$\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}' := \mathfrak{X}_{D, \text{Sen}} \times_{\mathfrak{X}_D} \mathfrak{X}_{D, \mathcal{I}}.$$

The formal smoothness of P_{Sen} follows from Lemma 3.14 and Prop. 3.7. It implies the surjectivity of P_{Sen} in the sequence of the statement, as well as the exactness of the whole sequence for dimensions reasons. It also implies that \mathfrak{X}' is formally smooth over L , hence that $\mathfrak{X}_{D, \text{crys}} = \mathfrak{X}'$ as they have the same dimension. \square

The other kind of paraboline deformation functor that will play a role in the sequel is the “miraboline” case. Assume that D is a crystalline (φ, Γ) -module over \mathcal{R}_L satisfying (i) and let $\phi \in L^*$ be an eigenvalue of φ on $\mathcal{D}_{\text{crys}}(D)$. By Lemma 3.10, there is a unique (φ, Γ) -submodule D_ϕ of rank 1, as well as a unique (φ, Γ) -module D^ϕ of rank $n - 1$, both direct summands as \mathcal{R}_L -modules, such that

$$\mathcal{D}_{\text{crys}}(D_\phi)^{\varphi=\phi} \neq 0 \quad \text{and} \quad \mathcal{D}_{\text{crys}}(D/D^\phi)^{\varphi=\phi} \neq 0.$$

Consider the filtration \mathcal{P}_ϕ (resp. \mathcal{P}^ϕ) of D whose unique proper subspace is $\text{Fil}_1 = D_\phi$ (resp. $\text{Fil}_1 = D^\phi$). It will be convenient to modify a little bit those functors by fixing some Hodge-Tate weight. Assume that (iv) holds and let $k_1 < k_2 < \dots < k_n$ be the Hodge-Tate weights of D . We define $\mathfrak{X}_{D, \phi \downarrow}$ (resp. $\mathfrak{X}_{D, \phi \uparrow}$) as the subfunctor of $\mathfrak{X}_{D, \mathcal{P}_\phi}$ (resp. $\mathfrak{X}_{D, \mathcal{P}^\phi}$) defined by $\text{P}_{\text{Sen}}(k_1) = 0$ (resp. $\text{P}_{\text{Sen}}(k_n) = 0$).

PROPOSITION 3.16. – *Assume that D is a crystalline (φ, Γ) -module of rank n over \mathcal{R}_L satisfying (i), (ii), (iii) and (iv) and let ϕ be an eigenvalue of φ on $\mathcal{D}_{\text{crys}}(D)$. Then $\mathfrak{X}_{D, \phi \downarrow}$ is a pro-representable subfunctor of \mathfrak{X}_D , formally smooth over L of dimension $n^2 - n + 1$.*

Assume that $\mathcal{D}_{\text{crys}}(D)^{\varphi=\phi}$ is non-critical in $\mathcal{D}_{\text{crys}}(D)$. Then $\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}_{D, \phi \downarrow}$. Furthermore, for any object A in \mathcal{C} , $\mathfrak{X}_{D, \phi \downarrow}(A)$ is the subset of $(D_A, \pi) \in \mathfrak{X}_D(A)$ such that there exists $\tilde{\phi} \in A^$ lifting ϕ and such that $\mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}}$ is free of rank 1 over A .*

Proof. – By Lemma 3.11 (i), $\mathfrak{X}_{D, \mathcal{P}_\phi}$ is a subfunctor of \mathfrak{X}_D , formally smooth over L of dimension $n^2 - n + 2$. The assertion on $\mathfrak{X}_{D, \phi \downarrow}$ follows from this, property (ii), Prop. 3.7 and Lemma 3.14. The inclusion $\mathfrak{X}_{D, \text{crys}} \subset \mathfrak{X}_{D, \phi \downarrow}$ follows from Prop. 3.13.

Before checking the last assertion, remark that for an element $(D_A, \text{Fil}_i, \pi) \in \mathfrak{X}_{D, \mathcal{P}_\phi}(A)$, and $\tilde{\phi} \in A^*$ lifting ϕ , the natural inclusion induces a bijection

$$(3.6) \quad \mathcal{D}_{\text{crys}}(\text{Fil}_1(D_A))^{\varphi=\tilde{\phi}} = \mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}}.$$

Indeed, $- \mapsto \mathcal{D}_{\text{crys}}(-)^{\varphi=\tilde{\phi}}$ is left exact over the category of (φ, Γ) -modules over \mathcal{R}_A and $\mathcal{D}_{\text{crys}}(D_A/\text{Fil}_1)^{\varphi=\tilde{\phi}}$ vanishes by a dévissage as $\mathcal{D}_{\text{crys}}(D/\text{Fil}_1(D))^{\varphi=\phi} = 0$.

Let us check the last assertion. Define $\mathfrak{X}'(A)$ as the subset of $(D_A, \pi) \in \mathfrak{X}_D(A)$ having the property given in the statement. Assume first that $(D_A, \text{Fil}_i, \pi) \in \mathfrak{X}_{D, \phi \downarrow}(A)$. Write $\text{Fil}_1 \simeq \mathcal{R}_A(\delta)$ for some character $\delta : \mathbb{Q}_p^* \rightarrow A^*$. As $\mathcal{D}_{\text{crys}}(D_\phi)$ is non-critical $\text{P}_{\text{Sen}/L}(D_\phi) = T - k_1$, so the reduction $\bar{\delta} : \mathbb{Q}_p^* \rightarrow L^*$ of δ modulo m_A is the algebraic character $x \mapsto x^{-k_1}$ over \mathbb{Z}_p^* (Lemma 3.14 (b)). As k_1 is a simple root of $\text{P}_{\text{Sen}L}(D)$ by (iv), the assumption $\text{P}_{\text{Sen}/A}(D_A)(k_1) = 0$ and Lemma 3.14 ensure that

$$\text{P}_{\text{Sen}/A}(\text{Fil}_1(D_A)) = \text{P}_{\text{Sen}/A}(\mathcal{R}_A(\delta)) = T - k_1.$$

This implies that $\delta(x) = x^{-k_1}$ for any $x \in \mathbb{Z}_p^*$, so $\text{Fil}_1(D_A)$ is crystalline, hence a crystalline deformation of D_ϕ . By (3.6), this implies that $(D_A, \pi) \in \mathfrak{X}'(A)$ (even with $\tilde{\phi} = \delta(p)p^{-k_1}$).

We now check the other inclusion $\mathfrak{X}'(A) \subset \mathfrak{X}_{D, \phi \downarrow}(A)$. Fix $(D_A, \pi) \in \mathfrak{X}'(A)$ and let $\tilde{\phi} \in A^*$ be as in the statement. We claim first that

$$\text{Fil}^i(\mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}}) = 0 \quad \text{for all } i > k_1.$$

Indeed, let us argue by induction on $\dim_L A$. It holds when $\dim_L A = 1$ as D_ϕ is non-critical. When $\dim_L A > 1$ we may find an ideal $I \subset A$ of L -dimension 1, and as D_A is free over \mathcal{R}_A

we may consider the exact sequence $0 \rightarrow ID_A = D \rightarrow D_A \rightarrow D_{A/I} \rightarrow 0$ of (φ, Γ) -modules over \mathcal{R}_L (and of \mathcal{R}_A -modules). Applying $\mathcal{D}_{\text{crys}}(-)^{\varphi=\tilde{\phi}}$ we obtain for any i an exact sequence $0 \rightarrow \text{Fil}^i \mathcal{D}_{\text{crys}}(D)^{\varphi=\phi} \rightarrow \text{Fil}^i \mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}} \rightarrow \text{Fil}^i \mathcal{D}_{\text{crys}}(D_{A/I})^{\varphi=\tilde{\phi}}$ from which we get the claim by induction. As a consequence, the crystalline (φ, Γ) -submodule $D'_A \subset D_A$ corresponding to the filtered submodule $\mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}}$ is free over \mathcal{R}_A by Lemma 3.12. The natural map $\pi : D'_A \rightarrow D$ surjects onto D_ϕ as it is so after applying $\mathcal{D}_{\text{crys}}(-)$, and we are done. (For an alternative argument, see [1, Lemma 2.5.2].) \square

REMARK 3.17. – The duality $E \mapsto E^\vee = \text{Hom}_{\mathcal{R}_A}(E, \mathcal{R}_A)$ on (φ, Γ) -modules E over \mathcal{R}_A induces an isomorphism $\iota : \mathfrak{X}_D \xrightarrow{\sim} \mathfrak{X}_{D^\vee}$. Note that D^\vee satisfies (i), (ii), (iii), or (iv) if D does. If \mathcal{P} is a filtration on D , then there is a natural dual filtration \mathcal{P}^\vee on D^\vee . When D is crystalline, \mathcal{P} is non-critical if and only if \mathcal{P}^\vee is. The isomorphism ι induces an isomorphism $\mathfrak{X}_{D, \mathcal{P}} \xrightarrow{\sim} \mathfrak{X}_{D^\vee, \mathcal{P}^\vee}$, as well as an isomorphism $\mathfrak{X}_{D, \phi \uparrow} \xrightarrow{\sim} \mathfrak{X}_{D^\vee, \phi^{-1} \downarrow}$ (when it makes sense). In particular, the proposition above has an obvious analogue for $\mathfrak{X}_{D, \phi \uparrow}$ whose statement is left as an exercise to the reader.

3.18. The main theorem

Let D be a crystalline (φ, Γ) -module of rank $n \geq 1$ over \mathcal{R}_L . We say that D is *generic* if it satisfies (i), (ii) of § 3.9, and if *all of its $n!$ refinements are non-critical*. As we shall see below, such a D necessarily satisfies (iii) and (iv) as well. In the algebraic variety of all filtered φ -modules of dimension n over L whose filtration admits n jumps, the generic condition is Zariski-open and Zariski-dense.

Let us denote by \mathfrak{t} the tangent space $\mathfrak{X}_D(L[\varepsilon])$, and for $* \in \{\mathcal{F}, \phi \downarrow, \text{crys} \dots\}$ let \mathfrak{t}_* denote the tangent space $\mathfrak{X}_{D, *}(L[\varepsilon])$. By Lemma 3.11, \mathfrak{t} contains the L -subvector-space $\mathfrak{t}_{\mathcal{F}}$ for any of the $n!$ refinements \mathcal{F} of D .

THEOREM 3.19. – *If D is generic then $\mathfrak{t} = \sum_{\mathcal{F}} \mathfrak{t}_{\mathcal{F}}$.*

In other words, *any first order deformation of a generic crystalline representation is a linear combination of trianguline deformations*.

We will actually prove a more general statement, whose formulation requires some more definitions. Let $I \subset \{1, \dots, n\}$ be a non-empty subset and denote by $i_1 < \dots < i_r$ its elements. An element σ in \mathfrak{S}_n will be called an *ordered cycle* on I if either $\sigma = (i_1, i_2, \dots, i_r)$ or $\sigma = (i_r, i_{r-1}, \dots, i_1)$. By a *nested* sequence of intervals in $\{1, \dots, n\}$ we mean a decreasing sequence of subsets

$$I_0 = \{1, \dots, n\} \supset I_1 \supset \dots \supset I_{n-1}$$

such that for each $1 \leq i \leq n-1$, $I_i = I_{i-1} \setminus \{y_i\}$ where $y_i \in \{\text{Min}(I_{i-1}), \text{Max}(I_{i-1})\}$. Such a sequence is uniquely determined by the sequence (y_i) entering in its definition. For each $1 \leq i \leq n-1$ we define as well an element $y_i^* \in \{1, \dots, n\}$ by $\{y_i, y_i^*\} = \{\text{Min}(I_{i-1}), \text{Max}(I_{i-1})\}$.

We will say that a sequence of n permutations $\sigma_0, \sigma_1, \dots, \sigma_{n-1} \in \mathfrak{S}_n$ is *nested* if $\sigma_0 = \text{id}$ and if there exists a nested sequence of intervals (I_i) of $\{1, \dots, n\}$ such that

$$\forall i \in \{1, \dots, n-1\}, \quad \sigma_i = c_i \sigma_{i-1},$$

where c_i is the ordered cycle on I_{i-1} such that $c_i(y_i^*) = y_i$ (the y_i and y_i^* being associated to I_i as above). For $n > 1$ there are exactly 2^{n-2} nested sequences in \mathfrak{S}_n . For instance, id , $(1, 2, 3)$, $(2, 3)(1, 2, 3)$ and id , $(3, 2, 1)$, $(1, 2)(3, 2, 1)$ are the two nested sequences in \mathfrak{S}_3 .

We will say that a sequence of $2n - 1$ permutations

$$\sigma_0, \sigma_1, \sigma_1^*, \sigma_2, \sigma_2^*, \dots, \sigma_{n-1}, \sigma_{n-1}^* \in \mathfrak{S}_n$$

is *weakly nested* if $\sigma_0 = \text{id}$ and if there exists a nested sequence of intervals (I_i) of $\{1, \dots, n\}$ such that

$$\forall i \in \{1, \dots, n-1\}, \quad \sigma_i = \tau_i \sigma_{i-1} \quad \text{and} \quad \sigma_i^* = c_i \sigma_{i-1},$$

where c_i is the ordered cycle on I_{i-1} such that $c_i(y_i^*) = y_i$, and where $\tau_i \in \mathfrak{S}_n$ is any element such that $\tau_i(y_i^*) = y_i$ and such that $\tau_i(j) = j$ for all $j \in \{1, \dots, n\} \setminus I_{i-1}$. For example, the nested sequence of intervals defined by $y_i = i$ for each i , and the elements $\tau_i = (i, n)$ for $1 \leq i \leq n-1$, define a unique weakly nested sequence in \mathfrak{S}_n .

Of course, a nested sequence defines a weakly nested sequence if we set $\sigma_i^* = \sigma_i$, but there are far more weakly nested sequences in general. When $\tau_i(y_i) \neq y_i^*$ for $1 \leq i < n-1$ (so for most weakly nested sequences, but not for the one in the example above), note that the elements σ_i determine uniquely the nested sequence of intervals entering in their definition, hence the σ_i^* as well.

Let D be a crystalline (φ, Γ) -module of rank n over \mathcal{R}_L satisfying (i). Any refinement \mathcal{F} of D determines an ordering $(\varphi_1, \dots, \varphi_n)$ of the eigenvalues of φ on $\mathcal{D}_{\text{crys}}(D)$ defined by $\det(\varphi|_{\text{Fil}_i(\mathcal{D}_{\text{crys}}(D))}) = \prod_{j=1}^i \varphi_j$, which of course determines \mathcal{F} , so we shall also write $\mathcal{F} = (\varphi_1, \dots, \varphi_n)$. In particular \mathfrak{S}_n acts on the set of refinements of D , via $\sigma((\varphi_i)) = (\varphi_{\sigma^{-1}(i)})$, this action being simply transitive. We say that a sequence of refinements $\mathcal{F}_0, \dots, \mathcal{F}_{n-1}$ is *nested* if there exists a nested sequence of permutations σ_i of \mathfrak{S}_n such that $\mathcal{F}_i = \sigma_i(\mathcal{F}_0)$ for each i . We say that a sequence of refinements $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_1^*, \dots, \mathcal{F}_{n-1}, \mathcal{F}_{n-1}^*$ is *weakly nested* if there exists a weakly nested sequence of permutations σ_i, σ_i^* of \mathfrak{S}_n such that $\mathcal{F}_i = \sigma_i(\mathcal{F}_0)$ and $\mathcal{F}_i^* = \sigma_i^*(\mathcal{F}_0)$ for each $i = 1, \dots, n-1$.

THEOREM 3.20. – *Let D be a crystalline (φ, Γ) -module of rank n over \mathcal{R}_L satisfying (i) and (ii). If $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_1^*, \dots, \mathcal{F}_{n-1}, \mathcal{F}_{n-1}^*$ is a weakly nested sequence of non-critical refinements of D , then $\mathfrak{t} = \sum_{i=0}^{n-1} \mathfrak{t}_{\mathcal{F}_i}$.*

In particular, if $\mathcal{F}_0, \dots, \mathcal{F}_{n-1}$ is a nested sequence of non-critical refinements of D , then $\mathfrak{t} = \sum_{i=0}^{n-1} \mathfrak{t}_{\mathcal{F}_i}$.

Note that this result implies Theorem 3.19. Let us settle first the underlying representability questions (property (iii)).

LEMMA 3.21. – *Let D be a crystalline (φ, Γ) -module satisfying (i). If D admits a weakly nested sequence of non-critical refinements, then (iii) and (iv) hold. Any subquotient of a generic (φ, Γ) -module is generic.*

Proof. – If D admits a non critical refinement, then (iv) is easily checked. By (i), $\text{End}_{(\varphi, \Gamma)}(D) \subset \text{End}_{L[\varphi]}(\mathcal{D}_{\text{crys}}(D)) \simeq L^n$ (a diagonal L -algebra), so (iii) for D is equivalent to the fact that D is not the direct sum of two (φ, Γ) -modules over \mathcal{R}_L .

Assume that $D = D_1 \oplus D_2$ and let $k_1 < k_2 < \dots < k_n$ be the Hodge-Tate weights of D . This defines a partition $\{1, 2, \dots, n\} = A \amalg B$ by $i \in A$ if, and only if, k_i is a

Hodge-Tate weight of D_1 . Let $(\varphi_1, \dots, \varphi_n)$ be a non-critical refinement of D . Observe that for all i , φ_i is an eigenvalue of φ on $\mathcal{D}_{\text{crys}}(D_1)$ if and only if $i \in A$: indeed, if $(\text{Fil}_i(D))$ is the triangulation of D associated to \mathcal{F} , then for all j the jump of the Hodge filtration on the line $\mathcal{D}_{\text{crys}}(D/\text{Fil}_{j-1}(D))^{\varphi=\varphi_j}$ is exactly k_j as \mathcal{F} is non-critical, and we are done by induction on j . As a consequence, if \mathcal{F} and $\sigma(\mathcal{F})$ are both non-critical, then $\sigma(A) \subset A$. If σ is a cycle, $A = \emptyset$ or $A = \{1, \dots, n\}$, which concludes the proof of the first part. The second part follows at once. \square

The remainder of this section is devoted to the proof of Theorem 3.20. Recall that for each non critical refinement \mathcal{F} we have an inclusion

$$\mathfrak{t}_{\text{crys}} \subset \mathfrak{t}_{\mathcal{F}} \subset \mathfrak{t},$$

with respective dimensions $\frac{n(n-1)}{2} + 1$, $\frac{n(n+1)}{2} + 1$ and $n^2 + 1$. So $\mathfrak{t}_{\mathcal{F}}/\mathfrak{t}_{\text{crys}}$ is a subspace of dimension n inside $\mathfrak{t}/\mathfrak{t}_{\text{crys}}$, which has dimension $\frac{n(n+1)}{2}$. The idea of the proof is to show that such an \mathcal{F} being given, the tangent space of a suitable paraboline deformation functor of type $(n-1, 1)$ or $(1, n-1)$ is in direct sum with $\mathfrak{t}_{\mathcal{F}}$ modulo $\mathfrak{t}_{\text{crys}}$, and then argue by induction on $\text{rk}_{\mathcal{R}_L}(D)$ in the $(n-1) \times (n-1)$ square. Note that even if we start from an étale D , we will lose this property in the induction process, which justifies the generality adopted in §3.9.

In the following lemma, D is a crystalline (φ, Γ) -module over \mathcal{R}_L satisfying (i), (ii), (iii) and (iv). Fix \mathcal{F} a refinement of D and set $c = (1, 2, \dots, n) \in \mathfrak{S}_n$.

LEMMA 3.22. – *If \mathcal{F} and $c(\mathcal{F})$ are non critical, and if ϕ is the last element of \mathcal{F} ,*

$$\mathfrak{t} = \mathfrak{t}_{\mathcal{F}} + \mathfrak{t}_{\phi\downarrow} \quad \text{and} \quad \mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}_{\phi\downarrow} = \mathfrak{t}_{\text{crys}}.$$

Similarly, if \mathcal{F} and $c^{-1}(\mathcal{F})$ are non critical, and if ϕ is the first element of \mathcal{F} , then $\mathfrak{t} = \mathfrak{t}_{\mathcal{F}} + \mathfrak{t}_{\phi\uparrow}$ and $\mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}_{\phi\uparrow} = \mathfrak{t}_{\text{crys}}$.

Proof. – The second part follows from the first one by duality by Remark 3.17, so we focus on the first part. First of all, by Prop. 3.15 and 3.16, $\mathfrak{t}_{\text{crys}} \subset \mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}_{\phi\downarrow}$ and

$$(3.7) \quad \dim(\mathfrak{t}_{\mathcal{F}}/\mathfrak{t}_{\text{crys}}) + \dim(\mathfrak{t}_{\phi\downarrow}/\mathfrak{t}_{\text{crys}}) = \frac{n(n+1)}{2} = \dim(\mathfrak{t}/\mathfrak{t}_{\text{crys}}).$$

(Prop. 3.15 applies as $\mathcal{D}_{\text{crys}}(D)^{\varphi=\phi}$ is non-critical, since ϕ is the first eigenvalue of the non critical refinement $c(\mathcal{F})$.) Thus it only remains to show that $\mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}_{\phi\downarrow} \subset \mathfrak{t}_{\text{crys}}$. As \mathcal{F} is non-critical, it is enough to show that

$$(3.8) \quad \mathfrak{X}_{D, \mathcal{F}} \cap \mathfrak{X}_{D, \phi\downarrow} \subset \mathfrak{X}_{D, \text{Sen}}$$

by Prop. 3.15. Fix an object A of \mathcal{C} , let $(D_A, \text{Fil}_i, \pi) \in \mathfrak{X}_{D, \mathcal{F}}(A)$, and assume that $(D_A, \pi) \in \mathfrak{X}_{D, \phi}(A)$ as well. By property (iv), we may write

$$P_{\text{Sen}/A}(D_A) = \prod_{i=1}^n (T - \kappa_i) \in A[T],$$

where $\kappa_i \in A$ lifts k_i . Choose $\tilde{\phi} \in A^*$ lifting ϕ such that $\mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}}$ is free of rank 1 over A and let $0 \leq j \leq n-1$. As ϕ is the last element of \mathcal{F} , $\mathcal{D}_{\text{crys}}(\text{Fil}_j(D))^{\varphi=\phi} = 0$ and an immediate dévissage shows that $\mathcal{D}_{\text{crys}}(\text{Fil}_j(D_A))^{\varphi=\tilde{\phi}} = 0$. Applying the left exact

functor $\mathcal{D}_{\text{crys}}(-)^{\varphi=\tilde{\phi}}$ to the exact sequence $0 \rightarrow \text{Fil}_j(D_A) \rightarrow D_A \rightarrow D_A/\text{Fil}_j(D_A) \rightarrow 0$ of (φ, Γ) -modules over \mathcal{R}_A , we obtain an $A[\varphi]$ -linear injection

$$(3.9) \quad \mathcal{D}_{\text{crys}}(D_A)^{\varphi=\tilde{\phi}} \hookrightarrow \mathcal{D}_{\text{crys}}(D_A/\text{Fil}_j(D_A))^{\varphi=\tilde{\phi}}.$$

On the other hand, as ϕ is an eigenvalue of multiplicity 1 in $\mathcal{D}_{\text{crys}}(D)$, another immediate dévissage shows that the length of the A -module $\mathcal{D}_{\text{crys}}(D_A/\text{Fil}_j(D_A))^{\varphi=\tilde{\phi}}$ is $\leq \text{length}(A)$, thus (3.9) is an isomorphism. As $c(\mathcal{F})$ is non critical, the Hodge-Tate weights of $D/\text{Fil}_j(D)$ are k_{j+1}, \dots, k_n and $\mathcal{D}_{\text{crys}}(D/\text{Fil}_j(D))^{\varphi=\phi}$ is non-critical in $\mathcal{D}_{\text{crys}}(D/\text{Fil}_j(D))$. Thus Prop. 3.16 applies, and shows that for the obvious π' we have $(D_A/\text{Fil}_j(D_A), \pi') \in \mathfrak{X}_{D/\text{Fil}_j(D), \phi \downarrow}(A)$. In particular,

$$P_{\text{Sen}/A}(D_A/\text{Fil}_j(D_A))(k_{j+1}) = 0.$$

As $P_{\text{Sen}/A}(D_A/\text{Fil}_j(D_A)) = \prod_{i=j+1}^n (T - \kappa_i)$ by Lemma 3.14, we conclude that $\kappa_{j+1} = k_{j+1}$ for all $0 \leq j \leq n-1$: D_A is Hodge-Tate, and we are done. \square

We now prove Theorem 3.20 by induction on $n = \text{rk}_{\mathcal{R}_L} D$. When $n = 1$ the theorem is obvious as $\mathfrak{X}_{D, \mathcal{F}} = \mathfrak{X}_D$, so assume $n > 1$. Let $\mathcal{F}_0, \mathcal{F}_1, \mathcal{F}_1^*, \dots, \mathcal{F}_{n-1}, \mathcal{F}_{n-1}^*$ be a weakly nested sequence of non-critical refinements of D . Applying the above lemma to $\mathcal{F} = \mathcal{F}_0$, we obtain a fortiori

$$(3.10) \quad \mathfrak{t} = \mathfrak{t}_{\mathcal{F}_0} + \mathfrak{t}_{\mathcal{F}}$$

where either $\mathcal{P} = \mathcal{P}_{\phi}$ and ϕ is the last element of \mathcal{F}_0 (case $\sigma_1^* = c$), or $\mathcal{P} = \mathcal{P}^{\phi}$ and ϕ is the first element of \mathcal{F}_0 (case $\sigma_1^* = c^{-1}$). Set $D' = D/D_{\phi}$ in the first case and $D' = D^{\phi}$ in the second case. By definition, for $i = 1, \dots, n-1$ the refinements \mathcal{F}_i and \mathcal{F}_i^* have the form (ϕ, \mathcal{F}'_i) and (ϕ, \mathcal{F}'_i^*) (resp. (\mathcal{F}'_i, ϕ) and (\mathcal{F}'_i^*, ϕ)) in the first case (resp. second case). Moreover,

$$\mathcal{F}'_1, \mathcal{F}'_2, \mathcal{F}'_2^*, \dots, \mathcal{F}'_{n-1}, \mathcal{F}'_{n-1}^*$$

is a weakly nested sequence of refinements of D' . Each \mathcal{F}'_i (resp. \mathcal{F}'_i^*) is non-critical for D' as \mathcal{F}_i (resp. \mathcal{F}_i^*) is non critical for D . As D' obviously still satisfies (i) and (ii), we obtain by induction:

$$(3.11) \quad \mathfrak{t}' = \sum_{i=1}^{n-1} \mathfrak{t}'_{\mathcal{F}'_i},$$

where $\mathfrak{t}'_* = \mathfrak{X}_{D', *}(L[\varepsilon])$. Note that for $i = 1, \dots, n-1$ we have $\mathfrak{X}_{D, \mathcal{F}_i} \subset \mathfrak{X}_{D, \mathcal{F}}$. By Prop. 3.7 and property (ii), the natural map $\mathfrak{X}_{D, \mathcal{F}} \rightarrow \mathfrak{X}_{D'}$ induces isomorphisms

$$(3.12) \quad \mathfrak{X}_{D, \mathcal{F}_i} = \mathfrak{X}_{D, \mathcal{F}} \times_{\mathfrak{X}_{D'}} \mathfrak{X}_{D', \mathcal{F}'_i}.$$

By (3.11) and (3.12) we have $\mathfrak{t}_{\mathcal{F}} = \sum_{i=1}^{n-1} \mathfrak{t}_{\mathcal{F}_i}$, and we are done by (3.10). \square

Another consequence of the proof of Lemma 3.22 is the following proposition.

PROPOSITION 3.23 (Transversality of trianguline deformation functors)

Assume that D satisfies (i), (ii) and let \mathcal{F} and \mathcal{F}' be two refinements of D . If \mathcal{F}' starts with the last element of \mathcal{F} , and if \mathcal{F} , $c(\mathcal{F})$ and \mathcal{F}' are non-critical, $\mathfrak{X}_{D, \mathcal{F}} \cap \mathfrak{X}_{D, \mathcal{F}'}$ is exactly the subfunctor of deformations of D which are crystalline up to a twist.

3.24. Some examples

Note that any (φ, Γ) -module of rank 2 satisfying (i), (ii), (iii) and (iv) is generic. The following other example plays some role in the main global result of this paper.

EXAMPLE 3.25. – Assume that $D = D_{\text{rig}}(V)$ for some representation V of dimension 3 over L satisfying (i), (iii) and (iv). Then D admits a nested sequence of non-critical refinements.

Proof. – Let X be the set of Frobenius eigenvalues of φ on $D_{\text{crys}}(V)$, and let $k_1 < k_2 < k_3$ be the Hodge-Tate weights of V . A line $L \subset D_{\text{crys}}(V)$ (resp. a plane P) is critical if $L \subset \text{Fil}^{k_2}(D_{\text{crys}}(V))$ (resp. $P \supset \text{Fil}^{k_3}(D_{\text{crys}}(V))$).

Assume first that there is some $\phi \in X$ such that the line $L_\phi := D_{\text{crys}}(V)^{\varphi=\phi}$ is critical. Then we claim that there is a unique such ϕ in X and that no φ -stable plane $P \subset D_{\text{crys}}(V)$ is critical. In particular, the critical refinements of D are exactly the two ones starting with ϕ . This implies the lemma in this case as each refinement of the nested sequence

$$(\phi', \phi, \phi''), (\phi'', \phi', \phi), (\phi'', \phi, \phi')$$

is non-critical. If there is no critical line L_ϕ as above, then either D is generic or there is a φ -stable critical plane $L_{\phi'} \oplus L_{\phi''}$. It follows from the preceding claim applied to V^* that in the latter case such a plane is unique, hence the critical refinements of D are the ones ending by ϕ and

$$(\phi', \phi, \phi''), (\phi, \phi'', \phi'), (\phi'', \phi, \phi')$$

is a nested sequence of non-critical refinements, and we are done again.

Let us check the claim. As V is indecomposable, $D_{\text{crys}}(V)$ is not the direct sum of an admissible line and of an admissible plane. Write $X = \{\phi, \phi', \phi''\}$ and do not assume anything about ϕ for the moment. If $L_\phi \oplus L_{\phi'} = \text{Fil}^{k_2}(D_{\text{crys}}(V))$, the weak admissibility property of $D_{\text{crys}}(V)$ implies that $L_{\phi''}$ and $L_\phi \oplus L_{\phi'}$ are admissible, which is absurd. Similarly, $L_\phi \neq \text{Fil}^{k_3}(D_{\text{crys}}(V))$ otherwise L_ϕ and $L_{\phi'} \oplus L_{\phi''}$ would be admissible. Assume now that L_ϕ is critical. If P is a φ -stable critical plane, then $P \neq \text{Fil}^{k_2}(D_{\text{crys}}(V))$ and P does not contain L_ϕ by what we just proved, so $P = L_{\phi'} \oplus L_{\phi''}$ and the jump indices of the Hodge filtration on P are k_1 and k_3 . But weak admissibility implies that L_ϕ and P are admissible, and we are done. \square

EXAMPLE 3.26. – Let D be a crystalline (φ, Γ) -module of rank 2 over \mathcal{R}_L satisfying (i), (iii) and such that the ratio r between its two Frobenius eigenvalues satisfies $r^i \neq 1$ for $i = 1, \dots, n$.

Then $\text{Sym}_{\mathcal{R}_L}^n(D)$ satisfies (i) and all of its refinements are non critical.

Proof. – We leave as an exercise to the reader to check that the statement follows from the following claim: if $I \subset \{0, \dots, n\}$ has i elements and if we have a polynomial identity $Q(x)(1+x)^i = \sum_{i \in I} a_i x^i$ where $a_i \in L$ and $Q \in L[x]$ has degree $\leq n+1-i$, then $Q = 0$. To check the claim, consider the i successive derivatives at -1 of the right hand side. This is a linear system in the unknown $(a_i)_{i \in I}$ whose determinant (in absolute value) is the Vandermonde determinant over the i elements of I . \square

Let us end this paragraph by introducing some *regularity conditions* that will arise for technical reasons in the subsequent global applications. Let D be a crystalline (φ, Γ) -module over \mathcal{R}_L satisfying (i). We say that a refinement $\mathcal{F} = (\varphi_1, \dots, \varphi_n)$ is *regular* if for all $1 \leq j \leq n$, the element $\varphi_1 \varphi_2 \cdots \varphi_j$ is a simple root of the polynomial $\prod_I (T - \prod_{i \in I} \varphi_i)$, the product being over the $I \subset \{1, \dots, n\}$ with $|I| = j$.⁽⁹⁾

DEFINITION 3.27. – *We say that D is weakly generic and regular if it satisfies (i), (ii), and if it possesses a weakly nested sequence of non-critical refinements $\{\mathcal{F}_i, \mathcal{F}_i^*\}$ such that \mathcal{F}_i is regular for each $i = 0, \dots, n-1$. If $D = D_{\text{rig}}(V)$, we say that V is regular if D is.*

The condition of being weakly generic and regular is Zariski-open and Zariski-dense in the algebraic variety of all filtered φ -modules of dimension n over L (whose filtration admits n jumps). This is actually still true if we restrict to filtered φ -modules satisfying a self-duality condition (orthogonal or symplectic, possibly with a similitude factor, in which cases the filtration is fixed by a Lagrangian in L^n). Indeed, set $k = \lfloor n/2 \rfloor$, let $(X_i)_{i=1, \dots, k}$ be some indeterminates and consider the sequence $s = (X_1, \dots, X_k, X_k^{-1}, \dots, X_1^{-1})$ if $n = 2k$, or $s = (X_1, \dots, X_k, 1, X_k^{-1}, \dots, X_1^{-1})$ if $n = 2k+1$. Observe that s is regular (for the obvious definition), moreover each permutation of s whose first k terms do not contain both some X_i and its inverse X_i^{-1} is regular as well (for instance, there are $2^k k!$ such permutations when n is even). We conclude as we may find a weakly nested sequence of permutations of \mathfrak{S}_n , associated to the nested sequence of intervals $y_i = i$, whose σ_i have the property above: for instance take $\tau_i = (i, n-i, n)$ for all $i \leq k$ if n is even.

EXAMPLE 3.28. – *Let D be generic of rank 4 over \mathcal{R}_L and assume that it admits a refinement of the form $(\mu, \mu x, \mu x^2, \mu x^3)$ for some $\mu, x \in L^*$. If $x^j \neq 1$ for $1 \leq j \leq 4$, then D is weakly generic regular.*

Proof. – The refinement \mathcal{F}_0 of the statement, $\mathcal{F}_1 = (\mu x^3, \mu x, \mu, \mu x^2)$, $\mathcal{F}_2 = (\mu x^3, \mu x^2, \mu x, \mu)$ and $\mathcal{F}_3 = (\mu x^3, \mu x^2, \mu, \mu x)$ form a weakly nested sequence of regular refinements of D (associated to $y_i = i$). \square

4. The eigenvariety at non-critical tempered classical points

Our main goal in this section is to show that the eigenvarieties of $U(3)$ or even $U(n)$ are étale over the weight space at the (stable, tempered) non-critical classical points.

4.1. An infinitesimal classicity criterion

Let $n \geq 1$ be any integer and let U/F be a unitary group in n variables attached to E/F . We need to recall some definitions of the theory of p -adic automorphic forms for U . The reader may consult [12, §2] for a detailed discussion and complete proofs of the statements of this section, as well as [10] and [1, §7.3]. We assume that $U(F_v)$ is compact for all Archimedean places v and that $U(F_v) \simeq \text{GL}_n(\mathbb{Q}_p)$ if v divides p . In particular p splits in E and for each v in the set S_p of finite places of F dividing p we fix such an isomorphism, which defines as well a place \tilde{v} of E above p (so $\mathbb{Q}_p = F_v = E_{\tilde{v}}$).

⁽⁹⁾ When $n \leq 3$, this condition is a consequence of property (i) of D .

We need to introduce some group theoretic notations concerning $\mathrm{GL}_n(\mathbb{Q}_p)^{S_p}$, that we shall view as an algebraic group over \mathbb{Q}_p . Let B be its upper triangular Borel subgroup, T its diagonal torus, $X^*(T)$ the algebraic characters of T , I the subgroup of $\mathrm{GL}_n(\mathbb{Z}_p)^{S_p}$ whose elements are upper triangular modulo p , \overline{N}_0 the subgroup of lower triangular elements of I , $T^0 = T \cap I$, $T^- \subset T$ the submonoid whose elements t satisfy $t^{-1}\overline{N}_0 t \subset \overline{N}_0$, and $M = IT^-I$ (it is a submonoid of $\mathrm{GL}_n(\mathbb{Q}_p)^{S_p}$).

For each B -dominant weight $\chi \in X^*(T)$, let W_χ be the irreducible algebraic \mathbb{Q}_p -representation of $\mathrm{GL}_n(\mathbb{Q}_p)^{S_p}$ with highest weight χ . For any $\chi \in X^*(T)$, let \mathcal{C}_χ be the standard analytic principal series of the Iwahori subgroup I . It may be defined as follows. If J is a finite set, a function $f : \mathbb{Z}_p^J \rightarrow \mathbb{Q}_p$ is said analytic if $f((x_j))$ belongs to the Tate algebra $\mathbb{Q}_p\langle (x_j)_j \rangle$. Let J be the set of triples (i, j, v) with $1 \leq i < j \leq n$ and $v \in S_p$. Identifying \mathbb{Z}_p^J with \overline{N}_0 via the bijection $(x_{(i,j,v)}) \mapsto (px_{i,j})_v$, we obtain a notion of analytic function on \overline{N}_0 and set:

$$\mathcal{C}_\chi = \left\{ f : IB \longrightarrow \mathbb{Q}_p, \ f(xb) = \chi(b)f(x) \ \forall x \in IB, \ b \in B, \right. \\ \left. f|_{\overline{N}_0} \text{ is analytic.} \right\}.$$

The product in $\mathrm{GL}_n(\mathbb{Q}_p)^{S_p}$ induces an isomorphism $\overline{N}_0 \times B \xrightarrow{\sim} IB$. Moreover, as $t^{-1}\overline{N}_0 t \subset \overline{N}_0$ for $t \in T^-$, we have $M^{-1}IB \subset IB$, thus left translations $(m.f)(x) := f(m^{-1}x)$ define a representation of M on \mathcal{C}_χ . When χ is dominant, and if v is a highest weight vector in W_χ , the map $\varphi \mapsto (g \mapsto \varphi(gv))$ defines a $\mathbb{Q}_p[M]$ -equivariant embedding $W_\chi^\vee \rightarrow \mathcal{C}_\chi$, whose image is the subspace of polynomial elements in \mathcal{C}_χ .

The spaces of p -adic automorphic forms for U are built from these representations \mathcal{C}_χ . Let K be any compact open subgroup of $U(\mathbb{A}_{F,f})$ of the form $K = I \times K^{S_p}$. For any $\mathbb{Q}_p[M]$ -module W , consider the \mathbb{Q}_p -vector space

$$F(W) := \left\{ f : U(F) \backslash U(\mathbb{A}_{F,f}) \longrightarrow W, \right. \\ \left. f(gk) = (\prod_{v|p} k_v)^{-1} f(g), \ \forall g \in U(\mathbb{A}_{F,f}), \ \forall k \in K. \right\}.$$

This defines an exact functor from $\mathbb{Q}_p[M]$ -modules to \mathbb{Q}_p -vector spaces. Moreover, $F(W)$ is in a natural way a module over the Atkin-Lehner algebra \mathfrak{A}^- which is the \mathbb{Q} -subalgebra of the Iwahori Hecke-algebra of $\mathrm{GL}_n(\mathbb{Q}_p)^{S_p}$ relative to I , consisting of functions with support in M . Recall that the map $T^-/T^0 \rightarrow \mathfrak{A}^-$, $t \mapsto \mathbb{1}_{ItI}$, is a multiplicative homomorphism inducing an isomorphism

$$\mathbb{Q}[T^-/T^0] \xrightarrow{\sim} \mathfrak{A}^-$$

(in particular \mathfrak{A}^- is commutative). Using this latter isomorphism we will view elements of T^- as elements of \mathfrak{A}^- . Last but not least, $F(W)$ also admits a natural structure of module over the Hecke \mathbb{Q} -algebra of $(U(\mathbb{A}_{F,f}^{S_p}), K^{S_p})$ that commutes with \mathfrak{A}^- . We fix a commutative subring \mathcal{H} of this latter Hecke-algebra that contains the spherical Hecke-algebra for almost all finite primes v of F .

Let $\chi \in X^*(T)$. The space of p -adic analytic automorphic forms of U of weight χ and level K is the space $F(\mathcal{C}_\chi)$. It is a \mathbb{Q}_p -Banach space in a natural way, on which the Hecke-operators $\mathfrak{A}^- \otimes \mathcal{H}$ act continuously. Moreover, if $T^{--} \subset T^-$ denotes the submonoid whose elements t are such that $t^{-1}\overline{N}_0 t$ vanishes mod p , then any $t \in T^{--}$ acts compactly on $F(\mathcal{C}_\chi)$.

When χ is dominant, $F(\mathcal{C}_\chi)$ contains as $\mathfrak{A}^- \otimes \mathcal{H}$ -submodule the subspace $F(W_\chi^\vee)$ of automorphic forms of weight χ and level K , often called the subspace of *classical* p -adic

automorphic forms. If we fix a $\iota = (\iota_p, \iota_\infty)$ as in §1.7, this subspace is a canonical \mathbb{Q}_p -structure of the complex $\mathbb{Q}[U(\mathbb{A}_{F,f})//K]$ -module

$$(4.13) \quad \bigoplus_{\pi} m(\pi) \pi_f^K,$$

where π varies in the set of irreducible automorphic representations of $U(\mathbb{A}_F)$ such that $\pi_\infty \simeq W_\chi$, and with finite multiplicity $m(\pi)$ in $L^2(U(F) \backslash U(\mathbb{A}_F))$.

We shall also need to use the global base-change from U/F to GL_n/E . It is currently known in the following cases:

- $n \leq 3$ (Rogawski) or,
- there is a finite place v such that $U(F_v)$ is the group of invertible elements of a central division algebra over F_v (Clozel-Labesse, Harris-Labesse) or,
- $[F : \mathbb{Q}] \geq 2$ (Labesse),

and we definitely assume that we are in one of these three cases. Thanks to [38] and [15, 24], it follows that for any automorphic representation π of U there is a semi-simple continuous Galois representation

$$\rho_\pi : G_E \longrightarrow \mathrm{GL}_n(\overline{\mathbb{Q}_p}),$$

which is unramified above any finite place $v \notin S_p$ of F such that π_v and U/F_v are unramified, and which is characterized by the following property: for any such v which splits⁽¹⁰⁾ in E the characteristic polynomial of $\mathrm{Frob}_{\tilde{v}}$ coincides (via ι) with the characteristic polynomial of the Langlands conjugacy class of $\pi_v| \cdot |^{\frac{1-n}{2}}$. It has the following extra properties:

- $\rho_\pi^{\vee,c} \simeq \rho_\pi(n-1)$ (note that n is not necessarily odd in this paragraph).
- For each finite place $v \notin S_p$ of F such that $U(F_v) \xrightarrow{\sim}_{\tilde{v}} \mathrm{GL}_n(F_v)$ the Frobenius semi-simplification of $\rho_{\pi,\tilde{v}}$ corresponds to $\iota_p \iota_\infty^{-1} \pi_v| \cdot |^{\frac{1-n}{2}}$ via the local Langlands correspondence ([38],[15, 24],[9]).
- If $v \in S_p$ and π_v is unramified, then ρ_π is crystalline at \tilde{v} . Its Hodge-Tate weights are related to Π_∞ via ι and the usual recipe, and the characteristic polynomial of its crystalline Frobenius is the characteristic polynomial of the Langlands conjugacy class of $\pi_v| \cdot |^{\frac{1-n}{2}}$ (of course via ι , see [14]).

Before stating the *infinitesimal classicity criterion* we need to recall the relation between refinements and eigenvectors of \mathfrak{A}^- . Let $f \neq 0 \in F(W_\chi^\vee) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$ be a common eigenvector for all the elements of $\mathfrak{A}^- \otimes \mathcal{H}$. Let π be any automorphic representation of U with $\pi_\infty \simeq W_\chi$ and which occurs as a summand of the representation generated by $\iota_\infty \iota_p^{-1} f$; the Galois representation $\rho_f := \rho_\pi$ does not depend on the choice of π by Cebotarev's theorem. According to the recipe described in [1, §6.4], the action of \mathfrak{A}^- on f determines for each $v \in S_p$ a canonical ordering $(\varphi_{i,v})_{i=1}^n$ of the eigenvalues of the Langlands conjugacy class associated to the representation $\pi_v| \cdot |^{\frac{1-n}{2}}$ (which has Iwahori invariants by construction). When $\varphi_{i,v} \varphi_{j,v}^{-1} \neq p$ for all $i \neq j$ then π_v is unramified, so $\rho_{f,\tilde{v}}$ is crystalline. If furthermore $\varphi_{i,v} \varphi_{j,v}^{-1} \neq 1$ for all $i \neq j$, then $(\varphi_{i,v})_{i=1}^n$ defines a refinement of $\rho_{f,\tilde{v}}$ in the sense of §3.9 (see the discussion preceding Thm 3.20).

⁽¹⁰⁾ So $U(F_v) \simeq \mathrm{GL}_n(F_v)$ and the choice of such an isomorphism defines a unique place \tilde{v} of E above v .

PROPOSITION 4.2. – *Let $f \in F(W_\chi^\vee) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$ be an eigenvector for all the elements of $\mathfrak{A}^- \otimes \mathcal{H}$. Assume that for each $v \in S_p$, the sequence $(\varphi_{i,v})_{i=1}^n$ defined above satisfies $\varphi_{i,v}\varphi_{j,v}^{-1} \neq 1, p$ for any $i \neq j$, and defines a non-critical, regular, refinement of $\rho_{f,\widetilde{v}}$.*

Then the generalized $\mathfrak{A}^- \otimes \mathcal{H}$ -eigenspace of f inside $F(\mathcal{C}_\chi) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$ is included in $F(W_\chi^\vee) \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}$.

When $(\varphi_{i,v})_{i=1}^n$ is numerically non-critical, this result follows from the standard classicity criterion (compare [1, Rem. 2.4.6] with [12, Thm. 1.6 (vi)]). However, this simple case is not enough for our purpose in this paper, and the general case is much deeper. It is the analogue of the classical fact that an ordinary modular eigenform which is in the image of the theta map is split at p (Mazur-Wiles).

The first ingredient of the proof is the following result of Jones, which is a (locally) analytic version of the Bernstein-Gelfand-Gelfand resolution, and plays the role of the theta map in the context of p -adic modular forms.

LEMMA 4.3 ([26, Thm. 29]). – *Assume that $\chi \in X^*(T)$ is dominant. There exists an exact sequence of $\mathbb{Q}_p[M]$ -modules:*

$$0 \longrightarrow W_\chi^\vee \longrightarrow \mathcal{C}_\chi \longrightarrow \prod_{l(\sigma)=1} \mathcal{C}_{\sigma(\chi+\rho)-\rho}.$$

In this statement, σ is an element in the Weyl group $\mathfrak{S} = \mathfrak{S}_n^{S_p}$ of T , ρ is the half-sum of the positive roots (this is not quite an element of $X^*(T)$, but $\sigma(\chi + \rho) - \rho$ is anyway). By exactness of the functor F we obtain an exact sequence of $\mathfrak{A}^- \otimes \mathcal{H}$ -modules $0 \rightarrow F(W_\chi^\vee) \rightarrow F(\mathcal{C}_\chi) \rightarrow \prod_{l(\sigma)=1} F(\mathcal{C}_{\sigma(\chi+\rho)-\rho})$. Let $\psi : \mathfrak{A}^- \otimes \mathcal{H} \rightarrow \overline{\mathbb{Q}_p}$ be the \mathbb{Q} -algebra homomorphism defined by $f : a(f) = \psi(a)f$ for all $a \in \mathfrak{A}^- \otimes \mathcal{H}$. If E is any $\mathbb{Q}_p \otimes \mathfrak{A}^- \otimes \mathcal{H}$ -module we denote by $E[\psi]$ the generalized eigenspace associated to ψ :

$$E[\psi] = \{e \in E \otimes_{\mathbb{Q}_p} \overline{\mathbb{Q}_p}, \exists n \in \mathbb{N}, \forall a \in \mathfrak{A}^- \otimes \mathcal{H}, (a - \psi(a)\text{id})^n e = 0\}.$$

As f is classical, $\psi(T^-) \subset \overline{\mathbb{Q}_p}^*$, and by compactness of any $t \in T^{--}$ we obtain an exact sequence of finite dimensional $\overline{\mathbb{Q}_p}$ -vector spaces:

$$0 \longrightarrow F(W_\chi^\vee)[\psi] \longrightarrow F(\mathcal{C}_\chi)[\psi] \longrightarrow \prod_{l(\sigma)=1} F(\mathcal{C}_{\sigma(\chi+\rho)-\rho})[\psi].$$

We claim that for each simple reflection $\sigma = (i, i+1)_v$, we have $F(\mathcal{C}_{\sigma(\chi+\rho)-\rho})[\psi] = 0$. Indeed, it is enough to show that there is no $g \neq 0 \in F(\mathcal{C}_{\sigma(\chi+\rho)-\rho})[\psi]$ that is an eigenvector for $\mathfrak{A}^- \otimes \mathcal{H}$. Assume for a contradiction that there are such $i, v \in S_p$ and g .

As g and f share the same system of Hecke-eigenvalues under $\mathfrak{A}^- \otimes \mathcal{H}$, the form g is of finite slope and its associated p -adic Galois representation $\rho_g : G_E \rightarrow \text{GL}_n(\overline{\mathbb{Q}_p})$ coincides with ρ_f . However, the form f has weight χ whereas its “companion form” g has weight $\sigma(\chi + \rho) - \rho$. Let us denote by

$$k_{1,v} < k_{2,v} < \cdots < k_{n,v}$$

the Hodge-Tate weights of $\rho_{f,\widetilde{v}}$. They are related to the weight χ of f by the usual recipe, namely an identification $X^*(T) \simeq (\mathbb{Z}^n)^{S_p}$ making the dominant elements of $X^*(T)$ correspond to increasing sequences as above. According to the same recipe, the ordering of the

Hodge-Tate numbers of $\rho_{g,v}$ corresponding to $\sigma(\chi + \rho) - \rho$ is the same one except that $k_{i,v}$ and $k_{i+1,v}$ are interchanged. As the actions of \mathfrak{A}^- on g and f coincide, Kisin's result [29, §5] on the continuity of crystalline periods in refined families ensures that ρ_g has the following property⁽¹¹⁾:

$$(4.14) \quad \mathrm{Fil}^{k_{1,v}+k_{2,v}+\cdots+k_{i-1,v}+k_{i+1,v}}(D_{\mathrm{crys}}(\Lambda^i \rho_{g,v})^{\varphi=\varphi_{1,v}\varphi_{2,v}\cdots\varphi_{i,v}}) \neq 0.$$

As $(\varphi_{i,v})_{i=1}^n$ is regular, if $F_i = \bigoplus_{j=1}^i D_{\mathrm{crys}}(\rho_{g,v})^{\varphi=\varphi_{j,v}} \subset D_{\mathrm{crys}}(\rho_{g,v})$ we have

$$D_{\mathrm{crys}}(\Lambda^i \rho_{g,v})^{\varphi=\varphi_{1,v}\varphi_{2,v}\cdots\varphi_{i,v}} = \Lambda^i(F_i) \subset \Lambda^i D_{\mathrm{crys}}(\rho_{g,v}).$$

On the other hand, $(\varphi_{i,v})_{i=1}^n$ is non critical so the jumps of the filtration on F_i induced by the Hodge filtration on $D_{\mathrm{crys}}(\rho_{g,v})$ are $k_{1,v}, k_{2,v}, \dots, k_{i,v}$, which contradicts (4.14). \square

4.4. Non critical classical points are étale over the weight space

The preceding result, combined with some properties of automorphic forms, allows to show that eigenvarieties are étale over the weight space at non critical classical points.

Let $\ell \neq p$ be a prime and E_ℓ a finite extension of \mathbb{Q}_ℓ . We denote by W_{E_ℓ} the Weil-group of E_ℓ and $I_{E_\ell} \subset W_{E_\ell}$ its inertia subgroup. If $\rho : W_{E_\ell} \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$ is continuous, it admits an associated Frobenius semi-simple Weil–Deligne representations (r, N) . Recall that we have fixed some embeddings ι_p, ι_∞ . We say that ρ is defined over $\overline{\mathbb{Q}}$ if r is. In this case, the local Langlands correspondence associates to (r, N) (and ι_p, ι_∞) a canonical irreducible smooth complex representation of $\mathrm{GL}_n(E_\ell)$. For two continuous representations $\rho, \rho' : W_{E_\ell} \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}}_p)$, with associated Weil–Deligne representations (r, N) and (r', N') , we write $\rho \prec_I \rho'$ if $r|_{I_{E_\ell}} \simeq r'|_{I_{E_\ell}}$ and if ρ has less monodromy than ρ' in the following sense: for each irreducible $\overline{\mathbb{Q}}_p$ -representation τ of I_{E_ℓ} , the nilpotent conjugacy class of the monodromy operator is greater on the τ -isotypic component of r' than on the τ -isotypic component of r (for the dominance ordering, the conjugacy class of 0 being by convention the smallest one). Recall that if $r = \sum_{i=1}^s r_i$ and $r' = \sum_{j=1}^{s'} r'_j$ are irreducible decompositions of r and r' , then $r|_{I_{E_\ell}} \simeq r'|_{I_{E_\ell}}$ if and only if $s = s'$ and, up to renumbering the r_i, r'_i is an unramified twist of r_i for each $i \in \{1, \dots, s\}$ (see, e.g., [12, Lemme 3.14 (i)]).

LEMMA 4.5. – *Let X be an irreducible affinoid over \mathbb{Q}_p , $Z \subset X(\overline{\mathbb{Q}}_p)$ a Zariski-dense subset, and let $\rho : W_{E_\ell} \rightarrow \mathrm{GL}_n(\mathcal{O}(X))$ be a continuous representation.*

- (i) *There is a Zariski-dense subset $Z' \subset Z$ such that $\rho_x \prec_I \rho_z$ and $\rho_z|_{I_{E_\ell}} \simeq \rho_{z'}|_{I_{E_\ell}}$ for all $z, z' \in Z'$ and $x \in X$.*

⁽¹¹⁾ To deduce this result from the literature, we may refer as follows to [1]. First, choose by Lemma 7.8.11 an affinoid neighborhood Ω of the point x corresponding to g in the eigenvariety of U of level K . There is a surjective alteration $\Omega' \rightarrow \Omega$ and a locally free \mathcal{O}_Ω -module \mathcal{M} with a continuous $\mathcal{O}_{\Omega'}$ -linear action of G_E which is generically semi-simple and whose trace is the pull-back of the natural family of pseudo-characters on Ω . Apply Theorem 3.3.3 to $\mathcal{O}_{\Omega'}^i(\mathcal{M})$ and any point $x' \in \Omega'$ above x , its assumptions are satisfied by Prop. 7.5.13 and §4.2.4. We obtain (4.14) with $\overline{\mathcal{M}}_{x'}$ instead of $\rho_{g,v}$. But $\rho_{g,v}$ is the semi-simplification of $\overline{\mathcal{M}}_{x'} \otimes \overline{\mathbb{Q}}_p$, so (4.14) follows from the left-exactness of the functor $\mathrm{Fil}^k(D_{\mathrm{crys}}(-)^{\varphi=\phi})$.

- (ii) For each $z \in Z$ assume that ρ_z is defined over $\overline{\mathbb{Q}}$ and that its Langlands correspondent π_z is tempered. Then for any $z, z' \in Z$, we have

$$\rho_z|_{I_{E_\ell}} \simeq \rho_{z'}|_{I_{E_\ell}} \quad \text{and} \quad \pi_z|_{\mathrm{GL}_n(\mathcal{O}_{E_\ell})} \simeq \pi_{z'}|_{\mathrm{GL}_n(\mathcal{O}_{E_\ell})}.$$

Proof. – Part (i) is [1, Prop. 7.8.19] (in the notations there, apply it to the tautological pseudo-character $T : \mathrm{GL}_n(\mathcal{O}(X)) \rightarrow \mathcal{O}(X)$ and to the morphism $W_{E_\ell} \rightarrow \mathrm{GL}_n(\mathcal{O}(X))$ of the statement). Let us check part (ii).

Let π and π' be two irreducible *tempered* representations of $\mathrm{GL}_n(E_\ell)$, with associated Weil–Deligne representations (r, N) and (r', N') . If $(r|_I, N)$ and $(r'|_I, N')$ are isomorphic, then π and π' are fully induced from discrete series Δ and Δ' of a same Levi subgroup of $\mathrm{GL}_n(E_\ell)$, and Δ and Δ' only differ by a unitary unramified twist (Langlands, Zelevinski). In particular, π and π' are isomorphic restricted to $\mathrm{GL}_n(\mathcal{O}_{E_\ell})$. As a consequence, the second assertion in part (ii) follows from the first one, that we check now. We may assume $z' \in Z'$. Moreover, by the inequality in part (i) it is enough to check that $\rho_z|_{I_{E_\ell}} \simeq \rho_{z'}|_{I_{E_\ell}}$ holds after a finite base change. As base change preserves temperedness, we may assume that for each z in Z , if (r_z, N_z) is the Weil–Deligne representation associated to ρ_z , then r_z is trivial over I_{E_ℓ} .

Let $\varphi \in W_{E_\ell}$ be a lift of a geometric Frobenius and let $P \in \mathcal{O}(X)[T]$ be the characteristic polynomial of $\rho(F)$. Up to replacing X by a finite covering and Z by its inverse image, we may assume that P splits in $\mathcal{O}(X)[T]$; let $F_1, \dots, F_n \in \mathcal{O}(X)$ be its roots. By part (i) the conjugacy class of N_z does not depend on $z \in Z'$, let us call it N ; it is determined by a partition $n_1 + n_2 + \dots + n_s = n$. For each $z \in Z'$, there is a renumbering F'_i of the F_i such that whenever $1 \leq j < n_1$ or $n_1 + n_2 + \dots + n_r + 1 \leq j < n_1 + n_2 + \dots + n_{r+1}$ we have $F'_j(z) = qF'_{j+1}(z)$, where q is the cardinal of the residue field of E_ℓ . For z in a Zariski-dense subset $Z'' \subset Z'$, this numbering will be the same, hence up to renumbering the F_i once and for all we may assume that if $1 \leq j < n_1$ or if $n_1 + n_2 + \dots + n_r + 1 \leq j < n_1 + n_2 + \dots + n_{r+1}$,

$$(4.15) \quad F_j(z) = qF_{j+1}(z) \quad \forall z \in Z''.$$

Thus $F_{j+1} = qF_j$ in $\mathcal{O}(X)$ (we may assume X is reduced) for any j as above, hence (4.15) holds as well for any $z \in Z$. But the temperedness of π_z and the inequality $N_z \prec N$ imply in turn that N_z is conjugate to N . (Note the funny interplay between the complex and p -adic sides in this proof.) \square

Assume now that E_ℓ is a quadratic extension of F_ℓ (ℓ is still prime to p) and let $\mathrm{U}(F_\ell)$ be the quasi-split unitary group in n variables attached to E_ℓ/F_ℓ . For $n \leq 3$, the standard base-change between $\mathrm{U}(F_\ell)$ and $\mathrm{GL}_n(E_\ell)$, conjectured by Langlands and proved by Rogaswki in [35, Prop. 11.4.1, Thm. 13.2.1], is an injection from the set of L -packets of $\mathrm{U}(F_\ell)$ to the irreducible smooth representations π of $\mathrm{GL}_n(E_\ell)$ such that $\pi^{\vee, w} \simeq \pi$ ($w \in W_{F_\ell} \backslash W_{E_\ell}$) and whose central character is trivial over F_ℓ^* if n is odd.

LEMMA 4.6. – We keep the assumptions of Lemma 4.5 and assume that $n \leq 3$.

For each $z \in Z$ assume that ρ_z is defined over $\overline{\mathbb{Q}}$ and that there is a tempered L -packet Π_z of $\mathrm{U}(F_\ell)$ whose base-change to $\mathrm{GL}_3(E_\ell)$ corresponds to ρ_z . Then for any compact open subgroup K_v of $\mathrm{U}(F_\ell)$ and any $z, z' \in Z$, we have

$$\left(\bigoplus_{\pi \in \Pi_z} \pi \right)_{|K_v} \simeq \left(\bigoplus_{\pi \in \Pi_{z'}} \pi \right)_{|K_v}.$$

If χ is a character of E_ℓ^* , we shall denote by χ^\perp the character $x \mapsto \chi(c(x))^{-1}$ where c generates $\text{Gal}(E_\ell/F_\ell)$.

Proof. – Assume $n = 3$. The lemma is obvious when the isomorphism class of ρ_z is constant when z varies in Z , as then Π_z does not depend on z . We claim that this occurs in particular when there is some $z \in Z$ such that $\rho_z \otimes \overline{\mathbb{Q}}_p$ is either irreducible, of the form $r \oplus \chi$ with r irreducible of dimension 2, or of the form $\chi_1 \oplus \chi_2 \oplus \chi_3$ where the χ_i are distinct characters such that $\chi_i^\perp = \chi_i$ for each i (the associated L -packet of $U(F_\ell)$ is discrete and has respectively 1, 2 or 4 elements by [35, §13]).

Indeed, in the first case part (i) shows that for any $x \in X(\overline{\mathbb{Q}}_p)$ there is an unramified character χ_x such that $\rho_x \otimes \overline{\mathbb{Q}}_p = (\rho_z \otimes \overline{\mathbb{Q}}_p) \otimes \chi_x$, and the self-duality like condition forces χ_x to vary in a finite set when x varies. This implies that for all $g \in W_{E_\ell}$, $\text{trace}(\rho(g))$ is *constant*, *i.e.* belongs to the biggest finite extension of \mathbb{Q}_p in the domain $\mathcal{O}(X)$, hence the statement. The second case is similar. In the last case, part (i) implies that $\text{trace}(\rho)$ factors through a pseudo-character of $W_{E_\ell}^{\text{ab}} = E_\ell^*$. We may find a normal affinoid Y , a finite map $Y \rightarrow X$, as well as continuous characters $\tilde{\chi}_i : W_{E_\ell} \rightarrow \mathcal{O}(Y)^*$ such that

$$(4.16) \quad \text{trace}(\rho) = \sum_{i=1}^3 \tilde{\chi}_i,$$

thus up to replacing X by Y and Z by its inverse image in Y we may assume that $X = Y$. The self-duality like condition and the assumption that the evaluation of the $\tilde{\chi}_i$ at z are all self- \perp and distinct imply that $\tilde{\chi}_i^\perp = \tilde{\chi}_i$ for each i . As they are constant on I_{E_ℓ} by part (i), we see again that there is only a finite number of possibilities for the $\rho_x \otimes \overline{\mathbb{Q}}_p$ with $x \in X$, and we conclude as before.

In the non-constant case, then for any z in Z we have

$$\rho_z^{\text{ss}} \otimes \overline{\mathbb{Q}}_p = \chi_z \oplus \nu_z \oplus \nu_z^\perp$$

for some characters $\chi_z, \nu_z : E_\ell^* \rightarrow \overline{\mathbb{Q}}_p^*$, with $\chi_z^\perp = \chi_z$. This implies that χ_z is trivial over F_ℓ^* (as $\chi_z \nu_z \nu_z^\perp$ is), hence is the base change to E_ℓ^* of a unique character $\tilde{\chi}_z$ of $U(1)$. This case is more subtle as there are in general four possible kinds of L -packets for Π_z , labelled as (1), (2), (6) and (8) in [35, p. 174] and to which we shall refer several times. We shall argue according to the generic monodromy of ρ :

(a) If the monodromy of ρ_z has nilpotent index 3, then from that list we see that Π_z is some twist of the Steinberg representation (case (8)). If this occurs for some $z \in Z'$, Lemma 4.5 (i) implies that this also occurs for all $z \in Z'$, so that Π_z is independent of $z \in Z'$ in this case (the twist alluded to above does not depend on z by an argument similar to the ones above). Assume it is so and fix $x \in Z$. This implies

$$(4.17) \quad \{\nu_x, \nu_x^\perp\} = \{\chi_x|\cdot|, \chi_x|\cdot|^{-1}\}.$$

We claim that Rogawski's list shows that Π_x is the twist of the Steinberg representation by $\tilde{\chi}_x$, so $\Pi_x = \Pi_z$ for any $z \in Z'$. This would conclude the proof in case (a). To check the claim note that property (4.17) of ν_x excludes the packets of type (1) (*irreducible principal series*), as well as those of type (6) (*l.d.s. L-packets*), so it only remains to exclude the type (2) (*endoscopic transfer of a twist of the Steinberg representation of $U(2) \times U(1)$, an L -packet with 2 elements*).

But the Weil–Deligne representation (r, N) , $r : W_{E_\ell} \rightarrow \mathrm{GL}_3(\mathbb{C})$, of the base-change of an L -packet of type (2) has a non trivial N and r is the sum of three characters

$$(4.18) \quad \chi \oplus \nu|\cdot|^{-1/2} \oplus \nu|\cdot|^{1/2}$$

where $\chi = \chi^\perp$, $\nu = \nu^\perp$, $\chi|_{F_\ell^*} = 1$ and $\nu|_{F_\ell^*} \neq 1$. These conditions contradict (4.17).

(b) If the monodromy of ρ_z has nilpotent index 2, the unique possibility from the list is that the L -packet Π_z is of type (2). If this occurs for some $z \in Z'$, an argument similar to the one in case (a) shows again that Π_z is independent of $z \in Z$.

(c) In the remaining case, ρ_z has no monodromy for all $z \in Z$, so we see from Rogawski's description that Π_z is the set of irreducible constituents of the parabolic induced representation $\mathrm{Ind}(\eta_z)$ of the character $\eta_z = \nu_z \chi_z^{-1} \times \tilde{\chi}_z$ of the diagonal torus $E_\ell^* \times U(1)$ of $U(F_\ell)$ (this covers the two possibilities (1) and (6)). Arguing as for (4.16), we may assume that χ_z and ν_z have been chosen so that they vary analytically when $z \in Z \subset X$. But then $\eta_z|_{\mathcal{O}_{E_\ell}^* \times U(1)}$ does not depend on $z \in Z$ by part (i), so we check at once that $\mathrm{Ind}(\eta_x)|_{K_v} = \mathrm{Ind}(\eta_z)|_{K_v}$ for all $x, z \in Z$.

This concludes the case $n = 3$. The case $n = 2$ is similar and only easier (use the description of L -packets given in [35, Chap. 11]) and $n = 1$ is trivial. \square

LEMMA 4.7. – Lemmas 4.5 and 4.6 still hold if ρ is replaced by the restriction to W_{E_ℓ} of a continuous pseudo-character $T : \mathrm{Gal}(\overline{E}/E) \rightarrow \mathcal{O}(X)$ of dimension n .

In this setting, if $x \in X(\overline{\mathbb{Q}_p})$ we mean by ρ_x the restriction to W_{E_ℓ} of the (unique) semi-simple representation $\mathrm{Gal}(\overline{E}/E) \rightarrow \mathrm{GL}_n(\overline{\mathbb{Q}_p})$ whose trace is the evaluation of T at x .

Proof. – Indeed, Lemmas 4.5 (ii) and 4.6 only relied on the existence of a ρ such that $\mathrm{trace}(\rho) = T$ via its Corollary Lemma 4.5 (i). But part (i) in this more general setting follows again from [1, Prop. 7.8.19]. \square

We now prove the main results of this section. Recall that p splits in E and that U is a unitary group in n variables attached to E/F as in §4.1. Recall also that we have fixed a compact open subgroup $K \subset U(\mathbb{A}_{F,f})$ that we assume now of the form $\prod_v K_v$. Let S be a finite set of finite primes of F containing the primes ramified in E and such that K_v is maximal hyperspecial for each $v \notin S$. Let \mathcal{E} be the p -adic eigenvariety of U of level K associated to the set S_p of all places above p and to the global Hecke-algebra \mathcal{H} unramified outside S .

THEOREM 4.8. – ($n \leq 3$) Let π be an automorphic representation of U which is unramified above p and such that $\pi^K \neq 0$. We assume that the base-change of π to $\mathrm{GL}_n(\mathbb{A}_E)$ is cuspidal and that for each $v \in S_p$ the crystalline Frobenius has n distinct eigenvalues on $D_{\mathrm{crys}}(\rho_{\pi, \widetilde{v}})$.

Let $\{\mathcal{F}_v\}$ be a collection of refinements of the $\rho_{\pi, \widetilde{v}}$ for $v \in S_p$ and let $x \in \mathcal{E}$ be the point associated to $(\pi, \{\mathcal{F}_v\})$. If \mathcal{F}_v is non-critical for each $v \in S_p$, then \mathcal{E} is étale at x over weight space.

This solves conjecture (CRIT) of [1] §7.6, for $n \leq 3$, to which we refer for a more complete discussion and motivations. The *non-critical* assumption is very important in this statement.

Proof. – For each point $y \in \mathcal{E}$ (resp. $y \in \mathcal{E}(\overline{\mathbb{Q}_p})$) we denote by $k(y)$ its residue field and by $\psi_y : \mathfrak{A}^- \otimes \mathcal{H} \rightarrow k(y)$ (resp. $\psi_y : \mathfrak{A}^- \otimes \mathcal{H} \rightarrow \overline{\mathbb{Q}_p}$) the evaluation at y of the structural ring homomorphism $\mathfrak{A}^- \otimes \mathcal{H} \rightarrow \mathcal{O}(\mathcal{E})$ at y .

By construction of the eigenvariety, we may find a connected affinoid neighborhood B of the weight χ_0 of x (as defined in the statement) in the weight space and a finite locally free $\mathcal{O}(B)$ -module M equipped with an $\mathcal{O}(B)$ -linear action of $\mathfrak{A}^- \otimes \mathcal{H}$ such that:

- (a) The affinoid spectrum V of $\text{Im}(\mathcal{O}(B) \otimes \mathfrak{A}^- \otimes \mathcal{H} \rightarrow \text{End}_{\mathcal{O}(B)}(M))$ is an affinoid neighborhood of x in \mathcal{E} . Denote by $\kappa : V \rightarrow B$ the natural map (so $\kappa(x) = \chi_0$).
- (b) For each algebraic weight $\chi \in B$, there is an isomorphism of $\overline{\mathbb{Q}_p} \otimes \mathfrak{A}^- \otimes \mathcal{H}$ -modules

$$M \otimes_{\mathcal{O}(B)} k(y) = \bigoplus_{y \in \kappa^{-1}(\chi)} F(\mathcal{C}_\chi)[\psi_y],$$

up to some normalizing twist depending on χ for the action of \mathfrak{A}^- .

- (c) $\kappa^{-1}(\kappa(x))^{\text{red}} = \{x\}$ (in particular, V is connected) and the natural surjection $\mathcal{O}(V) \rightarrow k(x)$ has a section.

The classicity criterion implies that for $\chi \in B$ algebraic, dominant and sufficiently far from the walls of its Weyl chamber, $M \otimes_{\mathcal{O}(B)} k(\chi) \subset F(W_\chi^\vee)$, so for $z \in \kappa^{-1}(\chi)$ we have

$$(4.19) \quad F(W_\chi^\vee)[\psi_z] = F(\mathcal{C}_\chi)[\psi_z].$$

Let $Z_0 \subset B$ be a Zariski-dense subset of such weights, and let $Z \subset V$ be the union of $\{x\}$ and $\kappa^{-1}(Z_0)$, this a Zariski-dense subset of classical points of V . Up to reducing Z_0 if necessary, we may assume that the refinements associated to any $z \in Z$ and any $v \in S_p$ are regular, and that κ is étale at each point in $Z \setminus \{x\}$.

The first important fact is that

$$(4.20) \quad M \otimes_{\mathcal{O}(B)} k(\chi_0) = F(W_{\chi_0}^\vee)[\psi_x] = F(\mathcal{C}_{\chi_0})[\psi_x].$$

Indeed, by (b) and (c) above it is enough to check that $F(\mathcal{C}_{\chi_0})[\psi_x] \subset F(W_{\chi_0}^\vee)$ but this is Proposition 4.2 as the \mathcal{F}_v are regular and non-critical (note that $\varphi_{i,v} \varphi_{j,v}^{-1} \neq p$ as π_v is tempered and unramified).

For each $z \in Z(\overline{\mathbb{Q}_p})$, let $\tilde{\Pi}_z$ be the (unique) global discrete A -packet of representations of $\text{U}(\mathbb{A}_F)$ containing the representations π which are unramified outside $S \setminus S_p$, with $\pi_\infty \simeq W_{\kappa(z)}$, and whose system of $\mathfrak{A}^- \otimes \mathcal{H}$ -eigenvalues on $\pi_{S \setminus S_p}^{K_{S \setminus S_p}}$ is $\iota_\infty \iota_p^{-1}(\psi_z)$. Let $\tilde{\Pi}_z \subset \Pi_z$ be the (finite) subset of representations having these last three properties. We deduce from (4.13) that for each $z \in Z(\overline{\mathbb{Q}_p})$, if we set $\delta(z) := \dim_{\overline{\mathbb{Q}_p}} F(\mathcal{C}_{\kappa(z)})[\psi_z]$, then

$$(4.21) \quad \delta(z) = \sum_{\pi \in \Pi_z} m(\pi) \dim_{\mathbb{C}}(\pi_{S \setminus S_p}^{K_{S \setminus S_p}}),$$

where $m(\pi)$ denotes the multiplicity of π in the discrete spectrum of U . To obtain the formula above, we have used that $\pi_v^{K_v}$ has dimension 1 for $v \notin S$ and that for $v \in S_p$ the generalized eigenspace of $\psi_z|_{\mathfrak{A}^-}$ on $\pi_v^{K_v}$ is also one-dimensional as the $\varphi_{i,v}$ are distinct for each $v \in S_p$ by assumption.

By (4.19) and (4.20), we have $F(W_\chi^\vee)[\psi_z] = F(\mathcal{C}_\chi)[\psi_z]$ for each $z \in Z$. Note that the action of $\mathfrak{A}^- \otimes \mathcal{H}$ on each $F(W_\chi^\vee)[\psi_z]$ for $z \in Z$ is actually scalar. As a consequence, if we can show that $\forall \chi \in Z_0$, $\kappa^{-1}(\chi)$ has a single element, and that the residue field of this element

is $k(x)$, then (a), (b) and (c) above, as well as the local freeness of M over $\mathcal{O}(B)$, ensure that $\mathcal{O}(V) = \mathcal{O}(B) \otimes_{\mathbb{Q}_p} k(x)$ and we are done. By the local freeness of M again, it is enough to show that $\delta(z)$ does not depend on $z \in Z(\overline{\mathbb{Q}_p})$, or at least that $\delta(z) \geq \delta(x)$ for any z in $Z(\overline{\mathbb{Q}_p})$. We shall use for this the natural n -dimensional pseudo-character $T_Y : \text{Gal}(\overline{E}/E) \rightarrow \mathcal{O}(Y)$ for each irreducible component Y of V ; these components all meet at x by (c).

We have not used so far that π has a cuspidal base change to $\text{GL}_n(\mathbb{A}_E)$. This implies that $\rho_\pi = \rho_x$ is absolutely irreducible (by Ribet, Blasius and Rogawski). As the absolutely irreducible locus is Zariski open in V , up to reducing Z_0 if necessary we may assume that for all $z \in Z(\overline{\mathbb{Q}_p})$ the packet $\tilde{\Pi}_z$ has a cuspidal base change as well. In this case, Π_z is tempered (hence a "tensor product" of local tempered L -packets) and each element $\pi \in \Pi_z$ is automorphic and has multiplicity $m(\pi) = 1$ by [35, Thm. 13.3.3 (c), Thm. 14.6.1] (when $n = 2$ this is due to Labesse-Langlands). By Lemmas 4.5, 4.6 and 4.7, as well as Formula (4.21), we obtain that $\delta(z)$ is independent of $z \in Z(\overline{\mathbb{Q}_p})$, which concludes the proof. These lemmas apply as for each π' with cuspidal base-change Π' , π' is tempered and $\Pi' \mapsto \rho_{\pi'}$ is compatible with the Frobenius semi-simplified local Langlands correspondence at all primes prime to p (that follows from Rogawski's strong base change).

Note that strictly, we have not considered the case of a split place $v \in S$ such that $U(F_v)$ is the group of units of a central simple algebra over F_v . But in this case the statement of Lemma 4.6 still makes sense and holds: it follows from Lemma 4.5 (ii) and the fact that if π_1 and π_2 are two representations of $U(F_v)$ whose Weil–Deligne representations are inertially equivalent, then π_1 and π_2 differ by an unramified twist, hence coincide over the maximal compact subgroups of $U(F_v)$. \square

REMARK 4.9. – A similar statement is probably true in many other cases (all?) when π is only assumed to be tempered, but this would have forced us to look at Rogawski's multiplicity formula in unpleasant detail. This is however simple enough if $S \setminus S_p$ does not contain any non split prime. Indeed, in this case the multiplicity formula [35, Thm. 14.6.5] implies that if Π is a discrete tempered (possibly endoscopic) packet of U , then all the elements $\pi \in \Pi$ which are unramified at p and such that $\pi^K \neq 0$ have multiplicity one, so the argument above still applies.

We end this paragraph with a result valid for any n . Assume that E/F is unramified everywhere and that U is quasi-split at each finite place of F (so $[F : \mathbb{Q}]$ is even if n is even, by Hasse's principle for unitary groups). Assume that S only contains primes which split in E . Let again \mathcal{E} be the p -adic eigenvariety of U of level K associated to the set S_p of all places above p and to the global Hecke-algebra \mathcal{H} unramified outside S .

THEOREM 4.10. – *Assume that U , E/F and K are as above. Let π be a tempered automorphic representation of U which is unramified above p and such that $\pi^K \neq 0$. We assume that π has multiplicity 1 in the discrete spectrum of U and that for each $v \in S_p$ the crystalline Frobenius has n distinct eigenvalues on $D_{\text{crys}}(\rho_{\pi, \tilde{v}})$.*

Let $\{\mathcal{F}_v\}$ be a collection of refinements of the $\rho_{\pi, \tilde{v}}$ for $v \in S_p$ and let $x \in \mathcal{E}$ be the point associated to $(\pi, \{\mathcal{F}_v\})$. If \mathcal{F}_v is non-critical and regular for each $v \in S_p$, then \mathcal{E} is étale at x over weight space.

By a result of Labesse [30, Thm. 5.4], such a π has multiplicity 1 if its base-change to $\mathrm{GL}_n(\mathbb{A}_E)$ is cuspidal, in which case it is necessarily tempered. It follows from Langlands' conjectures that any π as in the statement should have multiplicity one.

Proof. – We start with an observation. Let π and π' be any two automorphic representations of U . Assume that π and π' are unramified at each finite place v of F which is inert in E , and that $\pi_v \simeq \pi'_v$ if v is Archimedean and for almost all finite places v of F . We claim that $\pi \simeq \pi'$. Indeed, let Π and Π' be their respective base change to $\mathrm{GL}_n(\mathbb{A}_E)$. By Labesse [30, Cor. 5.3], Π and Π' are induced from discrete automorphic representations of a Levi subgroup of $\mathrm{GL}_n(\mathbb{A}_E)$, and $\Pi_v \simeq \Pi'_v$ for almost all the finite places v of E . By Mœglin–Waldspurger classification of the discrete spectrum of GL_m and by Jacquet–Shalika [25, Thm. 4.4], this implies that $\Pi \simeq \Pi'$ (at all places). By Labesse's theorem above again, this implies that $\pi \simeq \pi'$. Indeed, the global base change is compatible with the obvious local base change at all the finite places. At the split places it is the identity and at the inert places it is the spherical base change, so in both cases it is injective, and we are done. As a consequence, if ψ is a system of eigenvalues of $\mathcal{H} \otimes \mathfrak{A}^-$ on $F(W_\chi^\vee)$, then there is a unique automorphic representation $\pi(\psi)$ of U of weight χ such that \mathcal{H} acts on $\pi(\psi)_f^S$ as ψ .

We now go back to the proof of the theorem. The arguments of the proof of Thm. 4.8 apply verbatim until “We have not used so far that π has a cuspidal base-change” if for any $z \in Z(\overline{\mathbb{Q}}_p)$ we set $\Pi_z = \{\pi(\psi_z)\}$. If x and π are as in the statement then $\pi(\psi_x) \simeq \pi$. As $m(\pi(\psi_x)) = 1$ by assumption, we have

$$\delta(x) := \dim_{\overline{\mathbb{Q}}_p} F(\mathcal{C}_{\chi_0})[\psi_x] = \dim_{\mathbb{C}} (\pi_{S \setminus S_p}^{K_{S \setminus S_p}}).$$

By Lemma 4.7 applied to each irreducible component of V , we obtain that for each $z \in Z(\overline{\mathbb{Q}}_p)$ we have $\delta(z) = m(\pi(\psi_z))\delta(x) \geq \delta(x)$, which concludes the proof. This lemma applies as $\pi(\chi_z)$ is tempered for each $z \in Z(\overline{\mathbb{Q}}_p)$: at x it is the assumption, and at $z \neq x$ it follows from [30, Cor. 5.3] and [9, Thm. 1.2] as $\kappa(z)$ is not in a Weyl wall of $X^*(T)$. \square

5. End of the proof of Theorem A and other global applications

5.1. Proof of Theorem 2.11

Fix $x \in \mathfrak{X}(\bar{\rho})$ a modular point. For each place v above p , set $V_v := \rho_x|_{G_{\bar{\mathbb{Q}}_v}}$. Assume that for each v , $\mathrm{End}_{G_{\bar{\mathbb{Q}}_v}}(V_v) = L$, and that the eigenvalues of the crystalline Frobenius of $D_{\mathrm{cris}}(V_v)$ are distinct and in $k(x)$.

Let $y \in \mathcal{E}(\bar{\rho})$ be a refined modular point above x . Those y are in natural bijection with the set of collections of refinements \mathcal{F}_v of V_v , $v \in S_p$, in the sense of § 3.9. Indeed, they have the form $y = (x, \delta)$ and this bijection is

$$\delta \mapsto (\mathcal{F}_v)_{v \in S_p} = (\delta_{1,v}(p)p^{k_{1,v}}, \delta_{2,v}(p)p^{k_{2,v}}, \delta_{3,v}(p)p^{k_{3,v}})_{v \in S_p},$$

where $k_{1,v} < k_{2,v} < k_{3,v}$ are the Hodge-Tate weights of V_v . For this reason we shall also denote those y above x by $(x, \{\mathcal{F}_v\})$. We use the notations of § 3.9.

PROPOSITION 5.2. – *Let $(x, \{\mathcal{F}_v\}) \in \mathcal{E}(\bar{\rho})$ be a refined modular point such that \mathcal{F}_v is non-critical for each $v \in S_p$. For any Artinian thickening $\mathrm{Spec}(A) \hookrightarrow \mathcal{E}(\bar{\rho})$ at x , the associated Galois representation $\rho_A : G_E \rightarrow \mathrm{GL}_3(A)$ satisfies*

$$\rho_A|_{G_{\bar{v}}} \in \mathfrak{X}_{V_v, \mathcal{F}_v}(A).$$

Note that this statement makes sense as $\mathrm{End}_{G_{\bar{v}}}(V_v) = L$ by Remark 3.5.

Proof. – This is [1, Thm. 4.4.1]. □

Let us set $t_{v,*} := \mathfrak{X}_{V_v,*}(L[\varepsilon])$. Fix $y = (x, \{\mathcal{F}_v\})$ a refined modular point and consider the natural map on tangent spaces:

$$(5.22) \quad T_y(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho})) \xrightarrow{(\mathrm{res}_{\mathbf{v}})_{v \in S_p}} \prod_{v \in S_p} t_v.$$

By the proposition above, we know that if \mathcal{F}_v is non critical for each v then the image of this map falls inside $\prod_{v \in S_p} t_{v, \mathcal{F}_v}$. However, this information is still too weak to localize this image: the space on the left, which is of global nature, tends to have dimension $3 \cdot |S_p|$, whereas $\prod_{v \in S_p} t_{v, \mathcal{F}_v}$, which is purely locally defined, has dimension $7 \cdot |S_p|$. A key idea is to neglect the crystalline deformations in the range of (5.22), as they should be conjecturally transversal to the global ones by the Bloch-Kato conjecture (see [1, Conj. 7.6.5]). This is confirmed by the following proposition.

PROPOSITION 5.3. – *Assume again that \mathcal{F}_v is non-critical for each v , then the linear map (5.22) induces an isomorphism*

$$T_{(x, \{\mathcal{F}_v\})}(\mathcal{E}(\bar{\rho})) \xrightarrow{\sim} \prod_{v \in S_p} t_{v, \mathcal{F}_v} / t_{v, \mathrm{crys}}.$$

Proof. – By Theorem 4.8 and Prop. 5.2, the composite of the natural maps

$$T_{(x, \{\mathcal{F}_v\})}(\mathcal{E}(\bar{\rho})) \longrightarrow \prod_{v \in S_p} t_{v, \mathcal{F}_v} / t_{v, \mathrm{crys}} \xrightarrow{\mathrm{P}_{\mathrm{sen}}} \prod_{v \in S_p} L^3$$

is an isomorphism. But by Prop 3.15 the second map is an isomorphism, hence so is the first one. □

COROLLARY 5.4. – *The image of $\bigoplus_{y=(x, \{\mathcal{F}_v\})} T_y(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho}))$ has $k(x)$ -dimension at least $6[F : \mathbb{Q}]$.*

Proof. – Consider the natural map

$$\bigoplus_{y=(x, \{\mathcal{F}_v\})} T_y(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho})) \longrightarrow \prod_{v \in S_p} t_v / t_{v, \mathrm{crys}}$$

where we restrict in the sum on the left to those $\{\mathcal{F}_v\}$ such that \mathcal{F}_v is non-critical for all v . By Theorem 3.20, Example 3.25 and Prop. 5.3, the composite of the two maps above is surjective. The result follows as the space on the right has dimension $\frac{n(n+1)}{2} \cdot |S_p| = 6 \cdot |S_p|$ by Prop. 3.15. □

5.5. Generalizations and other settings

Assume now that n is any odd (say) integer, and keep the notations of §1.1. Assume again that $\bar{\rho}$ is modular. We expect that the strategy employed when $n = 3$ will lead to a proof of the natural variant of Thm. 1.14 for any n (with 3 and 6 replaced by n and $\frac{n(n+1)}{2}$). It would actually follow by the same proof if we could prove the following conjecture on the genericity of global Galois representations that we believe in:

CONJECTURE 5.6 (Genericity conjecture). – *For any modular point $x \in \mathfrak{X}(\bar{\rho})$ and any affinoid neighborhood U of x in $\mathfrak{X}(\bar{\rho})$, there exists a modular point $y \in U$ such that $\rho_{y,\tilde{v}}$ is generic for each $v \in S_p$.*

Of course, we expect to find such a point y by starting from x and moving in the piece of the infinite fern inside U . It would even be enough for our purpose to have a similar statement with “is generic” replaced by “has a nested sequence of non-critical refinements”. When $n = 3$, Lemma 2.9 and Example 3.25 give a positive answer to this weak version, but even in this case the above conjecture is still open as far as we see. Note that it would not be too difficult to show that we may find an y in U such that the Zariski-closure of the image of $\rho_{y,\tilde{v}}$ is the whole of $\mathrm{GL}_n(\overline{k(x)})$ (extending Lemma 2.9 for any n). However, this does not seem to imply that $\rho_{y,\tilde{v}}$ is generic enough. In particular, there does not seem to be any variant of Example 3.25 when $n > 3$. We hope to come back to this conjecture in the future. Here is another natural question.

QUESTION 5.7. – *How non-generic can geometric Galois representations be? Are there some (even conjectural) global conditions implying the genericity at each place above p ?*

The example we have in mind is the classical conjecture that if a p -adic Galois representation $\rho_f : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}_p})$ attached to a classical modular eigenform f is split at p (hence non-generic), then the form f is CM.

In another direction, our infinitesimal approach also shed some lights back to the standard case studied by Gouvêa and Mazur. The following result is a simple consequence of our method and of Prop. 3.23.

PROPOSITION 5.8. – *Let $\rho_f : G_{\mathbb{Q},S} \rightarrow \mathrm{GL}_2(\overline{\mathbb{Q}_p})$ be a p -adic Galois representation attached to a modular eigenform f of weight $k > 1$ and level prime to p . Assume that $\rho_f|_{G_{\mathbb{Q}_p}}$ is indecomposable and that the two eigenvalues of its crystalline Frobenius are distinct. Then the two leaves of the infinite fern of Gouvêa-Mazur cross transversally at ρ_f .*

Let us end this paper by considering the Hilbert modular analogue of Thm 1.14, to which the arguments of Gouvêa-Mazur do not extend as well when $[F : \mathbb{Q}] > 1$ (as far as we know).

Here F is a totally real field which is *totally split at the odd prime p* , S a finite set of places of F containing the places above p and ∞ , and $\bar{\rho} : G_{F,S} \rightarrow \mathrm{GL}_2(\mathbb{F}_q)$ is absolutely irreducible and totally odd. In this context, we say that ρ is modular if it is isomorphic to the p -adic Galois representation ρ_{Π} attached to a cuspidal automorphic Galois representation Π of $\mathrm{GL}_2(\mathbb{A}_F)$ which is cohomological, unramified above p and outside S (hence to a Hilbert modular eigenform), and we say that $\bar{\rho}$ is modular if it is isomorphic to the residual

representation of a modular ρ . Let $\mathfrak{X}(\bar{\rho})$ be the generic fiber of the universal $G_{F,S}$ -deformation of $\bar{\rho}$, as defined by Mazur. In the unobstructed case $H^2(G_{F,S}, \text{ad}(\bar{\rho})) = 0$, then $\mathfrak{X}(\bar{\rho})$ is the open unit ball over \mathbb{Q}_q in $1 + 2[F : \mathbb{Q}]$ variables. This unobstructedness assumption also implies that $H^2(G_{F,S}, \mathbb{F}_p) = 0$, thus Leopoldt's defect \mathfrak{d} of F at p vanishes. To avoid assuming Leopoldt's conjecture in general it is convenient to work with the notion of an *essentially modular* ρ , which we define as the twist $\rho = \rho_{\Pi} \otimes \chi$ of a modular ρ_{Π} by a continuous character $\chi : G_{F,S} \rightarrow \mathbb{Q}_p^*$. We obtain this way notions of modular and essentially modular points in $\mathfrak{X}(\bar{\rho})$. For simplicity we assume moreover that $[F : \mathbb{Q}]$ is even (see Rem. 5.10).

THEOREM 5.9. – *Assume that $\bar{\rho}$ is modular. Then the irreducible components of the Zariski-closure of the essentially modular points in $\mathfrak{X}(\bar{\rho})$ all have dimension at least $1 + \mathfrak{d} + 2[F : \mathbb{Q}]$. If $H^2(G_{F,S}, \text{ad}(\bar{\rho})) = 0$ then the modular points are Zariski-dense in $\mathfrak{X}(\bar{\rho})$.*

Of course, an important ingredient in the proof is the Hilbert modular eigenvarieties. These eigenvarieties have been studied by several authors: Hida, Kassaei, Buzzard, Kisin–Lai, Emerton, and Yamagami. We shall mostly rely on Buzzard's results.

Define $\mathfrak{X}^{\text{emod}} \subset \mathfrak{X}(\bar{\rho})$ as the subset of essentially modular points. If $x \in \mathfrak{X}^{\text{emod}}$ and $\rho_x = \rho_{\Pi} \otimes \chi$, then Π is well defined up to a twist by an algebraic Hecke character of F unramified above p and we shall denote it by Π_x . Let $\mathfrak{X}^{\text{gen}} \subset \mathfrak{X}^{\text{emod}}$ be the subset of x such that for each $v|p$, $\rho_{\Pi_x, v}$ is absolutely irreducible and with distinct crystalline Frobenius eigenvalues in $k(x)$ (in particular it is generic).

There is a finite index subgroup $\Gamma \subset \mathcal{O}_F^*$ of the totally positive units such that the universal character $G_{F,S}^{\text{ab}} \rightarrow R(\bar{\rho})^*$ is trivial over Γ via the reciprocity map. Let \mathcal{T} and \mathcal{W} be the rigid analytic spaces over \mathbb{Q}_p parameterizing respectively the continuous p -adic characters of $(F_p^*)^2$ and of $(\mathcal{O}_{F_p}^*)^2$ which are trivial on Γ diagonally embedded. The space \mathcal{W} is called the *weight space*. It has equidimension $\dim(\mathcal{W}) = 1 + \mathfrak{d} + [F : \mathbb{Q}]$. The $\bar{\rho}$ -Hilbert modular eigenvariety

$$\mathcal{E}(\bar{\rho}) \subset \mathfrak{X}(\bar{\rho}) \times \mathcal{T}$$

is the Zariski closure of the pairs (x, \mathcal{T}) of refined essentially modular points (using a translation similar as in §5.1 between refinements of an essentially modular ρ and elements of \mathcal{T}).

$\mathcal{E}(\bar{\rho})$ has been studied by Buzzard in [8, §III], by switching to a totally definite quaternion algebra D over F which is split at all the finite places of F and using the Jacquet-Langlands correspondence. Such a D exists as $[F : \mathbb{Q}]$ is even. The important properties of $\mathcal{E}(\bar{\rho})$ are the following:

(i) $\mathcal{E}(\bar{\rho})$ is equidimensional of dimension $\dim(\mathcal{W})$. The natural map to the weight space $\kappa : \mathcal{E}(\bar{\rho}) \rightarrow \mathcal{T} \rightarrow \mathcal{W}$ is locally finite. The set of (x, \mathcal{T}) with $x \in \mathfrak{X}^{\text{emod}}$ is a Zariski-dense accumulation subset of $\mathcal{E}(\bar{\rho})$. The same holds for the (x, \mathcal{T}) with x modular if (and only if) $\mathfrak{d} = 0$.

These properties are not stated this way, *loc. cit.*, but they follow simply from the construction there and an argument similar to the one of Thm. 2.3. There is a classicality criterion in this case as well. Arguing as in Thm. 4.10 (note that the strong multiplicity one property holds for D^*), we obtain that:

(ii) If $x \in \mathfrak{X}^{\text{emod}}$ is such that $\rho_{\Pi_x, v}$ has distinct crystalline Frobenius eigenvalues at each $v|p$, and if \mathcal{T}_v is a non-critical refinement of $\rho_{x, v}$ for each $v|p$, then κ is étale at $(x, \{\mathcal{T}_v\})$.

Moreover, the following (easier) variant of Lemma 2.9 holds:

(iii) *Let $x \in \mathfrak{X}^{\text{emod}}$ and assume that a collection $\mathcal{F} = (\mathcal{F}_v)_{v \in S_p}$ of refinements of the $\rho_{x,v}$ has the property that for each $v|p$ such that $\rho_{x,v}$ is ordinary, then the first Frobenius eigenvalue of \mathcal{F}_v is the one with the greatest p -adic valuation. Then the (y, \mathcal{F}') with $y \in \mathfrak{X}^{\text{gen}}$ accumulate at (x, \mathcal{F}) in $\mathcal{E}(\bar{\rho})$.*

Proof of Thm. 5.9. – The proof is only easier than in the $\text{U}(3)$ case, so we shall be rather sketchy.

Consider some $x \in \mathfrak{X}^{\text{gen}}$. Let $(x, \mathcal{F}) \in \mathcal{E}(\bar{\rho})$ be an associated refined point and consider the composite of the natural maps

$$(5.23) \quad T_{(x, \mathcal{F})}(\mathcal{E}(\bar{\rho})) \longrightarrow T_x(\mathfrak{X}(\bar{\rho})) \rightarrow \prod_{v \in S_p} \mathfrak{t}_v / \mathfrak{t}_{v, \text{crys}}$$

as in (5.22). Using the étaleness of $\mathcal{E}(\bar{\rho}) \rightarrow \mathcal{W}$ at (x, \mathcal{F}) , Prop. 3.15, and arguing as in §5.1, we see that the image of (5.23) is exactly the subspace $\mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}^0$ where $\mathfrak{t}_{\mathcal{F}} = \prod_{v \in S_p} \mathfrak{t}_{v, \mathcal{F}_v} / \mathfrak{t}_{v, \text{crys}}$ and $\mathfrak{t}^0 \subset \prod_{v \in S_p} \mathfrak{t}_v / \mathfrak{t}_{v, \text{crys}}$ is the subspace parameterizing the deformations of $(\rho_{x,v})_{v \in S_p}$ whose determinant, viewed as a morphism $F_p^* \rightarrow k(x)^*$, factors through Γ . Remark that

$$\dim_{k(x)} \mathfrak{t}^0 = 3[F : \mathbb{Q}] - ([F : \mathbb{Q}] - 1 - \mathfrak{d}) = 1 + \mathfrak{d} + 2[F : \mathbb{Q}].$$

Moreover $\sum_{\mathcal{F}} \mathfrak{t}_{\mathcal{F}} \cap \mathfrak{t}^0 = \mathfrak{t}^0$ by Thm. 3.19 (remark that $\cap_{\mathcal{F}} \mathfrak{t}_{\mathcal{F}} \rightarrow (\prod_{v \in S_p} \mathfrak{t}_v) / \mathfrak{t}^0$ is surjective as “families of twists are \mathcal{F} -trianguline for each \mathcal{F} ”). It follows that the image of the natural map

$$\bigoplus_{\mathcal{F}} T_{(x, \mathcal{F})}(\mathcal{E}(\bar{\rho})) \rightarrow T_x(\mathfrak{X}(\bar{\rho}))$$

has $k(x)$ -dimension at least $1 + \mathfrak{d} + 2[F : \mathbb{Q}]$. The first statement follows by the same argument as in §2.8. The second statement (unobstructed case) follows as well by property (i) of $\mathcal{E}(\bar{\rho})$ and by the local-global numerical coincidence

$$\dim \mathfrak{X}(\bar{\rho}) = \dim \mathfrak{t}^0. \quad \square$$

REMARK 5.10. – When $[F : \mathbb{Q}]$ is odd, the same result holds, with the same proof, if we restrict to essentially modular points $\rho_{\Pi} \otimes \chi$ such that Π_w is essentially square integrable at some finite place $w \in S \setminus S_p$, as we may switch to a suitable definite quaternion algebra in this case as well. However there may sometimes be no such modular point (e.g. when $S = S_p \cup S_{\infty}$). We could actually still conclude in most cases in general using some well chosen quadratic base-change and results of Kisin–Lai, but this would lead us too far away from our purposes here.

6. An application to adjoint Selmer groups

6.1. Bloch-Kato Selmer groups

Let p be a prime, L a finite extension of \mathbb{Q}_p . In what follows, an L -representation of a topological group G will always mean a continuous, finite-dimensional, L -linear representation. Let ℓ be a prime, M a finite extension of \mathbb{Q}_{ℓ} , and let V be an L -representation of G_M . Recall that Bloch and Kato defined in [5, §3] a subspace $H_f^1(M, V) \subset H^1(M, V)$. Set $H_s^1(M, V) = H^1(M, V) / H_f^1(M, V)$. Let F be a number field, let S be a finite set of finite

places of F containing the set S_p of places dividing p , the set S_∞ of Archimedean places, and let V be an L -representation of G_F unramified outside S . Let $H_{f'}^1(F, V)$ be the kernel of the natural map $H^1(G_{F,S}, V) \rightarrow \prod_{v \in S \setminus S_p} H_s^1(F_v, V)$, it does not depend on S as above. In what follows, we shall be mostly interested in the surjectivity of the natural map:

$$(6.24) \quad H_{f'}^1(F, V) \rightarrow \prod_{v|p} H_s^1(F_v, V).$$

The kernel of this map is usually denoted by $H_f^1(F, V)$. We may like viewing the surjectivity of (6.24) as some splitting of the Poitou-Tate exact sequence. By the global duality theorem, the image of (6.24) is Tate's orthogonal complement of the image of $H_f^1(F, V^\vee(1)) \rightarrow \prod_{v|p} H_f^1(F_v, V^\vee(1))$. In particular, (6.24) is surjective if and only if the natural map $H_f^1(F, V^\vee(1)) \rightarrow \prod_{v|p} H^1(F_v, V^\vee(1))$ is the zero map.

REMARK 6.2. – We shall be mainly concerned with cases where V is geometric (say in the strongest sense) of pure weight 0. In this case, the conjectures of Bloch-Kato ([23, §3.4], [5]) assert that $H_f^1(F, V) = 0$, so (6.24) is expected to be injective. Assume furthermore that F is totally real, let c_v denote a complex conjugation at the Archimedean place v , and denote by $h_v^{(0,0)}$ the dimension of the $(0, 0)$ -part of the F_v -Hodge structure associated to V . If

$$\text{trace}(c_v|_V) = -h_v^{(0,0)}$$

for each real place v , then those conjectures predict as well that $H_f^1(F, V^\vee(1)) = 0$ (compare them with [37, §3]). In particular, (6.24) should be an isomorphism in this case! As far as we know, very little is known about this conjecture, possibly nothing when $\dim(V) \geq 3$ before the results of this paper.

6.3. The adjoint' Selmer group of a Galois representation of type $U(n)$

Assume now that F is a totally real field. Let E be a totally imaginary quadratic extension of F and assume that p is totally split in E (hence in F). For each $v|p$ fix some place \tilde{v} of E above v , so $F_v = E_{\tilde{v}} = \mathbb{Q}_p$. Let $c \in G_F$ be a complex conjugation.

Let Π be a cuspidal automorphic representation of $\text{GL}_n(\mathbb{A}_E)$ such that $\Pi^\vee \simeq \Pi^c$ and such that Π_v is algebraic regular at each Archimedean place v of E . Let ρ_Π be the p -adic Galois representation of G_E associated to Π and ι , say with coefficients in L and normalized so that $\rho_\Pi^\vee \simeq \rho_\Pi(n-1)$. We make the following assumption.

ASSUMPTION (A). – For each $v|p$, Π_v is unramified, its Langlands class has n distinct eigenvalues, and $\text{End}_{G_v}(\rho_{\Pi|G_v}) = L$.

Note that it implies that ρ_Π is absolutely irreducible. We shall assume furthermore that L is big enough so as to contain (via ι) the eigenvalues alluded to above, which is harmless.

We are going to define below an L -representation $\text{Ad}'(\rho_\Pi)$ of G_F of dimension n^2 that extends the standard representation $\text{ad}(\rho_\Pi)$ of G_E . From the self-duality like condition, there is a $P \in \text{GL}_n(L)$, unique up to L^* , such that

$${}^t \rho_\Pi(cgc^{-1})^{-1} = P \rho_\Pi(g) P^{-1} \chi(g)^{n-1}, \forall g \in G_E$$

($\chi : G_F \rightarrow \mathbb{Q}_p^*$ is the cyclotomic character). It follows that ${}^tP = \pm P$, and the main theorem of [2] ensures that P is a *symmetric* matrix. Following [16, §2], let us consider the linear algebraic \mathbb{Q} -group

$$\mathcal{G}_n := (\mathrm{GL}_n \times \mathrm{GL}_1) \rtimes \mathrm{Gal}(E/F),$$

where c acts on $\mathrm{GL}_n \times \mathrm{GL}_1$ by $(g, \lambda) \mapsto ({}^t g^{-1} \lambda, \lambda)$. We check at once that there exists a unique morphism $\widetilde{\rho}_\Pi : G_F \rightarrow \mathcal{G}_n(L)$ such that $\widetilde{\rho}_\Pi(g) = (\rho_\Pi(g), \chi(g)^{1-n})$ for $g \in G_E$ and $\widetilde{\rho}_\Pi(c) = ({}^t P^{-1}, 1)c$. We denote by

$$\mathrm{Ad}'(\rho_\Pi)$$

the representation of G_F defined by the adjoint representation of $\widetilde{\rho}_\Pi$ on

$$M_n(L) = \mathrm{Lie}(\mathrm{GL}_n \times \{1\}) \otimes L \subset \mathrm{Lie}(\mathcal{G}_n) \otimes L.$$

The G_F -equivalence class of $\mathrm{Ad}'(\rho_\Pi)$ only depends on the G_E -equivalence class of ρ_Π . The map $(X, Y) \mapsto \mathrm{trace}(XY)$ defines a $\mathcal{G}_n(L)$ -equivariant pairing $M_n(L) \otimes M_n(L) \rightarrow L$, thus $\mathrm{Ad}'(\rho_\Pi)$ is selfdual. Note that the complex conjugation c acts on $M_n(L) = \mathrm{Ad}'(\rho_\Pi)$ as $X \mapsto -P^t X P^{-1}$. In particular, $\mathrm{Ad}'(\rho_\Pi)^{G_F} = 0$, and the homotheties induce an embedding $\varepsilon_{E/F} \rightarrow \mathrm{Ad}'(\rho_\Pi)$, where $\varepsilon_{E/F}$ is the character of order 2 of G_F associated to E/F .

In what follows we shall be mainly interested in the Selmer group

$$H_{f'}^1(F, \mathrm{Ad}'(\rho_\Pi)).$$

The purity of ρ_Π (see [9] for the most general case) ensures that $H^0(F_v, \mathrm{Ad}'(\rho_\Pi)^\vee(1))$ vanishes, hence so do $H^2(F_v, \mathrm{Ad}'(\rho_\Pi))$ and $H_s^1(F_v, \mathrm{Ad}'(\rho_\Pi))$, for each finite place v . In particular, for each finite set S containing S_p , S_∞ and the ramification of $\mathrm{Ad}'(\rho_\Pi)$, we have

$$(6.25) \quad H^1(G_{F,S}, \mathrm{Ad}'(\rho_\Pi)) = H_{f'}^1(F, \mathrm{Ad}'(\rho_\Pi)).$$

CONJECTURE 6.4. – *If $V = \mathrm{Ad}'(\rho_\Pi)$, then (6.24) is an isomorphism, and $H_f^1(F, V) = H_f^1(F, V^\vee(1)) = 0$.*

Indeed, we are in the case of Remark 6.2. Note that for each real place v , $h_v^{(0,0)} = n$ by the description of the Hodge-Tate numbers of ρ_Π , and $\mathrm{trace}(c_v|_V) = -n$ by the result of [2] recalled above.

The *adjoint'* Selmer group of ρ_Π has a natural description as the tangent space of a suitable deformation functor. Fix an S as above and let \mathcal{C} be the category defined in §3.1. For an object A of \mathcal{C} let $\mathfrak{X}_{\rho_\Pi}(A)$ be the set of A -isomorphism classes of continuous representations $\rho_A : G_{E,S} \rightarrow \mathrm{GL}_n(A)$ such that $\rho_A \otimes_A L \simeq \rho_\Pi$ and such that $\mathrm{trace}(\rho_A^\vee) = \mathrm{trace}(\rho_A^c) \chi^{n-1}$. This defines a functor $\mathfrak{X}_{\rho_\Pi} : \mathcal{C} \rightarrow \mathrm{Sets}$, equipped with a natural morphism

$$\mathfrak{X}_{\rho_\Pi} \rightarrow \prod_{v \in S_p} \mathfrak{X}_{\rho_{\Pi, \widetilde{v}}} =: \mathfrak{X}_p,$$

where we have set $\rho_{\Pi, \widetilde{v}} := \rho_\Pi|_{G_{\widetilde{v}}}$. This allows to define a collection of subfunctors of the left-hand side by pulling back the subfunctors of the right-hand side studied in §3. Let $\mathcal{F} = \{\mathcal{F}_v\}$ be a collection of refinements of the $\rho_{\Pi, \widetilde{v}}$ for $v \in S_p$. We set

$$\mathfrak{X}_{\rho_\Pi, \mathcal{F}} = \mathfrak{X}_{\rho_\Pi} \times_{\mathfrak{X}_p} \prod_{v|p} \mathfrak{X}_{\rho_{\Pi, \widetilde{v}}, \mathcal{F}_v} \quad \text{and} \quad \mathfrak{X}_{\rho_\Pi, f} = \mathfrak{X}_{\rho_\Pi} \times_{\mathfrak{X}_p} \prod_{v|p} \mathfrak{X}_{\rho_{\Pi, \widetilde{v}}, \mathrm{crys}}.$$

Note that conditions (i), (ii), (iii) and (iv) of §3.9 on $\rho_{\Pi, \sim}$ are satisfied by conditions (i) and (ii) above (and by the known description of its Hodge-Tate weights for (iv)). Similar functors were previously introduced in [1, §7.6].

PROPOSITION 6.5. – $\mathfrak{X}_{\rho_{\Pi}} : \mathcal{C} \rightarrow \text{Sets}$ is pro-representable and its tangent space is canonically isomorphic to $H_{f'}^1(F, \text{Ad}'(\rho_{\Pi}))$.

Moreover, for $* \in \{f, \mathcal{F}\}$, $\mathfrak{X}_{\rho_{\Pi}, *}$ is a pro-representable subfunctor of $\mathfrak{X}_{\rho_{\Pi}}$. The tangent space of $\mathfrak{X}_{\rho_{\Pi}, f}$ coincides with $H_f^1(F, \text{Ad}'(\rho_{\Pi}))$. When \mathcal{F}_v is non-critical for each $v \in S_p$, then $\mathfrak{X}_{\rho_{\Pi}, f} \subset \mathfrak{X}_{\rho_{\Pi}, \mathcal{F}}$.

Proof. – The functor $\mathfrak{X}_{\rho_{\Pi}}$ is pro-representable by Mazur as ρ_{Π} is absolutely irreducible (the condition $\text{trace}(\rho_*^{\vee}) = \text{trace}(\rho_*^c)\chi^{n-1}$ being obviously relatively representable). The identification of its tangent space with $H^1(G_{F, S}, \text{Ad}'(\rho_{\Pi}))$ is similar to Lemma 1.5 (using the \mathcal{G}_n here) and is checked in [16, §2]. The other assertions follow from Prop. 3.15. \square

DEFINITION 6.6. – For each collection of refinements $\mathcal{F} = \{\mathcal{F}_v\}$ for $v \in S_p$, define $H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_{\Pi}))$ (resp. $H_{\mathcal{F}_v}^1(F_v, \text{Ad}'(\Pi))$) as the tangent space of $\mathfrak{X}_{\rho_{\Pi}, \mathcal{F}}$ (resp. $\mathfrak{X}_{\rho_{\Pi, \sim}, \mathcal{F}_v}$).

6.7. The main results

Fix Π as in §6.3. The following statement (in some special cases) was conjectured in [1, §7.6].

THEOREM 6.8. – Assume that \mathcal{F}_v is non-critical and regular for each $v \in S_p$. Then the natural map

$$H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_{\Pi})) \longrightarrow \prod_{v|p} H_{\mathcal{F}_v}^1(F_v, \text{Ad}'(\rho_{\Pi})) / H_f^1(F_v, \text{Ad}'(\rho_{\Pi}))$$

is surjective. In particular, $\dim_L H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_{\Pi})) = \dim_L H_f^1(F, \text{Ad}'(\rho_{\Pi})) + n[F : \mathbb{Q}]$.

Proof. – Let us choose F'/F a totally real quadratic extension of F . Assume that F'/F is split above p , that $F'_v = E_v$ for each finite place v of F which is either ramified in E or which is inert and such that Π_v is ramified. Let Π' be Arthur-Clozel's quadratic base-change of Π to $E \cdot F'$, it is cuspidal by (i). As the f and \mathcal{F} conditions can be checked after any finite base-change which is split above p , the inflation-restriction sequence induces isomorphisms

$$H_*(F, V) = H_*(F', V|_{G_{F'}})^{\text{Gal}(F'/F)}$$

for $* \in \{f', f, \mathcal{F}\}$. As a consequence, it is enough to show the surjectivity of the map of the statement when F, E and Π are replaced by $F', E \cdot F'$ and Π' . In particular, we may assume that $[F : \mathbb{Q}]$ is even, that E/F is unramified everywhere, and that the places w of E such that Π_w is ramified are split over F . (This kind of trick has been used in another context by Blasius-Rogawski and Harris-Taylor.)

As $[F : \mathbb{Q}]$ is even, there exists a unitary group U/F as in §4.1 which is furthermore quasi-split at all finite places. By Labesse [30, Thm. 5.4], Π admits a strong descent to an automorphic representation π of U which has furthermore multiplicity 1. Let S be a finite set of places of F which split in E , containing S_p , and such that π_f is unramified outside S . Let $K = \prod_v K_v$ be a compact open subgroup of $U(\mathbb{A}_{F, f})$ as in §4.1 such that K_v is hyperspecial for each $v \notin S$ and small enough so that $\pi_f^K \neq 0$. Let X be the eigenvariety

of U of level K associated to ι and the spherical Hecke algebra outside S . Let $\{\mathcal{F}_v\}_{v \in S_p}$ be as in the statement and let $x \in X$ be the point associated to $(\pi, \{\mathcal{F}_v\})$. Up to extending the scalars in X , we may assume that X is defined over the field L of §6.3. The natural pseudo-character $G_{E,S} \rightarrow \mathcal{O}(X)$ (see e.g. [1, Prop. 7.5.4]) and the absolute irreducibility of ρ_Π define a canonical L -linear map on tangent spaces

$$(6.26) \quad T_x(X) \longrightarrow \mathfrak{X}_{\rho_\Pi}(L[\varepsilon]) = H^1(G_{F,S}, \text{Ad}'(\bar{\rho})(\rho_\Pi)) = H_{f'}^1(F, \text{Ad}'(\rho_\Pi)).$$

By [1, Thm. 4.4.1] (using that \mathcal{F}_v is non-critical and regular for each v), the image of (6.26) falls inside $H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_\Pi))$. Consider now the induced map:

$$(6.27) \quad T_x(X) \longrightarrow \prod_{v|p} H_{\mathcal{F}_v}^1(F_v, \text{Ad}'(\bar{\rho})(\rho_\Pi)) / H_f^1(F_v, \text{Ad}'(\bar{\rho})(\rho_\Pi)).$$

We claim that this map is an isomorphism, which will conclude the proof. Note that the subspace $T_x(X) \subset H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_\Pi))$ even furnishes a canonical section of the map of the statement. Consider the natural map

$$\prod_{v|p} H_{\mathcal{F}_v}^1(F_v, \text{Ad}'(\bar{\rho})(\rho_\Pi)) / H_f^1(F_v, \text{Ad}'(\bar{\rho})(\rho_\Pi)) \xrightarrow{(\text{P}_{\text{Sen}})_v} (L^n)^{S_p}.$$

By Prop. 3.15 this is an isomorphism, so it is enough to check that the composite of (6.27) with this map is an isomorphism. But this composite is the derivative of the map from X to the weight-space at x , which is an isomorphism by Thm. 4.10, and we are done. \square

THEOREM 6.9. – *Assume that $\forall v|p$, $\rho_{\Pi, \tilde{v}}$ is weakly generic and regular (§ 3.24). Then*

$$H_{f'}^1(F, \text{Ad}'(\rho_\Pi)) = \sum_{\mathcal{F}} H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_\Pi))$$

and the natural map

$$H_{f'}^1(F, \text{Ad}'(\rho_\Pi)) \longrightarrow \prod_{v|p} H_s^1(F_v, \text{Ad}'(\rho_\Pi))$$

is surjective.

For each $v|p$, let $\mathcal{F}_{i,v}$ be a weakly nested sequence of non-critical regular refinements of $\rho_{\Pi, \tilde{v}}$. The equality $H_{f'}^1(F, \text{Ad}'(\rho_\Pi)) = \sum_{\mathcal{F}} H_{\mathcal{F}}^1(F, \text{Ad}'(\rho_\Pi))$ even holds if we restrict the sum to the $n^{[F:\mathbb{Q}]}$ refinements \mathcal{F} such that \mathcal{F}_v is one of the $\mathcal{F}_{i,v}$ for each v .

Proof. – It is an immediate consequence of Thm. 3.20 and Thm. 6.8. \square

COROLLARY 6.10. – *Under the assumptions of Theorem 6.9, and for each S as in (6.25),*

$$\dim_L H_f^1(F, \text{Ad}'(\rho_\Pi)) = \dim_L H^2(G_{F,S}, \text{Ad}'(\rho_\Pi)).$$

In particular, if $H^2(G_{F,S}, \overline{\text{Ad}'(\rho_\Pi)}) = 0$ for some residual $\overline{\mathbb{F}}_p$ -representation $\overline{\text{Ad}'(\rho_\Pi)}$ associated to $\text{Ad}'(\rho_\Pi)$, then $H_f^1(F, \text{Ad}'(\rho_\Pi)) = 0$.

Proof. – By Thm. 6.9 and Prop. 3.15, we have

$$(6.28) \quad \dim_L H_{f'}^1(F, \text{Ad}'(\rho_\Pi)) - \dim_L H_f^1(F, \text{Ad}'(\rho_\Pi)) = \frac{n(n+1)}{2} [F : \mathbb{Q}].$$

As we already saw, $H^0(G_{F,S}, \text{Ad}'(\rho_\Pi)) = 0$ and $\text{Ad}'(\rho_\Pi)(c)(X) = -P^{-1}XP$ for some symmetric invertible matrix P by [2], so

$$\dim_L \text{Ad}'(\rho_\Pi)^{c=-1} = \frac{n(n+1)}{2}.$$

This numerical coincidence with the right-hand side of (6.28), as well as (6.25) and the global Euler characteristic formula conclude the proof. \square

Theorem 6.9 is very general as a lot of Selmer groups are direct summands of adjoint' Selmer groups. Here is an important special case.

Let F be a totally real field in which p is totally split and let Π be a regular algebraic cuspidal automorphic representation of $\text{GL}_n(\mathbb{A}_F)$ such that $\Pi^\vee \simeq \Pi\eta$ for some Hecke character η of F such that the sign $\eta_\infty(-1) := \eta_v(-1)$ is independent⁽¹²⁾ of the archimedean place v of F . Denote by $q \in \mathbb{Z}$ the integer such that $\eta| \cdot |^{-q}$ is an Artin character of F . Let $\rho_\Pi : G_F \rightarrow \text{GL}_n(L)$ be the p -adic Galois representation associated to Π and ι ([14]), and denote by $\eta_\iota : G_F \rightarrow L^*$ the character associated to η and ι . Recall that $\rho_\Pi^\vee \simeq \rho_\Pi \eta_\iota \chi^{n-1}$ and that ρ_Π is symplectic if n is even and $\eta_\infty(-1)(-1)^q = 1$, and orthogonal otherwise by [2, Cor. 1.3]. Assume that for each place v of F dividing p , Π_v is unramified and its Langlands class has n distinct eigenvalues. Let E be any totally imaginary quadratic extension of F in which p is totally split.

THEOREM 6.11. – *Let Π be as above and assume that for each $v|p$, $\rho_{\Pi|G_{F_v}}$ is weakly generic regular.*

(Symplectic case) If n is even and $\eta_\infty(-1)(-1)^q = 1$, then (6.24) is surjective for $V = \Lambda^2 \rho_\Pi \otimes \eta_\iota \varepsilon_{E/F} \chi^{n-1}$ and $V = \text{Sym}^2 \rho_\Pi \otimes \eta_\iota \chi^{n-1}$.

(Orthogonal case) Otherwise, (6.24) is surjective for $V = \Lambda^2 \rho_\Pi \otimes \eta_\iota \chi^{n-1}$ and $V = \text{Sym}^2 \rho_\Pi \otimes \eta_\iota \varepsilon_{E/F} \chi^{n-1}$.

Proof. – Indeed, let Π' be Arthur-Clozel's base change Π' of Π to E , it is cuspidal as $\rho_{\Pi|G_{F_v}}$ is indecomposable by assumption. By [16, Lemma 4.1.4], there is an algebraic Hecke character ν of E which is unramified at p and such that $\Pi' \otimes \nu$ is conjugate self-dual. This latter representation clearly satisfies assumption (A) of §6.3 by the corresponding assumption on Π . A simple computation left to the reader shows that

$$\text{Ad}'(\rho_{\Pi' \otimes \nu}) = \Lambda^2 \rho_\Pi \otimes \eta_\iota \varepsilon_{E/F} \chi^{n-1} \oplus \text{Sym}^2 \rho_\Pi \otimes \eta_\iota \chi^{n-1}$$

in the first case, and

$$\text{Ad}'(\rho_{\Pi' \otimes \nu}) = \Lambda^2 \rho_\Pi \otimes \eta_\iota \chi^{n-1} \oplus \text{Sym}^2 \rho_\Pi \otimes \eta_\iota \varepsilon_{E/F} \chi^{n-1}$$

in the second. \square

Let f be a classical modular eigenform of weight $k > 1$ and let $\rho_f : G_{\mathbb{Q},S} \rightarrow \text{GL}_2(\overline{\mathbb{Q}}_p)$ be the p -adic Galois representation associated to f and ι . Assume that the level of f is prime to p and let E be a quadratic imaginary field split at p .

⁽¹²⁾ Actually, Langlands' conjectures imply that this last condition should follow from the others.

THEOREM 6.12. – Assume that $\rho_f|_{G_{\mathbb{Q}_p}}$ is absolutely indecomposable and that the ratio of its two crystalline Frobenius eigenvalues is not a root of unity. Then (6.24) is an isomorphism for:

- (i) $V = \text{Sym}^n(\rho_f) \otimes \det(\rho_f)^{-n/2}$ if $n = 2, 6$.
- (ii) $V = \text{Sym}^n(\rho_f) \otimes \det(\rho_f)^{-n/2} \epsilon_{E/\mathbb{Q}}$ if $n = 0, 4$.

Moreover, for each such V we have $H_f^1(\mathbb{Q}, V) = H^2(G_{\mathbb{Q},S}, V)$.

Proof. – Let Π_0 be the cuspidal automorphic representation of $\text{GL}_2(\mathbb{A}_{\mathbb{Q}})$ associated to ρ_f and let η be its central character. We normalize Π_0 so that $\eta = |\cdot|^{2-k}$ times a Dirichlet character, so $\rho_f = \rho_{\Pi_0}$. As is well known, we have $\eta_{\infty}(-1)(-1)^{k-2} = 1$.

By Kim-Shahidi [28] the representation $\Pi := \text{Sym}^3 \Pi_0$ of $\text{GL}_4(\mathbb{A}_{\mathbb{Q}})$ defined at all places by the local Langlands correspondence is automorphic and cuspidal (note that f is not CM by the assumption at p). We have $\Pi^{\vee} \simeq \Pi \eta^{-3}$. Moreover Π is symplectic as $(\eta_{\infty}(-1)(-1)^{(k-2)})^3 = 1$. The first part of the theorem is then a consequence of Thm. 6.11 applied to Π . Its assumptions are satisfied by Examples 3.26 and 3.28. The last sentence follows from the global Euler characteristic formula as in the proof of Cor. 6.10. \square

Appendix: Some unobstructed Galois representations of type U(3)

Let A be an elliptic curve over \mathbb{Q} . Let $p \geq 5$ be a prime of good reduction of A and assume that the representation

$$G_{\mathbb{Q}} \rightarrow \text{Aut} A[p] \simeq \text{GL}_2(\mathbb{F}_p)$$

is surjective. Let E be a quadratic imaginary field and S the set of places containing ∞ , p and the primes dividing $\text{disc}(E)\text{cond}(A)$. Assume to simplify that p splits in E and that $\text{gcd}(\text{disc}(E), \text{cond}(A)) = 1$. By Example 1.9, the representation

$$\bar{\rho} = \text{Symm}^2 A[p](-1)|_{G_{E,S}}$$

is modular of type U(3). Let $\text{Ad}'(\bar{\rho})$ be the $G_{\mathbb{Q},S}$ -module associated to $\bar{\rho}$ as in §1.1.

LEMMA 6.13. – There is an isomorphism of $\mathbb{F}_p[G_{\mathbb{Q},S}]$ -modules:

$$\text{Ad}'(\bar{\rho}) \simeq \bar{\rho} \oplus \epsilon_{E/\mathbb{Q}} \oplus (\text{Symm}^4 A[p])(-2) \otimes \epsilon_{E/\mathbb{Q}},$$

where $\epsilon_{E/\mathbb{Q}}$ is the non-trivial character of $\text{Gal}(E/\mathbb{Q})$.

We are looking for a set of conditions ensuring that $H^2(G_{\mathbb{Q},S}, \text{Ad}'(\bar{\rho})) = 0$. Let K be the field of definition of some \mathbb{F}_p -line in $A[p]$ (so $[K : \mathbb{Q}] = p + 1$), $\chi : G_K \rightarrow \mathbb{F}_p^*$ the natural character on the quotient of $A[p]$ by this line, and K' the extension of K cut out by the character $\chi^4 \epsilon_{E/\mathbb{Q}}(-1)$. If ℓ is a prime of good reduction of A , we set $a_{\ell} = \ell + 1 - |A(\mathbb{F}_{\ell})|$.

PROPOSITION 6.14. – If the following conditions are satisfied, then $\bar{\rho}$ is unobstructed.

- (i) $\forall \ell \in S, H^0(\mathbb{Q}_{\ell}, \text{Ad}'(\bar{\rho})(1)) = 0$.
- (ii) p does not divide the degree of a modular parameterization of A ,
- (iii) The class numbers of E and K' are prime to p .

Moreover, $H^0(\mathbb{Q}_{\ell}, \text{Ad}'(\bar{\rho})(1)) = 0$ if:

- (a) $\ell = p$, unless A is ordinary at p , $A[p]|_{G_{\mathbb{Q}_p}}$ is split, and $a_p = \pm 1 \pmod{p}$.
- (b) $\ell \mid \text{disc}(E)$, $\ell \neq 1 \pmod{p}$ and $a_{\ell} \neq \pm(\ell + 1) \pmod{p}$.

(c) $\ell \mid \text{cond}(A)$, A is semistable at ℓ , $A[p]_{|G_{\mathbb{Q}_\ell}}$ is non-split, ℓ^2 and $\ell^3 \epsilon_{E/\mathbb{Q}}(\ell) \not\equiv 1 \pmod{p}$.

Proof. – As $\mathcal{O}_E^* \otimes \mathbb{F}_p = \text{Cl}(E) \otimes \mathbb{F}_p = 0$ we have $\text{III}_S^1(G_{\mathbb{Q},S}, \epsilon_{E/\mathbb{Q}}(1)) = 0$, thus assumption (i) ensures first that $H^2(G_{\mathbb{Q},S}, \epsilon_{E/\mathbb{Q}}) = 0$. By (i) again, $H^0(\mathbb{Q}_\ell, \text{Symm}^2 A[p]) = 0$ for each $\ell \in S$, thus assumption (ii) implies the vanishing of $H^2(G_{\mathbb{Q},S}, \bar{\rho})$ by a result of Flach [22, §3]. We now deal with the last term $U := \text{Symm}^4 A[p](-2) \otimes \epsilon_{E/\mathbb{Q}} = U^*$. Note that $H^0(G_{\mathbb{Q},S}, U^*(1)) = 0$ by assumption (i), so Tate's global duality theorem shows that $H^2(G_{\mathbb{Q},S}, U) = 0$ if and only if $\text{III}_S^1(G_{\mathbb{Q},S}, U(1)) = 0$.

Let $V = \mathbb{F}_p^2$ be the standard representation of $G = \text{GL}_2(\mathbb{F}_p)$, let $B \subset G$ denote the upper triangular subgroup and $\chi : B \rightarrow \mathbb{F}_p^*$ the unique quotient of $V|_B$. We have an $\mathbb{F}_p[G]$ -equivariant injection $\text{Symm}^4 V \rightarrow \text{Ind}_B^G(\chi^4)$ and the quotient Q is well known to be irreducible of dimension $p - 4$, isomorphic to $\det(V)^2$ when $p = 5$. In particular, in all cases $H^0(G_{\mathbb{Q},S}, Q(-1) \otimes \epsilon_{E/\mathbb{Q}}) = 0$ and we have natural injections

$$H^1(G_{\mathbb{Q},S}, U(1)) \longrightarrow H^1(G_{K,S}, \chi^4(-1)\epsilon_{E/\mathbb{Q}}) \longrightarrow H^1(G_{K',S}, \mathbb{F}_p).$$

Via this injection, $\text{III}_S^1(G_{\mathbb{Q},S}, U(1))$ is mapped into $\text{Cl}(K') \otimes \mathbb{F}_p$, hence (iii).

The second part of the statement is an immediate computation and is left to the reader. \square

We use the notations of Cremona's tables of elliptic curves [21].

PROPOSITION 6.15 (Under GRH). – *Let $p = 5$, $E = \mathbb{Q}(i)$ and assume that A belongs to one of the isogeny classes*

$$17A, 21A, 37B, 39A, 51A, 53A, 69A, 73A, 83A, \text{ and } 91B.$$

Then $H^2(G_{\mathbb{Q},S}, \text{Ad}'(\bar{\rho})) = 0$.

These curves are exactly the elliptic curves A of odd, square-free, conductor $N < 100$ such that each prime divisor of N is congruent to $\pm 2 \pmod{5}$, and such that $a_2(A) \neq \pm 2$ (this rules out the classes 37A, 43A, 67A and 91A). These last two conditions are the ones in (b) and part of (c) above, and are actually necessary local conditions for the unobstruction of the deformation functor of $A[5]$ itself.

The third row of the following table gives the degrees of the prime isogenies in a class, note that there is no isogeny of degree 5. The sixth row gives the j invariant of a certain element in the class, namely of: 17A4, 21A4, 37B3, 39A4, 51A1, 53A1, 69A1, 73A2, 83A1, and 91B1. Their valuation at any ℓ dividing $\text{cond}(A)$ is < 5 , so $A[p]_{|G_{\mathbb{Q}_\ell}}$ is not split for any such ℓ . This also shows that $G_{\mathbb{Q}} \rightarrow \text{Aut}(A[5])$ is surjective for each A . Using the table below, we see that the criteria (a), (b) and (c) of the proposition apply, hence (i) holds.

The seventh row gives the modular degree of the strong Weil curve in each class, (ii) follows (recall that there is no 5-isogeny within a class).

For each class we computed the field K and found $K = \mathbb{Q}(x)$ with:

$$\begin{aligned}
 x^6 + 4x^5 + 17x - 17 &= 0, \\
 x^6 - 2x^5 + 9x^4 - 12x^3 + 21x^2 - 15x + 3 &= 0, \\
 x^6 - 2x^5 - 5x^4 + 10x^3 - 15x^2 + 8x - 36 &= 0, \\
 x^6 - 2x^5 - 5x^4 + 20x^3 - 25x^2 - 25x + 75 &= 0, \\
 x^6 - 2x^5 - 5x^4 - 10x^3 + 5x^2 - 4x + 48 &= 0, \\
 x^6 - x^5 - 5x^4 + 20x^3 - 15x^2 - 46x + 101 &= 0, \\
 x^6 - x^5 - 5x^4 - 80x^3 + 35x^2 + 184x + 1521 &= 0, \\
 x^6 - 2x^5 - 5x^4 + 20x^3 - 25x^2 - 59x + 143 &= 0, \\
 x^6 - 3x^5 - 5x^4 - 10x^3 + 10x^2 + 75x + 245 &= 0, \\
 \text{and } x^6 - 2x^5 - 5x^4 + 30x^3 - 35x^2 + 20x + 80 &= 0.
 \end{aligned}$$

The last row below gives the computation by Pari [17] of the class number of $K' = K(\cos(\pi/10))$, which concludes the proof. This last computation (and only this one) depends on GRH; it would be interesting to make it unconditional!

class	17A	21A	37B	39A	51A	53A	69A	73A	83A	91B
cardinal	4	6	3	4	2	1	2	2	1	3
isogenies	2	2	3	2	3		2	2		3
a_2	-1	-1	0	1	0	-1	1	1	-1	0
a_5	-2	-2	0	2	3	0	0	2	-2	-3
j	$\frac{3^3 11^3}{17}$	$\frac{47^3}{3 \cdot 7}$	$\frac{2^{15} 5^3}{37}$	$\frac{23^3}{3 \cdot 13}$	$\frac{2^{15}}{3^3 17}$	$\frac{3^5 5^3}{53}$	$\frac{-5^6}{3^2 23}$	$\frac{3^3 19^3}{73}$	$\frac{47^3}{83}$	$\frac{-2^{15} 11^3}{7 \cdot 13}$
degree	1	1	2	2	2	2	2	3	2	4
$h_{K'}$	2	2	8	2	8	2	2	32	2	2

REFERENCES

- [1] J. BELLAÏCHE, G. CHENEVIER, Families of Galois representations and Selmer groups, *Astérisque* **324** (2009).
- [2] J. BELLAÏCHE, G. CHENEVIER, The sign of Galois representations attached to automorphic forms for unitary groups, *Compositio Math.* **147** (2011), 1137–1352.
- [3] L. BERGER, Représentations p -adiques et équations différentielles, *Invent. Math.* **148** (2002), 219–284.
- [4] L. BERGER, Équations différentielles p -adiques et (ϕ, N) -modules filtrés, *Astérisque* **319** (2008), 13–38.
- [5] S. BLOCH, K. KATO, L -functions and Tamagawa numbers of motives, in *The Grothendieck Festschrift, Vol. I*, Progr. Math. **86**, Birkhäuser, 1990, 333–400.
- [6] G. BÖCKLE, On the density of modular points in universal deformation spaces, *Amer. J. Math.* **123** (2001), 985–1007.

- [7] S. BOSCH, U. GÜNTZER, R. REMMERT, *Non-Archimedean analysis*, Grund. Math. Wiss. **261**, Springer, 1984.
- [8] K. BUZZARD, Eigenvarieties, in *L-functions and Galois representations*, London Math. Soc. Lecture Note Ser. **320**, Cambridge Univ. Press, 2007, 59–120.
- [9] A. CARAIANI, Local-global compatibility and the action of monodromy on nearby cycles, preprint [arXiv:1010.2188](https://arxiv.org/abs/1010.2188).
- [10] G. CHENEVIER, Familles p -adiques de formes automorphes pour GL_n , *J. reine angew. Math.* **570** (2004), 143–217.
- [11] G. CHENEVIER, Variétés de Hecke des groupes unitaires et représentations galoisiennes, cours Peccot au Collège de France, <http://www.math.polytechnique.fr/~chenevier/courspeccot.html>, 2008.
- [12] G. CHENEVIER, Une application des variétés de Hecke des groupes unitaires, in [24], 2009.
- [13] G. CHENEVIER, The p -adic analytic space of pseudo-characters of a profinite group, and pseudo-representations over arbitrary rings, preprint [arXiv:0809.0415](https://arxiv.org/abs/0809.0415).
- [14] G. CHENEVIER, M. HARRIS, Construction of automorphic Galois representations II, in [24], 2009.
- [15] L. CLOZEL et al. (eds.), *On the stabilization of the trace formula, I*, International Press, 2010.
- [16] L. CLOZEL, M. HARRIS, R. TAYLOR, Automorphy for some l -adic lifts of automorphic mod l Galois representations, *Publ. Math. I.H.É.S.* **108** (2008), 1–181.
- [17] H. COHEN, K. BELABAS, PARI/GP, <http://pari.math.u-bordeaux.fr/>.
- [18] R. COLEMAN, B. MAZUR, The eigencurve, in *Galois representations in arithmetic algebraic geometry (Durham, 1996)*, London Math. Soc. Lecture Note Ser. **254**, Cambridge Univ. Press, 1998, 1–113.
- [19] P. COLMEZ, Représentations triangulines de dimension 2, *Astérisque* **319** (2008), 213–258.
- [20] B. CONRAD, Irreducible components of rigid spaces, *Ann. Inst. Fourier (Grenoble)* **49** (1999), 473–541.
- [21] J. E. CREMONA, *Algorithms for modular elliptic curves*, second ed., Cambridge Univ. Press, 1997.
- [22] M. FLACH, A finiteness theorem for the symmetric square of an elliptic curve, *Invent. Math.* **109** (1992), 307–327.
- [23] J.-M. FONTAINE, B. PERRIN-RIOU, Autour des conjectures de Bloch et Kato: cohomologie galoisienne et valeurs de fonctions L , in *Motives (Seattle, WA, 1991)*, Proc. Sympos. Pure Math. **55**, Amer. Math. Soc., 1994, 599–706.
- [24] M. HARRIS et al., On the stabilization of the trace formula, book project of the Paris G.R.F.A. Seminar, Vol. I and II, <http://fa.institut.math.jussieu.fr/node/29>.
- [25] H. JACQUET, J. A. SHALIK, On Euler products and the classification of automorphic forms. II, *Amer. J. Math.* **103** (1981), 777–815.
- [26] O. T. R. JONES, An analogue of the BGG resolution for locally analytic principal series, *J. Number Theory* **131** (2011), 1616–1640.

- [27] A. J. DE JONG, Crystalline Dieudonné module theory via formal and rigid geometry, *Publ. Math. I.H.É.S.* **82** (1995), 5–96.
- [28] H. H. KIM, F. SHAHIDI, Functorial products for $GL_2 \times GL_3$ and the symmetric cube for GL_2 , *Ann. of Math.* **155** (2002), 837–893.
- [29] M. KISIN, Overconvergent modular forms and the Fontaine-Mazur conjecture, *Invent. Math.* **153** (2003), 373–454.
- [30] J.-P. LABESSE, Changement de base CM et séries discrètes, in [15, p. 429–470], 2009.
- [31] R. LIU, Cohomology and duality for (ϕ, Γ) -modules over the Robba ring, *Int. Math. Res. Not.* **2008** (2008), art. ID rnm 150, 32.
- [32] B. MAZUR, Deforming Galois representations, in *Galois groups over \mathbf{Q} (Berkeley, CA, 1987)*, Math. Sci. Res. Inst. Publ. **16**, Springer, 1989, 385–437.
- [33] B. MAZUR, An “infinite fern” in the universal deformation space of Galois representations, *Collect. Math.* **48** (1997), 155–193.
- [34] J. S. MILNE, *Arithmetic duality theorems*, Perspectives in Math. **1**, Academic Press Inc., 1986.
- [35] J. D. ROGAWSKI, *Automorphic representations of unitary groups in three variables*, Ann. of Math. Studies **123**, Princeton Univ. Press, 1990.
- [36] M. SCHLESSINGER, Functors of Artin rings, *Trans. Amer. Math. Soc.* **130** (1968), 208–222.
- [37] J.-P. SERRE, Facteurs locaux des fonctions zêta des variétés algébriques (définitions et conjectures), in *Séminaire de théorie des nombres Delange-Pisot-Poitou*, **11**, 1969–70, exp. n° 19, 1–15.
- [38] S. W. SHIN, Galois representations arising from some compact Shimura varieties, to appear in *Annals of math.*
- [39] T. WESTON, Unobstructed modular deformation problems, *Amer. J. Math.* **126** (2004), 1237–1252.

(Manuscrit reçu le 4 février 2010 ;
 accepté, après révision, le 18 mars 2011.)

Gaëtan CHENEVIER
 C.N.R.S et Centre de Mathématiques Laurent Schwartz
 École polytechnique
 91128 Palaiseau Cedex, France
 E-mail: chenevier@math.polytechnique.fr

