

quatrième série - tome 46 fascicule 5 septembre-octobre 2013

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Laurent CHARLES & Álvaro PELAYO & San VŨ NGỌC

Isospectrality for quantum toric integrable systems

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ISOSPECTRALITY FOR QUANTUM TORIC INTEGRABLE SYSTEMS

BY LAURENT CHARLES, ÁLVARO PELAYO AND SAN VŨ NGỌC

To Peter Sarnak on his sixtieth birthday, with admiration

ABSTRACT. – We give a full description of the semiclassical spectral theory of quantum toric integrable systems using microlocal analysis for Toeplitz operators. This allows us to settle affirmatively the isospectral problem for quantum toric integrable systems: the semiclassical joint spectrum of the system, given by a sequence of commuting Toeplitz operators on a sequence of Hilbert spaces, determines the classical integrable system given by the symplectic manifold and commuting Hamiltonians. This type of problem belongs to the realm of classical questions in spectral theory going back to pioneer works of Colin de Verdière, Guillemin, Sternberg and others in the 1970s and 1980s.

RÉSUMÉ. – Nous donnons une description complète du spectre de tout système intégrable torique quantique, au moyen de l'analyse microlocale des opérateurs de Toeplitz. Ceci résout la question de l'isospectralité pour cette classe de systèmes intégrables : le spectre semi-classique d'un système intégrable quantique torique détermine le système intégrable classique sous-jacent à symplectomorphisme près. Nous donnons aussi une description complète de la théorie spectrale semi-classique des systèmes intégrables toriques quantiques. Ces questions sont classiques en théorie spectrale et remontent aux travaux fondateurs de Colin de Verdière, Guillemin et Sternberg parmi d'autres dans les années 70 et 80.

1. Introduction

This paper gives a full description of the semiclassical spectral theory of *quantum toric integrable systems* in any finite dimension. The classical limits corresponding to quantum toric integrable systems are the so called symplectic toric manifolds or *toric systems*. Such a system consists of a compact symplectic $2n$ -manifold equipped with n commuting Hamiltonians f_1, \dots, f_n with periodic flows. The paper combines geometric techniques from the theory of toric manifolds, in the complex-algebraic and symplectic settings, with recently developed microlocal analytic methods for Toeplitz operators.

As a consequence of the spectral theory we develop, we answer the *isospectrality* question for quantum toric integrable systems, in any finite dimension: the semiclassical joint spectrum of a quantum toric integrable system, given by a sequence of commuting Toeplitz operators acting on quantum Hilbert spaces, determines the classical system given by the symplectic manifold and Poisson commuting functions, up to symplectic isomorphisms. This type of symplectic isospectral problem belongs to the realm of classical questions in inverse spectral theory and microlocal analysis, going back to pioneer works of Colin de Verdière [14, 13] and Guillemin-Sternberg [37] in the 1970s and 1980s. Colin de Verdière's works are an important inspiration for the present paper.

The question of isospectrality in Riemannian geometry may be traced back to Weyl [72, 73] and is most well known thanks to Kac's article [41], who himself attributes the question to Bochner. Kac popularized the sentence: “*can one hear the shape of a drum?*,” to refer to this type of isospectral problem. The spectral theory developed in this paper exemplifies a striking difference with Riemannian geometry, where this type of isospectrality rarely holds true, and suggests that *symplectic* invariants are much better encoded in spectral theory than *Riemannian* invariants. An approach to this problem for general integrable systems is suggested in the last two authors' article [60]. We refer to Section 8 for further remarks, and references, in these directions.

Joint spectrum

In order to state our results, let us introduce the required terminology. If (M, ω) is a symplectic manifold, a smooth map $\mu = (\mu_1, \dots, \mu_n) : M \rightarrow \mathbb{R}^n$ is called a *momentum map* for a Hamiltonian n -torus action on M if the Hamiltonian flows $t_j \mapsto \varphi_{\mu_j}^{t_j}$ are periodic of period 1, and pairwise commute :

$$\varphi_{\mu_j}^{t_j} \circ \varphi_{\mu_i}^{t_i} = \varphi_{\mu_i}^{t_i} \circ \varphi_{\mu_j}^{t_j},$$

so that they define an action of $\mathbb{R}^n / \mathbb{Z}^n$. If this action is effective and M is compact, $2n$ -dimensional and connected, we call (M, ω, μ) a *symplectic toric manifold*.

By the Atiyah and Guillemin-Sternberg theorem, for any torus Hamiltonian action on a connected compact manifold, the image of the momentum map is a rational convex polytope [1, 36]. For a symplectic toric manifold, the momentum polytope $\Delta \subset \mathbb{R}^n$ has the additional property that for each vertex v of Δ , the primitive normal vectors to the facets meeting at v form a basis of the integral lattice \mathbb{Z}^n . We call such a polytope a *Delzant polytope*.

A now standard procedure introduced by B. Kostant [43, 44, 45, 46] and J.-M. Souriau [62, 63] to quantize a symplectic compact manifold (M, ω) is to introduce a prequantum bundle $\mathcal{L} \rightarrow M$, that is a Hermitian line bundle with curvature $\frac{1}{\hbar}\omega$ and a complex structure j compatible with ω . One then defines the quantum space as the space

$$\mathcal{H}_k := H^0(M, \mathcal{L}^k)$$

of holomorphic sections of \mathcal{L}^k . The parameter k is a positive integer, the semiclassical limit corresponds to the large k limit. A description of this procedure, which is called *geometric quantization*, is given by Kostant and Pelayo in [47] from the angle of Lie theory and representation theory.

Not all symplectic manifolds have a complex structure or a prequantum bundle. However a symplectic toric manifold always admits a compatible complex structure, which is not unique. Furthermore a symplectic toric manifold M with momentum map $\mu : M \rightarrow \mathbb{R}^n$ is prequantizable if and only if there exists $c \in \mathbb{R}^n$ such that the vertices of the polytope $\mu(M) + c$ belong to $2\pi\mathbb{Z}^n$ (see Section 3). If it is the case, the prequantum bundle is unique up to isomorphisms.

In many papers, a prequantum bundle is defined as a line bundle with curvature $\frac{1}{2\pi i}\omega$. With this normalization, the cohomology class of ω is integral and the prequantization condition for toric manifolds is that, up to translation, the momentum polytope has integral vertices. This normalization may look simpler than ours, which includes a 2π -factor. Nevertheless, our choice is justified by the Weyl law. Indeed, with our normalization, the dimension of the quantum space \mathcal{H}_k is

$$\left(\frac{k}{2\pi}\right)^n \text{vol}(M, \omega) + \mathcal{O}(k^{n-1}).$$

Associated to such a quantization there is an algebra $\mathcal{T}(M, \mathcal{L}, j)$ of operators

$$T = (T_k : \mathcal{H}_k \rightarrow \mathcal{H}_k)_{k \in \mathbb{N}^*}$$

called *Toeplitz operators*. This algebra plays the same role as the algebra of semiclassical pseudodifferential operators for a cotangent phase space. Here the semiclassical parameter is $\hbar = 1/k$. A Toeplitz operator has a *principal symbol*, which is a smooth function on the phase space M . If T and S are Toeplitz operators, then $(T_k + k^{-1}S_k)_{k \in \mathbb{N}^*}$ is a Toeplitz operator with the same principal symbol as T . If T_k is Hermitian (i.e., self-adjoint) for k sufficiently large, then the principal symbol of T is real-valued. Two Toeplitz operators $(T_k)_{k \in \mathbb{N}^*}$ and $(S_k)_{k \in \mathbb{N}^*}$ commute if T_k and S_k commute for every k .




FIGURE 1. “Model image” of the spectrum of a normalized quantum toric integrable system.

We shall also need the following definitions. If P_1, \dots, P_n are mutually commuting endomorphisms of a finite dimensional vector space, then the *joint spectrum* of P_1, \dots, P_n is the set of $(\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$ such that there exists a non-zero vector v for which

$$P_j v = \lambda_j v,$$

for all $j = 1, \dots, n$. It is denoted by $\text{JointSpec}(P_1, \dots, P_n)$. The *Hausdorff distance* between two subsets A and B of \mathbb{R}^n is

$$d_H(A, B) := \inf\{\epsilon > 0 \mid A \subseteq B_\epsilon \text{ and } B \subseteq A_\epsilon\},$$

where for any subset X of \mathbb{R}^n , the set X_ϵ is $X_\epsilon := \bigcup_{x \in X} \{m \in \mathbb{R}^n \mid \|x - m\| \leq \epsilon\}$. If $(A_k)_{k \in \mathbb{N}^*}$ and $(B_k)_{k \in \mathbb{N}^*}$ are sequences of subsets of \mathbb{R}^n , we say that

$$A_k = B_k + \mathcal{O}(k^{-\infty})$$

if $d_H(A_k, B_k) = \mathcal{O}(k^{-N})$ for all $N \in \mathbb{N}^*$.

Our main result describes in full the joint spectrum of a quantum toric integrable system.

THEOREM 1.1 (Joint Spectral Theorem). – *Let $(M, \omega, \mu : M \rightarrow \mathbb{R}^n)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} and a compatible complex structure j . Let T_1, \dots, T_n be commuting Toeplitz operators of $\mathcal{T}(M, \mathcal{L}, j)$ whose principal symbols are the components of μ . Then the joint spectrum of T_1, \dots, T_n satisfies*

$$\text{JointSpec}(T_1, \dots, T_n) = g\left(\Delta \cap \left(v + \frac{2\pi}{k}\mathbb{Z}^n\right); k\right) + \mathcal{O}(k^{-\infty})$$

where $\Delta = \mu(M)$, v is any vertex of Δ and $g(\cdot; k) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ admits a C^∞ -asymptotic expansion of the form

$$g(\cdot; k) = \text{Id} + k^{-1}g_1 + k^{-2}g_2 + \dots$$

where each $g_j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is smooth. Moreover, for all sufficiently large k , the multiplicity of the eigenvalues of $\text{JointSpec}(T_1, \dots, T_n)$ is 1, and there exists a small constant $\delta > 0$ such that each ball of radius $\frac{\delta}{k}$ centered at an eigenvalue contains precisely only that eigenvalue.

Thus the joint spectrum of a quantum toric integrable system can be obtained by taking the $k^{-1}\mathbb{Z}^n$ lattice points in a polytope Δ (as in Figure 1), and applying a small smooth deformation g (as in Figure 2).

Isospectrality

We present next the isospectral theorem for toric systems. An easy consequence of the previous theorem is that the momentum polytope Δ is the Hausdorff limit of the joint spectrum of the quantum system, that is Δ , consists of the $\lambda \in \mathbb{R}^n$ such that for any neighborhood U of λ , $U \cap \text{JointSpec}(T_{1,k}, \dots, T_{n,k}) \neq \emptyset$ when k is sufficiently large.

Recall that two symplectic toric manifolds (M, ω, μ) and (M', ω', μ') are *isomorphic* if there exists a symplectomorphism $\varphi : M \rightarrow M'$ such that

$$\varphi^*\mu' = \mu.$$

By the Delzant classification theorem [21], a symplectic toric manifold is determined up to isomorphism by its momentum polytope. Furthermore, for any Delzant polytope Δ , Delzant constructed in [21] a symplectic toric manifold $(M_\Delta, \omega_\Delta, \mu_\Delta)$ with momentum polytope Δ . Now we are ready to state our isospectral theorem (see Figure 2 for an illustration of the semiclassical joint spectrum).

COROLLARY 1.2 (Isospectral Theorem). – *Let $(M, \omega, \mu : M \rightarrow \mathbb{R}^n)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} and a compatible complex structure j . Let T_1, \dots, T_n be commuting Toeplitz operators of $\mathcal{T}(M, \mathcal{L}, j)$ whose principal symbols are the components of μ . Then*

$$\Delta := \lim_{k \rightarrow \infty} \text{JointSpec}(T_1, \dots, T_n)$$




FIGURE 2. Sequence of images of the spectra of a quantum toric integrable systems as the semiclassical parameter \hbar goes to 0. The spectra lie on a plane, so they correspond to a four-dimensional integrable system with two degrees of freedom. In the Hausdorff limit corresponding to $\hbar = 0$, the spectra converge to a polytope; this is proved for a general quantum toric system in any dimension by Corollary 1.2. Therefore, one can recover the classical system from the semiclassical spectrum (i.e., the spectrum of the family of Toeplitz operators as \hbar approaches 0).

is a Delzant polytope and (M, ω, μ) is isomorphic with $(M_\Delta, \omega_\Delta, \mu_\Delta)$. In other words, one can recover the classical system from the limit of the joint spectrum.

This type of inverse result is classical and belongs to the realm of questions in inverse spectral theory, going back to similar questions raised (and in many cases answered) by pioneer works of Colin Verdière and Guillemin-Sternberg in the 1970s and 1980s. Many other contributions followed their works, for instance Datchev-Hezari-Ventura [20] and Iantchenko-Sjöstrand-Zworski [40]. A few global spectral results have also been obtained recently, for instance by Vũ Ngọc [71] for one degree of freedom pseudodifferential operators, or in the article by Dryden, Guillemin, and Sena-Dias [22] in which an equivariant spectrum of the Laplace operator is considered, and the references therein. See Section 8 for further references.

Metaplectic correction

Introducing a metaplectic correction refers to twisting the prequantum bundle (or its powers) by a half-form bundle. The metaplectic correction allows to obtain an easier control of the subprincipal terms in the semiclassical limit. In the following theorem we improve the previous Joint spectral Theorem by giving the explicit description of the spectrum up to $\mathcal{O}(k^{-2})$.

Recall that a half-form bundle of a complex manifold is a square root of its canonical bundle. Given a symplectic manifold (M, ω) with a compatible complex structure, a prequantum bundle \mathcal{L} and a half-form bundle δ , the associated quantum space is $\mathcal{H}_{m,k} = H^0(M, \mathcal{L}^k \otimes \delta)$.

We can define Toeplitz operators in this setting together with their principal symbols. To state our result, we also need the notion of subprincipal symbol of a Toeplitz operator whose definition is recalled in Section 5. Two Toeplitz operators with the same principal symbol are equal up to $\mathcal{O}(k^{-2})$ if and only if they have the same subprincipal symbols.

As we will see in Section 7, a symplectic toric manifold with moment polytope $\Delta \subset \mathbb{R}^n$ has a half-form bundle if and only if there exists a vector $u \in \mathbb{Z}^n$ such that for any one-codimensional face f of Δ , the scalar product of a primitive normal of f with u is odd. Such a vector, if it exists, is uniquely determined modulo $(2\mathbb{Z})^n$. We denote it by u_Δ .

THEOREM 1.3 (Joint Spectral Theorem with metaplectic correction)

Let $(M, \omega, \mu : M \rightarrow \mathbb{R}^n)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} , a compatible complex structure j and a half-form bundle δ . Let T_1, \dots, T_n be commuting Toeplitz operators of $\mathcal{H}_{m,k}$ whose principal symbols are the components of μ . Then the joint spectrum of T_1, \dots, T_n satisfies

$$\text{JointSpec}(T_1, \dots, T_n) = g\left(\Delta \cap \left(v + \frac{2\pi}{k}(\mathbb{Z}^n + u_\Delta/2)\right); k\right) + \mathcal{O}(k^{-\infty})$$

where $\Delta = \mu(M)$, v is any vertex of Δ and $g(\cdot; k) : \mathbb{R}^n \rightarrow \mathbb{R}^n$ admits a C^∞ -asymptotic expansion of the form

$$g(\cdot; k) = \text{Id} + k^{-1}g_1 + k^{-2}g_2 + \dots$$

where each $g_j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is smooth. Furthermore g_1 is determined by

$$g_1^i(E) = \int_0^1 f_1^i(\varphi_{\mu^i}^t(x)) dt, \quad \text{for all } E \in \Delta, x \in \mu^{-1}(E)$$

where $i = 1, \dots, n$, f_1^i is the subprincipal symbol of T^i and $\varphi_{\mu^i}^t$ is the Hamiltonian flow of μ^i . Moreover, for all sufficiently large k , the multiplicity of the eigenvalues of $\text{JointSpec}(T_1, \dots, T_n)$ is 1, and there exists a small constant $\delta > 0$ such that each ball of radius $\frac{\delta}{k}$ centered at an eigenvalue contains precisely only that eigenvalue.

Besides the average of the subprincipal symbols, it is interesting to note the shift by $u_\Delta/2$ so that no eigenvalue lies on the boundary of $g(\Delta)$ when k is sufficiently large, cf. Figure 5 for the spectrum of a model toric system with metaplectic correction.

Toeplitz quantization

A natural question is how to decide whether a given integrable system can be quantized. A discussion of this problem may be found in Garay-Van Straten [31] and the references therein (they work with pseudodifferential operators, instead of Toeplitz operators). Concretely, given a prequantizable symplectic manifold endowed with an integrable system (f_1, \dots, f_n) , it may not be possible to find a set of commuting Toeplitz operators whose principal symbols are f_1, \dots, f_n , respectively. However, in the case of toric integrable systems, we will obtain, as a byproduct of the proof of Theorem 1.1, the following existence result.

THEOREM 1.4 (Existence of Toeplitz quantization). – *Let $(M, \omega, \mu : M \rightarrow \mathbb{R}^n)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} and a compatible complex*

structure j . Then there exist mutually commuting Toeplitz operators T_1, \dots, T_n in $\mathcal{T}(M, \mathcal{L}, j)$ whose principal symbols are the components of μ .

The proofs in the paper combine geometric ideas from the theory of toric manifolds in the complex and symplectic settings with microlocal analytic methods dealing with semi-classical Toeplitz operators that were developed by the first author [8, 7, 9, 11].

2. Model for a symplectic toric manifold

We review the ingredients from the theory of symplectic toric manifolds which we need for this paper, namely the Delzant construction.

This section gives a fast review of the necessary background to read this paper.

Terminology from group actions

Let (M, ω) be a symplectic manifold, i.e., the pair consisting of a smooth manifold M and a symplectic form ω on M , that is, a non-degenerate differential closed 2-form on M .

Suppose that a Lie group G acts on a symplectic manifold (M, ω) *symplectically*, i.e., by diffeomorphisms which preserve the symplectic form. We denote by $(t, p) \mapsto t \cdot p$ the action $G \times M \rightarrow M$ of G on M . Let \mathfrak{g} be the Lie algebra of G . Any element $X \in \mathfrak{g}$ generates a vector field X_M on M , called the *infinitesimal generator*, given by

$$X_M(p) := \left. \frac{d}{dt} \right|_{t=0} \exp(tX) \cdot p,$$

where $\exp: \mathfrak{g} \rightarrow G$ is the exponential map of Lie theory. As usual, we write $\iota_{X_M} \omega := \omega(X_M, \cdot) \in \Omega^1(M)$ for the contraction 1-form. The G -action on (M, ω) is said to be *Hamiltonian* if there exists a smooth invariant map $\mu: M \rightarrow \mathfrak{g}^*$, called the *momentum map*, such that for all $X \in \mathfrak{g}$ we have that

$$\iota_{X_M} \omega = -d\langle \mu, X \rangle,$$

where $\langle \cdot, \cdot \rangle: \mathfrak{g}^* \times \mathfrak{g} \rightarrow \mathbb{R}$ is the duality pairing. If the first de Rham cohomology group $H_{\text{dR}}^1(M)$ is trivial, then any symplectic T -action on M is Hamiltonian.

The G -action is *effective* if the intersection of all stabilizer subgroups

$$G_p := \{g \in G \mid g \cdot p = p\}, \quad p \in M,$$

is the trivial group. The G -action is *free* if G_p is the trivial group for all points $p \in M$.

EXAMPLE 2.1. – The simplest example of a Hamiltonian group action is given by the standard symplectic sphere S^2 with the rotational S^1 -action. It is easy to check that the momentum map for this action is the height function $\mu(\theta, h) = h$. See Figure 3.

We will need the following classical result. Recall that a torus T is a compact, connected, commutative Lie group. As a Lie group, T is isomorphic to a finite product of circles S^1 .

THEOREM 2.2 (Atiyah [1], Guillemin and Sternberg [36]). – *If a torus T with Lie algebra \mathfrak{t} acts on a compact, connected $2n$ -dimensional symplectic manifold (M, ω) in a Hamiltonian fashion, then the image $\mu(M)$ under the momentum map $\mu: M \rightarrow \mathfrak{t}^*$ of the action is a convex polytope $\Delta \subset \mathfrak{t}^*$.*




FIGURE 3. The momentum map for the 2-sphere S^2 is the height function $\mu(\theta, h) = h$. The image of S^2 under the momentum map μ is the closed interval $\Delta := [-1, 1]$. Note that as predicted by the Atiyah-Guillemin-Sternberg Theorem, the interval $[-1, 1]$ is equal to the image under μ of the set $\{(0, 0, -1), (0, 0, 1)\}$ of fixed points of the Hamiltonian S^1 -action on S^2 by rotations about the vertical axis.

Delzant construction of toric systems

Let T be an n -dimensional torus. Denote by \mathfrak{t} the Lie algebra of T and by $\mathfrak{t}_{\mathbb{Z}}$ the kernel of the exponential map $\exp: \mathfrak{t} \rightarrow T$. We denote the isomorphism $\mathfrak{t}/\mathfrak{t}_{\mathbb{Z}} \rightarrow T$ also by \exp . A *symplectic toric manifold* $(M, \omega, T, \mu: M \rightarrow \mathfrak{t}^*)$ is a symplectic compact connected manifold (M, ω) of dimension $2n$ with an effective Hamiltonian action of T with momentum map μ .

When $T = \mathbb{R}^n/\mathbb{Z}^n$ so that $\mathfrak{t} \simeq \mathbb{R}^n$, we recover the definition given in the introduction. Two symplectic toric manifolds (M, ω, T, μ) and (M', ω', T, μ') are *isomorphic* if there exists a symplectomorphism $\varphi: M \rightarrow M'$ such that $\mu' \circ \varphi = \mu$. If it is the case, φ intertwines the torus actions.

We present the construction of Delzant [21] of symplectic toric manifolds as reduced phase spaces.

Step 1 (Starting from a Delzant polytope $\Delta \subset \mathfrak{t}^$).* – Let Δ be an n -dimensional convex polytope in the dual Lie algebra \mathfrak{t}^* . We denote by F and V the set of all codimension one faces and vertices of Δ , respectively. Every face of Δ is compact. For every $v \in V$, we write $F_v = \{f \in F \mid v \in f\}$. The polytope Δ is called a *Delzant polytope* if it has the following properties, see Guillemin [34, p. 8].

- i) For each $f \in F$ there exist $X_f \in \mathfrak{t}_{\mathbb{Z}}$ and $\lambda_f \in \mathbb{R}$ such that the hyperplane which contains f is equal to the set of all $\xi \in \mathfrak{t}^*$ such that $\langle X_f, \xi \rangle + \lambda_f = 0$, and Δ is contained in the set of all $\xi \in \mathfrak{t}^*$ such that $\langle X_f, \xi \rangle + \lambda_f \geq 0$.

Note: The vector X_f and constant λ_f are made unique by requiring that they are not an integral multiple of another such vector and constant, respectively.

- ii) For every vertex $v \in V$, the vectors X_f with $f \in F_v$ form a \mathbb{Z} -basis of the integral lattice $\mathfrak{t}_{\mathbb{Z}}$ in \mathfrak{t} .

It follows that

$$\Delta = \{\xi \in \mathfrak{t}^* \mid \langle X_f, \xi \rangle + \lambda_f \geq 0 \text{ for every } f \in F\}.$$

Also, $\#(F_v) = n$ for every $v \in V$.

Step 2 (The epimorphism $\mathbb{R}^F/\mathbb{Z}^F \rightarrow T$ and the subtorus N). – Let $\pi: \mathbb{R}^F \rightarrow \mathfrak{t}$ be defined by

$$\pi(t) := \sum_{f \in F} t_f X_f, \quad t \in \mathbb{R}^F.$$

Because, for any vertex v , the X_f with $f \in F_v$ form a \mathbb{Z} -basis of $\mathfrak{t}_{\mathbb{Z}}$, we have $\pi(\mathbb{Z}^F) = \mathfrak{t}_{\mathbb{Z}}$ and $\pi(\mathbb{R}^F) = \mathfrak{t}$. It follows that π induces an epimorphism

$$\pi': \mathbb{R}^F/\mathbb{Z}^F = (\mathbb{R}/\mathbb{Z})^F \rightarrow \mathfrak{t}/\mathfrak{t}_{\mathbb{Z}},$$

and we have the corresponding epimorphism $\exp \circ \pi': \mathbb{R}^F/\mathbb{Z}^F \rightarrow T$.

Write $\mathfrak{n} := \ker \pi$, a linear subspace of \mathbb{R}^F , and

$$N := \ker(\exp \circ \pi') \subseteq \mathbb{R}^F/\mathbb{Z}^F,$$

a compact commutative subgroup of the torus $\mathbb{R}^F/\mathbb{Z}^F$. One can check that N is connected, and therefore isomorphic to $\mathfrak{n}/\mathfrak{n}_{\mathbb{Z}}$, where $\mathfrak{n}_{\mathbb{Z}} := \mathfrak{n} \cap \mathbb{Z}^F$ is the integral lattice in \mathfrak{n} of the torus N .

Step 3 (Action of N on \mathbb{C}^F). – On the complex vector space \mathbb{C}^F , we have the action of the torus $\mathbb{R}^F/\mathbb{Z}^F$, where $t \in \mathbb{R}^F/\mathbb{Z}^F$ maps $z \in \mathbb{C}^F$ to the element $t \cdot z \in \mathbb{C}^F$ defined by

$$(t \cdot z)_f = e^{2\pi i t_f} z_f, \quad f \in F.$$

This action is Hamiltonian with momentum $\mu: \mathbb{C}^F \rightarrow (\mathbb{R}^F)^* \simeq \mathbb{R}^F$ given by

$$(1) \quad \mu(z)_f = |z_f|^2/2 - \lambda_f = (x_f^2 + y_f^2)/2 - \lambda_f, \quad f \in F.$$

Here $z_f = x_f + i y_f$, with $x_f, y_f \in \mathbb{R}$. Furthermore we work with the symplectic form

$$(2) \quad \omega := (i/4\pi) \sum_{f \in F} dz_f \wedge d\bar{z}_f = (1/2\pi) \sum_{f \in F} dx_f \wedge dy_f.$$

The factor $1/2\pi$ is introduced in order to avoid an integral lattice $(2\pi \mathbb{Z})^F$ instead of our \mathbb{Z}^F .

Hence N acts on \mathbb{C}^F Hamiltonianly and the corresponding momentum mapping is $\mu_N := \iota_{\mathfrak{n}}^* \circ \mu: \mathbb{C}^F \rightarrow \mathfrak{n}^*$, where $\iota_{\mathfrak{n}}: \mathfrak{n} \rightarrow \mathbb{R}^F$ denotes the identity viewed as a linear mapping from $\mathfrak{n} \subset \mathbb{R}^F$ to \mathbb{R}^F , and its transposed $\iota_{\mathfrak{n}}^*: (\mathbb{R}^F)^* \rightarrow \mathfrak{n}^*$ is the map which assigns to each linear form on \mathbb{R}^F its restriction to \mathfrak{n} .

Step 4 (The symplectic toric manifold $(M_\Delta, \omega_\Delta, T, \mu_\Delta)$). – It follows from Guillemin [34, Theorems 1.6 and 1.4] that 0 is a regular value of μ_N , hence the zero level set Z of μ_N is a smooth submanifold of \mathbb{C}^F , and that the action of N on Z is proper and free. The N -orbit space $M_\Delta := Z/N$ is a smooth $2n$ -dimensional manifold such that the projection $p : Z \rightarrow M_\Delta$ exhibits Z as a principal N -bundle over M_Δ . Moreover, there is a unique symplectic form ω_Δ on M_Δ such that $p^*\omega_\Delta = \iota_Z^*\omega$, where ι_Z is the identity viewed as a smooth mapping from Z to \mathbb{C}^F .

On $(M_\Delta, \omega_\Delta)$, the torus $(\mathbb{R}^F/\mathbb{Z}^F)/N \simeq T$ acts effectively and Hamiltonianly, with momentum mapping $\mu_\Delta : M_\Delta \rightarrow \mathfrak{t}^*$ determined by $\pi^* \circ \mu_\Delta \circ p = \mu|_Z$, and⁽¹⁾ $\mu_\Delta(M_\Delta) = \Delta$. In other words, $(M_\Delta, \omega_\Delta, T, \mu_\Delta)$ is a symplectic toric manifold with momentum map image equal to Δ .

THEOREM 2.3 (Delzant's Theorem). – *Any abstract symplectic toric manifold (M, ω, T, μ) with momentum polytope $\Delta \subset \mathfrak{t}^*$ is isomorphic to $(M_\Delta, \omega_\Delta, T, \mu_\Delta)$. Moreover, two symplectic toric manifolds $(M_\Delta, \omega_\Delta, T, \mu_\Delta)$ and $(M_{\Delta'}, \omega_{\Delta'}, T, \mu_{\Delta'})$ are isomorphic if and only if $\Delta = \Delta'$.*

Since the action of $\mathbb{R}^F/\mathbb{Z}^F$ preserves the complex structure of \mathbb{C}^F , M_Δ inherits by reduction a complex structure compatible with ω_Δ and invariant by the action of T (cf. [37, Theorem 3.5]). So M_Δ is a Kähler manifold.

3. Prequantization

Let us recall the basic facts we need on connections of Hermitian line bundles. A good reference for this material are Duistermaat's notes [25]. See also [47, Sections 8-14] and [49].

Let M be a manifold. Consider a Hermitian line bundle $\mathcal{L} \rightarrow M$, that is a complex line bundle endowed with a Hermitian metric. Let $C^\infty(M, \mathcal{L})$ be the space of smooth sections of \mathcal{L} and $\Omega^1(M, \mathcal{L})$ be the space of \mathcal{L} -valued 1-forms. A *connection* of \mathcal{L} is a linear operator

$$\nabla : C^\infty(M, \mathcal{L}) \rightarrow \Omega^1(M, \mathcal{L})$$

satisfying the Leibniz rule

$$\nabla(fs) = df \otimes s + f\nabla s, \quad \forall f \in C^\infty(M), \quad s \in C^\infty(M, \mathcal{L}).$$

For any vector field X of M , the *covariant derivative* of a section s of \mathcal{L} with respect to X is $\nabla_X s = \nabla s(X)$. The *curvature* of the connection is the unique 2-form R on M satisfying for any vector fields X, Y of M

$$[\nabla_X, \nabla_Y] - \nabla_{[X, Y]} = R(X, Y).$$

The connection is *compatible* with the Hermitian structure if for any sections s, t of \mathcal{L} ,

$$d(s, t) = (\nabla s, t) + (s, \nabla t),$$

where (\cdot, \cdot) denotes the Hermitian scalar product. In that case, $R = \frac{1}{i}\omega$ where ω is real valued. In this paper, we always assume the connections are compatible with the metric.

⁽¹⁾ See Guillemin [34, Theorem 1.7].

Assume that $\frac{1}{i}\omega$ is the curvature of a Hermitian line bundle connection. Then the cohomology class of $\omega/2\pi$ is integral, in the sense that it belongs to the image of the natural homomorphism $H^2(M, \mathbb{Z}) \rightarrow H^2(M, \mathbb{R})$. Conversely, given any $\omega \in \Omega^2(M, \mathbb{R})$ such that $[\omega]/2\pi$ is integral, there exists a Hermitian line bundle \mathcal{L} over M with a connection ∇ of curvature $\frac{1}{i}\omega$. Furthermore, \mathcal{L} and ∇ are unique up to isomorphism. For a proof of these results, we refer the reader to [25, Theorem 10.1] or [26, Section 15.3].

Let (M, ω) be a symplectic manifold endowed with a *prequantum bundle*, that is a Hermitian line bundle $\mathcal{L} \rightarrow M$ with a connection of curvature $\frac{1}{i}\omega$; such a symplectic manifold is called *prequantizable*. A *prequantum bundle automorphism* is a vector bundle automorphism of \mathcal{L} preserving the metric and the connection. Let G be a Lie group acting on \mathcal{L} by prequantum bundle automorphisms. This action lifts an action of G on M . One proves that the latter action is Hamiltonian and has a natural momentum map μ determined by the following condition: the induced action of the Lie algebra \mathfrak{g} on $C^\infty(M, \mathcal{L})$ is given by the Kostant-Souriau operators

$$(3) \quad f \rightarrow \nabla_{X^\#} f + i\langle \mu, X \rangle f, \quad X \in \mathfrak{g},$$

where we denote by $X^\#$ the infinitesimal action of X on M and by ∇ the covariant derivative of the prequantum bundle, cf. [26, Proposition 15.2]. If G and M are connected, the action on \mathcal{L} is conversely determined by the action on M and the momentum map μ . However, not all momentum maps generating a given action can be obtained in this way. Actually these momentum maps correspond to the Lie algebra representations on the prequantum bundle, through Equation (3).

Assume now that M is connected and T is a torus acting on M in a Hamiltonian way. The momentum map of this action is unique up to a translation by a vector of \mathfrak{t}^* . By [26, Proposition 15.4], this action can be lifted to \mathcal{L} in such a way that each element acts as a prequantum bundle automorphism. This lift is not unique, actually the momentum map corresponding to a lift is unique up to a translation by a vector of $2\pi\mathfrak{t}_{\mathbb{Z}}^*$. For a symplectic toric manifold, we can explicit everything in terms of the momentum polytope as follows.




FIGURE 4. Delzant polytopes corresponding to the complex projective spaces \mathbb{CP}^2 and \mathbb{CP}^3 equipped with scalar multiples of the Fubini-Study symplectic form.

PROPOSITION 3.1. – (a) *A symplectic toric manifold (M, ω, T, μ) with momentum polytope $\Delta \subset \mathfrak{t}^*$ admits a prequantum bundle \mathcal{L} if and only if the edges of Δ belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$.*
 (b) *When it exists, the prequantum bundle is unique up to isomorphism.*

- (c) *The momentum maps corresponding to the possible lifts (via (3)) are the ones such that the vertices of the associated polytope belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$.*

Proof. – Let (M, ω) be a symplectic toric manifold with momentum map μ . Let e be a closed edge of $\Delta = \mu(M)$. Then $\mu^{-1}(e)$ is a symplectic 2-sphere embedded in M with volume the length of e . Here the length of e is the largest positive real ℓ such that $e/\ell \in \mathfrak{t}_{\mathbb{Z}}^*$. A proof of this fact may be found in [42, Lemma 2.10].

Now if (M, ω) is prequantizable, then $[\omega]/2\pi$ is integral. So necessarily the length of any edge is an integral multiple of 2π . Since $H_2(M)$ is generated by the homology classes of the spheres corresponding to the various edges ([19, Proposition 10.6]), the condition is also sufficient. The uniqueness of the prequantum bundle follows from the fact that any symplectic toric manifold is simply connected (see [19, Th. 9.1]).

Assume now that (M, ω) has a prequantum bundle \mathcal{L} . We know that we can lift the action to \mathcal{L} . Then for any fixed point $p \in M$, the action preserves the fiber \mathcal{L}_p . Because of Formula (3), for any $\xi \in \mathfrak{t}$, $\exp(\xi)$ acts in \mathcal{L}_p by multiplication by $\exp(i\langle \mu(p), \xi \rangle)$. This implies that $\mu(p) \in 2\pi\mathfrak{t}_{\mathbb{Z}}^*$. Since the fixed points of a toric manifold are the preimages of the vertices of the polytope, this shows that the vertices belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$. \square

In the sequel, we will use the following explicit construction of the prequantum bundle for the Delzant model. We use the same notations as in Section 2. Assume that the vertices of Δ belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$. Consider the prequantum bundle of \mathbb{C}^F given by $\mathcal{L}_F := \mathbb{C}^F \times \mathbb{C}$ with connection

$$\nabla = d + \frac{1}{4\pi i} \sum_{f \in F} (x_f dy_f - y_f dx_f) = d + \frac{1}{8\pi} \sum_{f \in F} (z_f d\bar{z}_f - \bar{z}_f dz_f).$$

Since the vertices of Δ belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$, the λ_f 's defining the faces of Δ are integral multiple of 2π . So we can lift the action of $\mathbb{R}^F/\mathbb{Z}^F$ on \mathbb{C}^F to \mathcal{L}_F by

$$t \cdot (z, u) = (t \cdot z, u e^{i \sum_{f \in F} t_f \lambda_f}).$$

With a straightforward computation, one checks that this action preserves the prequantum bundle structure and that its associated momentum map is μ defined in (1).

Recall that the Delzant manifold $(M_{\Delta}, \omega_{\Delta})$ is the quotient of $Z = \mu_N^{-1}(0)$ by the action of subtorus N of $\mathbb{R}^F/\mathbb{Z}^F$. Then quotienting by N the restriction of \mathcal{L}_F to Z , we obtain a prequantum bundle \mathcal{L}_{Δ} over M_{Δ} (cf. [37, Theorem 3.2]). Furthermore the group $T = (\mathbb{R}^F/\mathbb{Z}^F)/N$ acts on \mathcal{L}_{Δ} . This action preserves the prequantum bundle structure, its associated momentum is the application μ_{Δ} defined in Theorem 2.3.

REMARK 3.2. – The construction of the prequantum bundle in the Delzant model gives an alternative proof of the fact that if the vertices of its momentum polytope Δ belong to $2\pi\mathfrak{t}_{\mathbb{Z}}^*$, a symplectic toric manifold is prequantizable. Moreover, the converse statement can be deduced directly from the last paragraph of the proof of Proposition 3.1: indeed, the polytope $(2\pi)^{-1}\Delta$ can be translated to a polytope with integral vertices if and only if its edges have integral lengths.

4. Quantum model

In this section we introduce a quantum model for quantum toric system and compute its spectrum.

Consider a Delzant polytope $\Delta \subset \mathfrak{t}^*$ with vertices in $2\pi\mathfrak{t}_{\mathbb{Z}}^*$. Then as explained in Section 3, the Delzant manifold $(M_{\Delta}, \omega_{\Delta}, T, \mu_{\Delta})$ admits a prequantum bundle \mathcal{L}_{Δ} unique up to isomorphisms. This line bundle has a unique holomorphic structure compatible with the complex structure of M_{Δ} provided by reduction and with the connection. So for any positive integer k , we can define the quantum space

$$\mathcal{H}_k^{\Delta} := H^0(M_{\Delta}, \mathcal{L}_{\Delta}^k)$$

which consists of the holomorphic sections of \mathcal{L}_{Δ}^k .

For any $X \in \mathfrak{t}$, consider the *rescaled Kostant-Souriau operator*

$$(4) \quad T_{X,k} := \langle \mu_{\Delta}, X \rangle + \frac{1}{ik} \nabla_{X^{\sharp}} : \mathcal{H}_k^{\Delta} \rightarrow \mathcal{H}_k^{\Delta}.$$

This operator is well-defined because the complex structure is invariant by the action of T on M_{Δ} . The rescaling has the effect that the $T_{X,k}$'s are self-adjoint and that their joint spectrum is the intersection of the polytope Δ and a rescaled lattice. The precise result is the following.

THEOREM 4.1. – *There is an orthogonal decomposition of the quantum space \mathcal{H}_k^{Δ} into a direct sum of lines:*

$$\mathcal{H}_k^{\Delta} = \bigoplus_{\ell \in (\frac{2\pi}{k}\mathfrak{t}_{\mathbb{Z}}^*) \cap \Delta} D_{\ell}^k$$

such that, for any $X \in \mathfrak{t}$,

$$T_{X,k}\Psi = \ell(X)\Psi, \quad \text{for all } \Psi \in D_{\ell}^k.$$

Proof. – The proof uses the Delzant construction recalled in Section 2 at the quantum level. Consider the same prequantum bundle $\mathcal{L}_F \rightarrow \mathbb{C}^F$ as in the proof of Proposition 3.1. The associated quantum space \mathcal{B}_k is the space of holomorphic sections ψ of \mathcal{L}_F^k such that

$$\int_{\mathbb{C}^F} |\psi|^2(z) \nu(z) < \infty,$$

where ν is the Liouville measure. Here holomorphic means that the covariant derivative with respect to antiholomorphic vectors vanishes. Such a holomorphic section can be written

$$\psi = e^{-k|z|^2/8\pi} f,$$

where f is a plain holomorphic function on \mathbb{C}^F . Here $|z|^2 = \sum_f |z_f|^2$. So \mathcal{B}_k can be identified with the usual Bargmann space, that is the space of holomorphic functions on a \mathbb{C}^p whose square is integrable with respect to a given gaussian weight. An orthogonal basis of \mathcal{B}_k is given by the family

$$\psi_{\alpha} = e^{-\frac{k}{8\pi}|z|^2} z^{\alpha}, \quad \alpha \in \mathbb{N}^F.$$

Consider the Kostant-Souriau operator associated to momentum μ given in (1)

$$S_{X,k} = \langle \mu, X \rangle + \frac{1}{ik} \nabla_{X^{\sharp}}.$$

If $X = e_f$ a straightforward computation shows that

$$S_{X,k} \left(e^{-\frac{k}{8\pi}|z|^2} g(z) \right) = e^{-\frac{k}{8\pi}|z|^2} \left(\frac{2\pi}{k} z_f \partial_{z_f} g - \lambda_f g \right).$$

We deduce that for any $X \in \mathbb{R}^F$ and $\alpha \in \mathbb{N}^F$,

$$(5) \quad S_{X,k}(\psi_\alpha) = \left\langle X, \frac{2\pi}{k} \alpha - \lambda \right\rangle \psi_\alpha.$$

Recall that the Delzant space was defined as the symplectic quotient of \mathbb{C}^F by the subtorus N of $\mathbb{R}^F / \mathbb{Z}^F$. The corresponding space at the quantum level is

$$\mathcal{B}_k^n := \{ \psi \in \mathcal{B}_k \mid S_{X,k} \psi = 0 \text{ for all } X \in \mathfrak{n} \}.$$

We call it the *reduced quantum space*. We deduce from Equation (5) that a basis of \mathcal{B}_k^n consists of the Ψ_α 's such that $\alpha \in \mathbb{N}^F$ satisfies

$$\left\langle X, \frac{2\pi}{k} \alpha - \lambda \right\rangle = 0$$

for all $X \in \mathfrak{n}$. Equivalently, $\frac{2\pi}{k} \alpha$ runs over $(\frac{2\pi}{k} \mathbb{N}^F) \cap (\lambda + \ker(\iota_n^*))$. This set is in bijection with $(\frac{2\pi}{k} \mathfrak{t}_{\mathbb{Z}}^*) \cap \Delta$.

LEMMA 4.2. – *The map $\pi^* + \lambda$ from \mathfrak{t}^* to $(\mathbb{R}^F)^*$ restricts to a bijection*

$$(\frac{2\pi}{k} \mathfrak{t}_{\mathbb{Z}}^*) \cap \Delta \longrightarrow (\frac{2\pi}{k} \mathbb{N}^F) \cap (\lambda + \ker(\iota_n^*)).$$

Furthermore, for any $X \in \mathbb{R}^F$ and $\ell \in \frac{2\pi}{k} \mathfrak{t}_{\mathbb{Z}}^* \cap \Delta$ we have that

$$S_{X,k}(\psi_\alpha) = \langle \ell, \pi(X) \rangle \psi_\alpha, \quad \text{if } \frac{2\pi}{k} \alpha = \pi^*(\ell) + \lambda.$$

Proof of Lemma 4.2. – We know that π^* is injective with image $\ker(\iota_n^*)$. Using the Delzant condition on a vertex (Step 1 in Section 2), one sees that π^* restricts to a bijection from $\mathfrak{t}_{\mathbb{Z}}^*$ to $(\mathbb{Z}^F)^* \cap \ker(\iota_n^*)$. By the prequantization condition, $\lambda \in 2\pi(\mathbb{Z}^F)^* \subset \frac{2\pi}{k}(\mathbb{Z}^F)^*$. So $\pi^* + \lambda$ restricts to a bijection from $\frac{2\pi}{k} \mathfrak{t}_{\mathbb{Z}}^*$ to $(\frac{2\pi}{k} \mathbb{Z}^F) \cap (\lambda + \ker(\iota_n^*))$. Furthermore, the proof to show that $\mu_\Delta(M_\Delta) = \Delta$ implies that

$$(6) \quad \pi^*(\Delta) + \lambda = \mathbb{R}_{\geq 0}^F \cap (\lambda + \ker(\iota_n^*)).$$

This implies the first part of the lemma. The second assertion follows from (5). \square

The end of the proof of Theorem 4.1 is an application of the “quantization commutes with reduction” theorem of Guillemin-Sternberg. Lifting the action of $\mathbb{T}^F = \mathbb{R}^F / \mathbb{Z}^F$ to \mathcal{L}_F as in the proof of Proposition 3.1, we get an action of \mathbb{T}^F on \mathcal{B}_k . The reduced quantum space is the subspace of \mathcal{B}_k of N -invariant vectors, in other words $\mathcal{B}_k^n = (\mathcal{B}_k)^N$. Now by Guillemin-Sternberg theorem ([37]), we have an isomorphism

$$\Phi_k : \mathcal{B}_k^n \rightarrow \mathcal{H}_k^\Delta.$$

The proof in [37] given in the compact case extends to our setting by [10] and [34]. Furthermore, under the isomorphism Φ_k , the action of the torus T on \mathcal{H}_k^Δ corresponds to the action

of \mathbb{T}^F/N on \mathcal{B}_k^n . Then, passing to the level of Lie algebras, we get the following relation between the Kostant-Souriau operators:

$$\Phi_k(S_{X,k}\Psi) = T_{\pi(X),k}\Phi_k(\Psi), \quad \text{for all } \Psi \in \mathcal{B}_k^n.$$

This concludes the proof of Theorem 4.1. \square

From the previous theorem, we deduce the following quantum normal form. Consider a Delzant polytope Δ in the Lie algebra \mathbb{R}^n of $\mathbb{R}^n/\mathbb{Z}^n$. Assume $(M_\Delta, \omega_\Delta)$ has a prequantum bundle \mathcal{L}_Δ and define the associated quantum spaces

$$\mathcal{H}_k^\Delta = H^0(M_\Delta, \mathcal{L}_\Delta^k).$$

Starting from the canonical basis (e_i) of \mathbb{R}^n , we get n operators

$$T_{i,k}^\Delta := T_{e_i,k} : \mathcal{H}_k^\Delta \rightarrow \mathcal{H}_k^\Delta, \quad k \in \mathbb{Z}_{>0}, \quad 1 \leq i \leq n,$$

defined by Kostant-Souriau Formula (4).

COROLLARY 4.3. – *For any k , $T_{1,k}^\Delta, \dots, T_{n,k}^\Delta$ are mutually commuting operators with simple joint eigenspaces. Their joint spectrum is $(v + \frac{2\pi}{k}\mathbb{Z}^n) \cap \Delta$ where v is any vertex of Δ .*

Proof. – Apply Theorem 4.1 to the polytope $\Delta - v$, whose vertices are integral. \square

5. Global quantum normal form

Toeplitz operators

We briefly review Toeplitz operators. Let (M, ω) be a compact connected symplectic manifold with a prequantum line bundle \mathcal{L} . Assume that M is endowed with a complex structure j compatible with ω , so that M is Kähler and \mathcal{L} is holomorphic. Here the holomorphic structure of the prequantum bundle is the unique one compatible with the connection. Recall that for a positive integer k , $\mathcal{H}_k := H^0(M, \mathcal{L}^k)$ is the space of holomorphic sections of \mathcal{L}^k .

Since M is compact, \mathcal{H}_k is a closed finite dimensional subspace of the Hilbert space $L^2(M, \mathcal{L}^k)$. Here the scalar product is defined by integrating the Hermitian pointwise scalar product of sections against the Liouville measure of M . Denote by Π_k the orthogonal projector of $L^2(M, \mathcal{L}^k)$ onto \mathcal{H}_k .

A *Toeplitz operator* is any sequence $(T_k : \mathcal{H}_k \rightarrow \mathcal{H}_k)_{k \in \mathbb{N}^*}$ of operators of the form

$$(7) \quad \left(T_k = \Pi_k f(\cdot, k) + R_k \right)_{k \in \mathbb{N}^*}$$

where $f(\cdot, k)$, viewed as a multiplication operator, is a sequence in $C^\infty(M)$ with an asymptotic expansion $f_0 + k^{-1}f_1 + \dots$ for the C^∞ topology, and the norm of R_k is $\mathcal{O}(k^{-\infty})$.

We denote by $\mathcal{T}(M, \mathcal{L}, j)$ the set of Toeplitz operators. The following result corresponds to [10, Theorem 1.2].

THEOREM 5.1. – *The set $\mathcal{T} = \mathcal{T}(M, \mathcal{L}, j)$ is a semiclassical algebra associated to (M, ω) in the following sense. The set \mathcal{T} is closed under the formation of product. So it is a star algebra, the identity is $(\Pi_k)_{k \in \mathbb{N}^*}$. The symbol map $\sigma_{\text{cont}} : \mathcal{T} \rightarrow C^\infty(M)[[\hbar]]$, sending T_k into the formal series $f_0 + \hbar f_1 + \dots$ where the functions f_i are the coefficients of the asymptotic expansion of the multiplier $f(\cdot, k)$, is well defined. It is onto and its kernel is the ideal consisting of $\mathcal{O}(k^{-\infty})$ Toeplitz operators. More precisely for any integer ℓ ,*

$$\|T_k\| = \mathcal{O}(k^{-\ell}) \text{ if and only if } \sigma_{\text{cont}}(T_k) = \mathcal{O}(\hbar^\ell).$$

*Furthermore, the induced product $*_{\text{cont}}$ on $C^\infty(M)[[\hbar]]$ is a star-product.*

We call the formal series

$$\sigma_{\text{cont}}(T_k) = f_0 + \hbar f_1 + \dots$$

the *contravariant symbol* of $(T_k)_{k \in \mathbb{N}^*}$. The first coefficient f_0 is the *principal symbol* of $(T_k)_{k \in \mathbb{N}^*}$. The *subprincipal symbol* of $(T_k)_{k \in \mathbb{N}^*}$ is the function

$$g_1 = f_1 + \frac{1}{2} \Delta f_0,$$

where Δ is the holomorphic Laplacian of M .

Consider two Toeplitz operators with principal and subprincipal symbols g_0, g_1 and g'_0, g'_1 respectively. Then the principal and subprincipal symbol of their composition is

$$g''_0 + \hbar g''_1 = (g_0 + \hbar g_1)(g'_0 + \hbar g'_1) + \frac{\hbar}{2i} \{g_0, g'_0\} + \mathcal{O}(\hbar^2),$$

where $\{\cdot, \cdot\}$ is the Poisson bracket of (M, ω) (cf. Theorem 1.4 of [9]).

The Kostant-Souriau operators considered in Section 4 are Toeplitz operators. More generally, let f be a function of M with Hamiltonian vector field X . Applying the Tuynman's trick ([68]), one proves that the sequence

$$T_k := \Pi_k \left(f + \frac{1}{ik} \nabla_X \right), \quad k \in \mathbb{N}^*,$$

is a Toeplitz operator with principal symbol f and subprincipal symbol $-\frac{1}{2} \Delta f$.

Normal Form

Recall that for each Delzant polytope $\Delta \subset \mathbb{R}^n$, we introduced in Section 2 a symplectic toric manifold $(M_\Delta, \omega_\Delta, \mathbb{R}^n/\mathbb{Z}^n, \mu_\Delta)$, a complex structure j_Δ on M_Δ compatible with ω_Δ . Assume that $\Delta + c$ has integral vertices for some $c \in \mathbb{R}^n$, so that $(M_\Delta, \omega_\Delta)$ has a prequantum bundle \mathcal{L}_Δ (unique up to isomorphism). We defined in Section 4 for any positive k , commuting operators $T_{1,k}^\Delta, \dots, T_{n,k}^\Delta$ acting on the Hilbert spaces $\mathcal{H}_k^\Delta = H^0(M_\Delta, \mathcal{L}_\Delta^k)$, $k \in \mathbb{Z}_{>0}$, and described explicitly their spectrum in Corollary 4.3.

THEOREM 5.2 (Global normal form for a quantum toric system)

Let $(M, \omega, \mathbb{R}^n/\mathbb{Z}^n, \mu)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} and a compatible complex structure j . Denote by Δ the momentum polytope $\mu(M) \subset \mathbb{R}^n$. Let T_1, \dots, T_n be commuting Toeplitz operators of $\mathcal{T}(M, \mathcal{L}, j)$ whose principal symbols are the components of μ .

Then there exists $k_0 > 0$, there exists a sequence $(g(\cdot; k))_{k \geq k_0}$ of smooth maps $\mathbb{R}^n \rightarrow \mathbb{R}^n$, and there exists an operator $U = (U_k : \mathcal{H}_k \rightarrow \mathcal{H}_k^\Delta)_{k \geq k_0}$ with U_k invertible for any k , such that

$$U_k(T_{1,k}, \dots, T_{n,k})U_k^{-1} = g(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k) + \mathcal{O}(k^{-\infty}).$$

Moreover, g admits an asymptotic expansion in the C^∞ topology of the form

$$g(\cdot; k) = \text{Id} + k^{-1}g_1 + k^{-2}g_2 + \dots.$$

If the operators T_j are self-adjoint (i.e., for any k , $T_{j,k}$ is a self-adjoint operator), then U_k may be chosen such that $U_k U_k^* = \text{Id}_{\mathcal{H}_k^\Delta}$.

REMARK 5.3. – For small k , the dimensions of \mathcal{H}_k and \mathcal{H}_k^Δ might be different. Theorem 5.2 does not give information about small values of k .

The proof of Theorem 5.2 will require the following technical lemma, which is a global version of a result of Eliasson [30, Corollary page 14]. Recall that in the case where E is a closed half-space, $h \in C^\infty(\mathbb{R}_x^n \times E)$ if and only if all the partial derivatives $\partial_x^k \partial_e^\ell h(x, e)$ for $(x, e) \in \mathbb{R}^n \times \overset{\circ}{E}$ have a limit at every point in $\mathbb{R}^n \times E$. (This is equivalent to saying that g has a smooth extension in a neighborhood of any point.)

LEMMA 5.4. – Let E be a vector space or a closed half-space. Let $f \in C^\infty(\mathbb{R}_{(x,\xi)}^2 \times E)$, and let

$$q(x, \xi, e) = x^2 + \xi^2.$$

Assume that $\{q, f\} = 0$ (here the Poisson bracket refers to the symplectic variables (x, ξ)). Then there exists $g \in C^\infty(\mathbb{R}_{\geq 0} \times E)$ such that

$$f(x, \xi, e) = g(q(x, \xi, e), e).$$

Proof. – Set-theoretically, there is a unique such function g . We need to prove that g is smooth. The Taylor expansion of f in the (x, ξ) variables has to commute with q . This implies that it has the form

$$\sum_{k \geq 0} q^k a_k(e),$$

where $a_k \in C^\infty(E)$. Hence by the Taylor formula, for any integer $r \geq 0$, there is a polynomial P_r in q with coefficients in $C^\infty(E)$, and a smooth function $\varphi \in C^\infty(\mathbb{R}^2 \times E)$ such that

$$f = P_r(q, e) + q^r \varphi(x, \xi, e).$$

Thus we get

$$g(t, e) = P_r(t, e) + t^r \varphi(\sqrt{t}, 0, e).$$

When $t > 0$, we simply compute the partial derivatives $\partial_t^k \partial_e^\ell g(t, e)$. They have a limit as $(t, e) \rightarrow (0, e)$ as long as $k \leq r$. Thus $g \in C^r(\mathbb{R}_{\geq 0} \times E)$, which proves the lemma. \square

Proof of Theorem 5.2. – We divide the proof into several steps.

Step 1. – By Theorem 2.3, there exists a symplectomorphism $\varphi : M \rightarrow M_\Delta$ such that $\mu = \varphi^* \mu_\Delta$. Since M_Δ is simply connected, the prequantum bundle \mathcal{L}_Δ is unique up to isomorphism. Hence φ can be lifted to a prequantum bundle isomorphism $\mathcal{L} \rightarrow \mathcal{L}_\Delta$. So φ can be quantized as an operator $U_k : \mathcal{H}_k \rightarrow \mathcal{H}_k^\Delta$ such that $U_k U_k^* = I_k$ for large k and such that for any Toeplitz operator $S = S_k$ with principal symbol s , $U_k S U_k^*$ is a Toeplitz operator whose principal symbol is $s \circ \varphi^{-1}$.

The operators U_k that we use here have been introduced in [8, Chapter 4] and similar ones have been considered by [74]. They are analogues of Fourier integral operators [39, 27].

Replacing T_j by $U T_j U^*$ we see that the problem is reduced to the case where T_1, \dots, T_n are commuting Toeplitz operators on $(M_\Delta, \mathcal{L}_\Delta)$, with joint principal symbol equal to μ_Δ .

Step 2. – We now prove the theorem by induction. Assume that, for some $N \in \mathbb{N}$, we have

$$(8) \quad (T_1, \dots, T_n) = g^{(N)}(T_1^\Delta, \dots, T_n^\Delta; k) + k^{-(N+1)} R_{N+1},$$

where R_{N+1} is a vector of n Toeplitz operators and $g^{(N)}$ is polynomial in k :

$$g^{(N)} = \text{Id} + k^{-1} g_1 + \dots + k^{-N} g_N,$$

and each $g_j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a smooth map. For simplicity we write

$$G_N := g^{(N)}(T_1^\Delta, \dots, T_n^\Delta; k).$$

Notice that, for $N = 0$, this is precisely the result of Step 1. We wish to prove that there exists an invertible Toeplitz operator $U = (U_k)_{k \in \mathbb{N}}$ such that

$$(9) \quad U(T_1, \dots, T_n)U^{-1} = g^{(N)}(T^\Delta; k) + k^{-(N+1)} h_{N+1}(T^\Delta) + k^{-(N+2)} R_{N+2}.$$

The procedure is standard and we only indicate the key points. It turns out that the case $N = 0$ is slightly different from the other cases $N > 0$. When $N = 0$, we plug (8) in the left-hand side of (9) and multiply on the right by U , and obtain

$$[U, T^\Delta] + k^{-1} U R_1 = k^{-1} G_1 U \mod k^{-2} \mathcal{T}.$$

Since both sides of the equation are Toeplitz operators of order 1, the equation is equivalent to the equality of the principal symbols :

$$\frac{1}{i} \{u, \mu_\Delta\} + u r_1 = g_1(\mu_\Delta) u.$$

Writing u of the form $u = e^{ia}$ we get the equation

$$\{\mu_\Delta, a\} = r_1 - g_1(\mu_\Delta).$$

For $N \geq 1$ we look for U in the form $U_k = \text{Id} + i k^{-N} A_N \mod k^{-(N+1)} \mathcal{T}$, where A_N is a Toeplitz operator. The same calculation as before gives the equation

$$i k^{-N} [A_N, G_N] + k^{-(N+1)} (R_{N+1} - K_{N+1}) = 0 \mod k^{-(N+2)} \mathcal{T}.$$

(We use here $2N + 1 \geq N + 2$ in order to eliminate higher order terms.) Since

$$G_N = (T_1^\Delta, \dots, T_n^\Delta) + \mathcal{O}(1),$$

the equation is equivalent to the following equation on the principal symbols :

$$\{\mu_\Delta, a_N\} = r_{N+1} - h_{N+1}(\mu_\Delta).$$

Step 3. – In order to complete the induction, we need to solve the following cohomological equation, where the unknown functions are a and g_j , $1 \leq j \leq n$:

$$(10) \quad \{\mu_j^\Delta, a\} = r_j - g_j \circ \mu_\Delta, \quad 1 \leq j \leq n.$$

The proof follows Eliasson's local argument in [30, Lemma 8], where he uses a formula due to Moser. Here we show that this local argument also works globally.

For any smooth function r on M_Δ , we define

$$M_j r = \int_0^1 r \circ \varphi_j^t dt, \quad P_j r := \int_0^1 t r \circ \varphi_j^t dt.$$

M_j and P_j are clearly linear operators sending $C^\infty(M_\Delta)$ into itself. Notice that, since the flows φ_j pairwise commute, the Fubini formula ensures that M_j and P_k commute for any j, k . The following Poisson bracket is easy to compute :

$$\begin{aligned} \{\mu_j^\Delta, P_j r\} &= \int_0^1 t \{\mu_j^\Delta, r \circ \varphi_j^t\} dt = \int_0^1 t \{\mu_j^\Delta \circ \varphi_j^t, r \circ \varphi_j^t\} dt = \int_0^1 t \{\mu_j^\Delta, r\} \circ \varphi_j^t dt \\ &= \int_0^1 t \frac{d}{dt} (r \circ \varphi_j^t) dt = r - \int_0^1 r \circ \varphi_j^t dt = r - M_j r. \end{aligned}$$

We shall need the following lemmas .

LEMMA 5.5. – Let r_1, \dots, r_n be smooth functions on M_Δ such that for all i, j ,

$$\{\mu_i^\Delta, r_j\} = \{\mu_j^\Delta, r_i\},$$

then for all $1 \leq i, j \leq n$, we have that $\{M_j r_j, \mu_i^\Delta\} = 0$.

LEMMA 5.6. – Let $f \in C^\infty(M)$ such that for all $1 \leq i \leq n$ we have that $\{\mu_i^\Delta, f\} = 0$. Then there exists $g \in C^\infty(\mathbb{R}^n)$ such that $f = g \circ \mu_\Delta$.

Proof of Lemma 5.5. – We have that

$$\begin{aligned} \{\mu_i^\Delta, M_j r_j\} &= \int_0^1 \{\mu_i^\Delta, r_j \circ \varphi_j^t\} dt = \int_0^1 \{\mu_i^\Delta, r_j\} \circ \varphi_j^t dt = \int_0^1 \{\mu_j^\Delta, r_i\} \circ \varphi_j^t dt \\ &= \int_0^1 \frac{d}{dt} (r_i \circ \varphi_j^t) dt = 0, \end{aligned}$$

as desired. \square

Proof of Lemma 5.6. – Since f is invariant by the action, set-theoretically, there exists a unique function g such that $f = g \circ \mu_\Delta$. We want to prove that g is smooth.

At a regular value of μ_Δ , this follows directly from the action-angle theorem. Let c be a critical value of μ_Δ , and let C be a small ball around c . Let $(z_1, \dots, z_k, I_1, \theta_1, \dots, I_\ell, \theta_\ell) \in \mathbb{C}^k \times (T^*S^1)^\ell$, with $k + \ell = n$ be Delzant coordinates on $(\mu_\Delta)^{-1}(C)$. Up to an affine transformation, we can assume

$$\mu_\Delta = (|z_1|^2/2, \dots, |z_k|^2/2, I_1, \dots, I_\ell).$$

By assumption, f does not depend on the θ_j coordinates, so there is a smooth function g_0 such that $f = g_0(z_1, \dots, z_k, I_1, \dots, I_\ell)$.

We apply Lemma 5.4 to the function g_0 with $(x, \xi) = z_1$ and $E = \mathbb{C}^{k-1} \times \mathbb{R}^\ell$. Thus there is a smooth function g_1 such that

$$f = g_1(|z_1|^2, z_2, \dots, z_k, I_1, \dots, I_\ell).$$

We may now apply the same lemma to

$$f_1(z, e) = g(e_1, z, e_2, \dots, e_k, e_{k+1}, \dots, e_n)$$

with $E = \mathbb{R}_{\geq 0} \times \mathbb{C}^{k-1} \times \mathbb{R}^\ell$ and get a smooth function g_2 such that

$$f = g_2(|z_1|^2, |z_2|^2, z_3, \dots, z_k, I_1, \dots, I_\ell).$$

We may repeat the argument and finally obtain a smooth function g_k such that

$$f = g_k(|z_1|^2, \dots, |z_k|^2, I_1, \dots, I_\ell).$$

This proves that g is smooth in C . Thus g is smooth on $\mu_\Delta(M_\Delta)$ (which means that there is a smooth extension of g in \mathbb{R}^n). \square

We return now to the cohomological Equation (10). By Lemmas 5.5 and 5.6, there exist smooth functions g_j on \mathbb{R}^n such that $M_j r_j = g_j \circ \mu_\Delta$. Let

$$a = P_1 r_1 + P_2 M_1 r_2 + P_3 M_2 M_1 r_3 + \dots + P_n M_{n-1} \dots M_1 r_n.$$

Notice that for any function h , $\{\mu_j^\Delta, M_j h\} = 0$. Hence, since the operators M_j and P_k commute, we get

$$\{\mu_1^\Delta, a\} = \{\mu_1^\Delta, P_1 r_1\} = r_1 - M_1 r_1 = r_1 - g_1 \circ \mu_\Delta,$$

so a solves the first equation of (10). Let

$$\tilde{a} = a - P_1 r_1.$$

We have $\{\mu_1^\Delta, \tilde{a}\} = 0$, and our system becomes

$$(11) \quad \{\mu_j^\Delta, \tilde{a}\} = \tilde{r}_j - g_j \circ \mu_\Delta, \quad j = 1, \dots, n.$$

with

$$\begin{aligned} \tilde{r}_j &:= r_j - \{\mu_j^\Delta, P_1 r_1\} = r_j - \int_0^1 t \{\mu_j^\Delta, r_1 \circ \varphi_1^t\} dt = r_j - \int_0^1 t \{\mu_1^\Delta, r_j\} \circ \varphi_1^t dt \\ &= r_j - \int_0^1 t \frac{d}{dt} (r_j \circ \varphi_1^t) dt = M_1 r_j. \end{aligned}$$

We notice that

$$\tilde{a} = P_2 \tilde{r}_2 + P_3 M_2 \tilde{r}_3 + \dots + P_n M_{n-1} \dots M_2 \tilde{r}_n,$$

so by induction \tilde{a} solves the complete system (11). Thus a solves (10).

The construction we have used to solve (10) does not require the functions r_j to be real-valued. In case they are real-valued, then a and g_j will be real-valued as well, and in Step 2 we may choose $U_k = \exp(ik^{-N} A_N)$, which is unitary.

Step 4. – From steps 2 and 3 we obtain, for any positive integer N , an invertible operator $U_N = (U_{N,k})_{\geq 0}$ (which is unitary in the case of self-adjoint operators T_j) and a smooth map

$$g^{(N)} = \text{Id} + k^{-1}g_1 + \cdots + k^{-N}g_N$$

such that

$$(12) \quad U_N(T_1, \dots, T_n)U_N^{-1} = g^{(N)}(T_1^\Delta, \dots, T_n^\Delta; k) + k^{-(N+1)}R_{N+1},$$

where R_{N+1} is a Toeplitz operator, and U_N is of the form

$$U_N = U^{(N)}U^{(N-1)} \dots U^{(0)}.$$

From step 3, we have

$$U^{(j)} = \text{Id} + ik^{-j}A_j \pmod{k^{-j}\mathcal{S}},$$

for $j \geq 1$. Therefore, one can construct by induction a sequence of symbols \tilde{a}_N such that for all $N \geq 1$, the operator $U^{(N)}U^{(N-1)} \dots U^{(1)}$ is, modulo $k^{-(N+1)}\mathcal{S}$, the Toeplitz quantization of the symbol

$$1 + ik^{-1}\tilde{a}_1 + \cdots + ik^{-N}\tilde{a}_N.$$

By the Borel summation procedure, one can find a Toeplitz operator \tilde{A} whose total symbol has the asymptotic expansion

$$\tilde{a}_1 + k^{-1}\tilde{a}_2 + \cdots + k^{-N+1}\tilde{a}_N + \cdots.$$

Moreover, one can find a smooth map \tilde{g} that admits the asymptotic expansion

$$g^{(N)} = \text{Id} + k^{-1}g_1 + \cdots + k^{-N}g_N + \cdots.$$

Now we let $\tilde{U} = (I + ik^{-1}\tilde{A})U^{(0)}$, so that for any N ,

$$\tilde{U} = U_N \pmod{k^{-(N+1)}\mathcal{S}}.$$

Thus from (14) we get, as required :

$$\tilde{U}(T_1, \dots, T_n)\tilde{U}^{-1} = \tilde{g}(T_1^\Delta, \dots, T_n^\Delta) + \mathcal{O}(k^{-\infty}).$$

In the case where the operators T_j are self-adjoint, one can change the construction of the sequence \tilde{a}_j in such a way that $U^{(N)}U^{(N-1)} \dots U^{(1)}$ is, modulo $k^{-(N+1)}\mathcal{S}$, the *exponential* of the Toeplitz quantization of the symbol $ik^{-1}\tilde{a}_1 + \cdots + ik^{-N}\tilde{a}_N$. Then we define

$$\tilde{U} := \exp(ik^{-1}\tilde{A})U^{(0)},$$

which is unitary. □

6. Isospectrality

In this section we prove Theorem 1.1 and Corollary 1.2.

Recall that the *joint spectrum* of n commuting matrices A_1, \dots, A_n is the set of

$$(\lambda_1, \dots, \lambda_n) \in \mathbb{C}^n$$

such that there exists a non-zero vector v for which

$$A_j v = \lambda_j v$$

for all $j = 1, \dots, n$. Such an n -uple $(\lambda_1, \dots, \lambda_n)$ will be called a *joint eigenvalue*. We begin with the following elementary observations.

LEMMA 6.1 ([14], Lemme 5.3). – *Let B_1, \dots, B_n be commuting self-adjoint $k \times k$ matrices. Let $\epsilon > 0$, and let $u \in \mathbb{C}^k \setminus \{0\}$ be such that $\|B_i u\| \leq \epsilon \|u\|$ for all $1 \leq i \leq n$. Then there exists $\lambda \in \mathbb{R}^n$ such that*

$$\lambda \in \text{JointSpec}(B_1, \dots, B_n) \cap [-\sqrt{n}\epsilon, \sqrt{n}\epsilon]^n.$$

Proof. – Consider the matrix

$$C := \sqrt{B_1^2 + \dots + B_n^2}.$$

Since $C \geq 0$, the minimum of the spectrum of C is equal to $\inf_{\|v\|=1} \|Cv\|$. Since

$$(13) \quad \forall v \in \mathbb{C}^k, \quad \|Cv\|^2 = \langle C^2 v, v \rangle = \|B_1 v\|^2 + \dots + \|B_n v\|^2,$$

we have $\|Cu\| \leq \sqrt{n}\epsilon \|u\|$, thus C admits an eigenvalue in $[0, \sqrt{n}\epsilon]$. The corresponding eigenspace being stable under the action of all B_j 's, it contains a common eigenvector, and by (13), the corresponding joint eigenvalue $(\lambda_1, \dots, \lambda_n)$ satisfies $|\lambda_j| \leq \sqrt{n}\epsilon$. \square

REMARK 6.2. – The constant factor \sqrt{n} in Lemma 6.1 is optimal. Indeed, let n and k be integers with $1 \geq n \geq k$. For each i with $1 \leq i \leq n$ consider the $k \times k$ diagonal matrix B_i whose j th column is equal to the zero vector for all $j \in \{1, \dots, n\} \setminus \{i\}$, and equal to the canonical basis vector e_j otherwise. Let $u = \frac{\sum_{i=1}^n e_i}{\sqrt{n}}$. Then $\|u\|^2 = 1$ and $B_i u = \frac{1}{\sqrt{n}} e_i$, and hence $\|B_i u\| = \frac{1}{\sqrt{n}}$ for all $i = 1, \dots, n$. From Lemma 6.1, the joint spectrum of B_1, \dots, B_n must contain a vector in $[-1, 1]^n$. This bound is sharp: indeed, the joint spectrum is equal to the set $\{f_1, \dots, f_n, \sum_{i=1}^n f_j, \dots, \sum_{i=1}^n f_j\}$, where (f_1, \dots, f_n) is the canonical basis of \mathbb{R}^n : thus its intersection with $[-t, t]^n$ is empty for any $0 < t < 1$.

LEMMA 6.3. – *The following statements hold.*

- (i) *If B_1, \dots, B_n are commuting self-adjoint matrices, and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ is such that*

$$\|(B_i - \alpha_i)u\| \leq \epsilon \|u\|$$

for all $1 \leq i \leq n$, then there exists a joint eigenvalue $\lambda \in \text{JointSpec}(B_1, \dots, B_n)$ such that $|\lambda_i - \alpha_i| \leq \sqrt{n}\epsilon$ for all $1 \leq i \leq n$.

- (ii) *Suppose that A_1, \dots, A_n is another collection of commuting self-adjoint matrices, and assume*

$$\|B_i - A_i\| \leq \epsilon \quad \text{for all } 1 \leq i \leq n.$$

Then the Hausdorff distance between $\text{JointSpec}(A_1, \dots, A_n)$ and $\text{JointSpec}(B_1, \dots, B_n)$ is at most $\sqrt{n}\epsilon$, i.e.,

$$d_H(\text{JointSpec}(A_1, \dots, A_n), \text{JointSpec}(B_1, \dots, B_n)) \leq \sqrt{n}\epsilon.$$

Proof. – The first statement is obtained from Lemma 6.1 applied to $B_j - \alpha_j I$. It implies that if $\alpha = (\alpha_1, \dots, \alpha_n)$ is a joint eigenvalue of (A_1, \dots, A_n) and $\|B_i - A_i\| \leq \epsilon$ for all $1 \leq i \leq n$, then there exists a joint eigenvalue $\lambda \in \text{JointSpec}(B_1, \dots, B_n)$ with $|\lambda_i - \alpha_i| \leq \sqrt{n}\epsilon$ for all $1 \leq i \leq n$ (and vice-versa), which gives the last statement. \square

If T_1, \dots, T_n are pairwise commuting Toeplitz operators, we call *joint spectrum* of T_1, \dots, T_n the sequence of joint spectra of the set of commuting matrices $(T_{1,k}, \dots, T_{n,k})$ acting on the Hilbert space \mathcal{H}_k .

Proof of Theorem 1.1. – By Theorem 5.2 applied to (T_1, \dots, T_n) , we get an integer $k_0 > 0$, a sequence $(g(\cdot; k))_{k \geq k_0}$ of smooth maps $\mathbb{R}^n \rightarrow \mathbb{R}^n$, and an operator $U = (U_k : \mathcal{H}_k \rightarrow \mathcal{H}_k^\Delta)_{k \geq k_0}$ with U_k unitary, such that

$$(14) \quad U_k(T_{1,k}, \dots, T_{n,k})U_k^{-1} = g(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k) + \mathcal{O}(k^{-\infty}).$$

Let $S_k := \text{JointSpec}(T_1, \dots, T_n)$. If we introduce the components of g ,

$$g =: (g_1, \dots, g_n),$$

then by Corollary 4.3, the joint spectrum of the commuting Toeplitz operators

$$(g_1(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k), \dots, g_n(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k))$$

is $\Sigma_k := g((v + \frac{2\pi}{k}\mathbb{Z}^n) \cap \Delta, k)$.

Equation (14) means that there exists a sequence $(C_N)_{N \in \mathbb{N}}$ of real numbers such that

$$(15) \quad \text{for all } N, \text{ for all } k \quad \|U_k(T_{1,k}, \dots, T_{n,k})U_k^{-1} - g(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k)\| \leq C_N k^{-N}.$$

Then, by virtue of Lemma 6.3, we have

$$(16) \quad \text{for all } N, \text{ for all } k, \quad d_H(S_k, \Sigma_k) \leq \sqrt{n} C_N k^{-N}$$

which means by definition $S_k = \Sigma_k + \mathcal{O}(k^{-\infty})$ as we wanted to show.

Now we prove that for all sufficiently large k , the multiplicity of the eigenvalues of $\text{JointSpec}(T_1, \dots, T_n)$ is 1, and there exists a small constant $\delta > 0$ such that each ball of radius $\frac{\delta}{k}$ centered at an eigenvalue contains precisely only that eigenvalue. First we observe that this statement holds when instead of $\text{JointSpec}(T_1, \dots, T_n)$ we consider Σ_k . The multiplicity one statement is a direct consequence of Theorem 4.1, and the separation property follows from the fact that the principal term of g is the identity map.

Consider $\lambda = \lambda(k) \in S_k \subseteq \mathbb{R}^n$ and the self-adjoint operators

$$C_k := \sqrt{(T_{1,k} - \lambda_1 \text{Id}_k)^2 + \dots + (T_{n,k} - \lambda_n \text{Id}_k)^2}$$

and

$$C_k^\Delta := \sqrt{(T_{1,k}^\Delta - \lambda_1 \text{Id}_k)^2 + \dots + (T_{n,k}^\Delta - \lambda_n \text{Id}_k)^2}.$$

Let

$$0 = \lambda_1(C_k) \leq \lambda_2(C_k) \leq \dots \leq \lambda_{d_k}(C_k)$$

be the eigenvalues of C_k repeated with multiplicity, where d_k is the dimension of \mathcal{H}_k . Let

$$\lambda_1(C_k^\Delta) \leq \lambda_2(C_k^\Delta) \leq \dots \leq \lambda_{d_k}(C_k^\Delta)$$

be the eigenvalues of C_k^Δ repeated with multiplicity. By the min-max formula we have that for any $j = 1, \dots, d_k$,

$$|\lambda_j(C_k) - \lambda_j(C_k^\Delta)| = |\lambda_j(U_k C_k U_k^{-1}) - \lambda_j(C_k^\Delta)| \leq \|U_k C_k U_k^{-1} - C_k^\Delta\| \leq \frac{M}{k^2}$$

for some constant $M > 0$. Therefore for any $j > 1$ the inequality $|\lambda_j(C_k^\Delta) - \lambda_1(C_k^\Delta)| \geq \frac{\delta}{k}$ implies

$$|\lambda_j(C_k)| = |\lambda_j(C_k) - \lambda_1(C_k)| \geq \frac{\delta}{k} - \frac{2M}{k^2} \geq \frac{\delta'}{k}$$

for some $\delta' > 0$. Thus 0 as an eigenvalue of C_k has multiplicity one, and is the only eigenvalue of C_k in the interval $[-\frac{\delta'}{k}, \frac{\delta'}{k}]$. Going back to the commuting operators T_1, \dots, T_n , we see that the joint eigenvalue $\lambda \in S_k$ has multiplicity one, and all other joint eigenvalues lie outside of the Euclidean ball centered at λ of radius $\frac{\delta'}{k}$. \square

Proof of Corollary 1.2. – Recall that we defined the limit of a sequence $(\mathcal{A}_k)_{k \in \mathbb{N}}$ of subsets of \mathbb{R}^n by

$$\lim \mathcal{A}_k := \left\{ c \in \mathbb{R}^n \mid \forall U \text{ neighborhood of } c, \exists k_0 \text{ such that } \forall k \geq k_0, U \cap \mathcal{A}_k \neq \emptyset \right\}.$$

We denote by $B(c, r)$ the open ball in \mathbb{R}^n centered at c and of radius r . Notice that if $\mathcal{A}_k \subset \mathcal{B}_k + B(0, Ck^{-1})$, for some constant C , then

$$\lim \mathcal{A}_k \subset \lim \mathcal{B}_k.$$

We use the same notation as in the proof of Theorem 1.1: $S_k = \text{JointSpec}(T_1, \dots, T_n)$ and

$$\Sigma_k = g\left(v + \frac{2\pi}{k}\mathbb{Z}^n \cap \Delta, k\right).$$

Step 1. – The estimate (15) for $N = 1$ gives

$$S_k \subset \Sigma_k + B(0, Ck^{-1}) \quad \text{and} \quad \Sigma_k \subset S_k + B(0, Ck^{-1}).$$

Therefore, $\lim S_k = \lim \Sigma_k$.

Step 2. – Now let us show that $\lim \Sigma_k = \Delta$. From Theorem 5.2, we know that g admits an asymptotic expansion in the C^∞ topology of the form

$$g(\cdot; k) = \text{Id} + k^{-1}g_1 + k^{-2}g_2 + \dots.$$

Therefore, for any compact $K \subset \mathbb{R}^n$, there exists a constant C such that

$$\max_{c \in K} \|g(c; k) - c\|_{\mathbb{R}^n} \leq Ck^{-1}.$$

We may choose K large enough so that it contains Δ , and we get the following inclusions :

$$(17) \quad \Sigma_k \subset \Delta \cap \left(v + \frac{2\pi}{k}\mathbb{Z}^n\right) + B(0, Ck^{-1}) \quad \text{and} \quad \Delta \cap \left(v + \frac{2\pi}{k}\mathbb{Z}^n\right) \subset \Sigma_k + B(0, Ck^{-1}).$$

Hence

$$\lim \Sigma_k = \lim \Delta \cap \left(v + \frac{2\pi}{k}\mathbb{Z}^n\right) = \Delta.$$

From steps 1 and 2 we conclude that

$$\lim S_k = \Delta$$

which finishes the proof. \square

COROLLARY 6.4 (Isospectrality). – *Two symplectic toric systems are isomorphic if and only if the limit of their joint spectra coincide. In particular, if two symplectic toric systems have the same joint spectra, then they are isomorphic.*

7. Metaplectic correction

Theorem 1.1 and Theorem 1.2 can be proven analogously to our proofs in the presence of a half form bundle. Below we explain the corresponding modifications needed.

Toeplitz quantization is often considered more natural in the presence of a half-form bundle. Let (M, ω, j) be a compact Kähler manifold. A half-form bundle of (M, j) is a square root of the canonical bundle of M . More precisely we consider a pair (δ, φ) consisting of a complex line bundle $\delta \rightarrow M$ and an isomorphism

$$\varphi : \delta^{\otimes 2} \rightarrow \wedge^{n,0} T^* M.$$

Here n is the complex dimension of M . Such a square root does not necessarily exist, and if it exists, the space of half-form bundles up to isomorphism is a principal homogeneous space for the group $H^1(M, \mathbb{Z}/2\mathbb{Z})$.

Let $\mathcal{L} \rightarrow M$ be a prequantum bundle and (δ, φ) be a half-form bundle. Observe that δ has a metric and a holomorphic structure determined by the condition that φ is an isomorphism of Hermitian holomorphic bundles. Define the quantum space $\mathcal{H}_{m,k}$ as the vector space of holomorphic sections of $\mathcal{L}^k \otimes \delta$. The space $\mathcal{H}_{m,k}$ has a natural scalar product obtained by integrating the pointwise scalar product of sections of $\mathcal{L}^k \otimes \delta$ against the Liouville measure. This scalar product is actually defined on the space of L^2 sections, and we have an orthogonal projector Π_k from the space of L^2 sections onto $\mathcal{H}_{m,k}$. The definition of Toeplitz operator is the same as before except that we use this new projector. So a Toeplitz operator is any family $(T_k : \mathcal{H}_{m,k} \rightarrow \mathcal{H}_{m,k})_{k \in \mathbb{N}^*}$ of endomorphisms of the form

$$T_k = \Pi_k f(\cdot, k) + R_k, \quad k \in \mathbb{N}^*$$

where $f(\cdot, k)$, viewed as a multiplication operator, is a sequence in $C^\infty(M)$ with an asymptotic expansion

$$f_0 + k^{-1} f_1 + \dots$$

for the C^∞ topology, and the norm of R_k is $\mathcal{O}(k^{-\infty})$. Theorem 5.1 still holds and we define the principal and subprincipal symbols of a Toeplitz operator with the same formula as before. The rule of composition of these symbols is still given by (18). We can also define Toeplitz operators by using the Kostant-Souriau formula. Consider a smooth function f on M whose Hamiltonian vector field X preserves the complex structure. Then the following operators are well defined

$$(18) \quad T_k = f + \frac{1}{ik} (\nabla_X^{\mathcal{L}^k} \otimes \text{Id} + \text{Id} \otimes L_X^\delta) : \mathcal{H}_{m,k} \rightarrow \mathcal{H}_{m,k}, \quad k \in \mathbb{N}^*.$$

Here L^δ is the Lie derivative of the half-form, that is the first order differential operator such that for any local section s of the half-form bundle one has

$$L_X(\varphi(s^{\otimes 2})) = 2\varphi(s \otimes L_X^\delta s).$$

One also shows that the family (T_k) is a Toeplitz operator with principal symbol f and vanishing subprincipal symbol.

Consider now the Delzant space M_Δ defined as in Section 2. Since M_Δ is simply connected, there exists at most one half-form bundle.

Recall that we denote by F the set of faces of Δ of codimension 1 and for each $f \in F$, $X_f \in \mathfrak{t}$ is the primitive normal vector to the face.

PROPOSITION 7.1. – *The Delzant space M_Δ admits an equivariant half-form bundle if and only if there exists $\gamma \in \mathfrak{t}_\mathbb{Z}^*$ such that $\gamma(X_f)$ is odd for any $f \in F$.*

Proof. – This criterion may be established using the divisor of the toric variety, cf. [17]. To each face $f \in F$ corresponds an irreducible divisor D_f of M_Δ . It is known that a divisor of the canonical bundle is $-\sum_{f \in F} D_f$. Recall also that the divisors D_f 's generate the Picard group, and that $\sum n_f X_f$ is principal if and only if $n_f = \gamma(X_f)$ for some $\gamma \in \mathfrak{t}_\mathbb{Z}^*$. So M_Δ admits a half-form bundle if and only if there exists a divisor

$$D = \sum n_f^\delta D_f$$

such that $2D + \sum_{f \in F} D_f$ is principal, that is

$$2n_f^\delta + 1 = \langle X_f, \gamma \rangle$$

for some $\gamma \in \mathfrak{t}_\mathbb{Z}^*$. □

Assume now that the vertices of Δ belongs to $2\pi\mathfrak{t}_\mathbb{Z}^*$ so that $(M_\Delta, \omega_{M_\Delta})$ admits a prequantum bundle \mathcal{L}_Δ unique up to isomorphism. Assume also that M_Δ is equipped with a half-form bundle δ_Δ . For any positive integer k , define the quantum space

$$\mathcal{H}_{\mathfrak{m},k}^\Delta = H^0(M_\Delta, \mathcal{L}_\Delta^k \otimes \delta_\Delta).$$

For any $X \in \mathfrak{t}$, consider the rescaled Kostant-Souriau operators

$$(19) \quad T_{X,k} := \langle \mu_\Delta, X \rangle + \frac{1}{ik} \left(\nabla_{X^\#} \otimes \text{Id} + \text{Id} \otimes L_X^\delta \right) : \mathcal{H}_{\mathfrak{m},k}^\Delta \rightarrow \mathcal{H}_{\mathfrak{m},k}^\Delta.$$

We can now state the analogue of Theorem 4.1. Introduce $\gamma \in \mathfrak{t}_\mathbb{Z}^*$ such that $\gamma(X_f)$ is odd for any $f \in F$.

THEOREM 7.2. – *There is an orthogonal decomposition of the quantum space $\mathcal{H}_{\mathfrak{m},k}^\Delta$ into a direct sum of lines:*

$$\mathcal{H}_{\mathfrak{m},k}^\Delta = \bigoplus_{\ell \in (\frac{2\pi}{k}(\mathfrak{t}_\mathbb{Z}^* + \frac{1}{2}\gamma)) \cap \Delta} D_\ell^k$$

such that, for any $X \in \mathfrak{t}$,

$$T_{X,k}\Psi = \ell(X)\Psi, \quad \text{for all } \Psi \in D_\ell^k.$$

Observe that the points of $(\frac{2\pi}{k}(\mathfrak{t}_\mathbb{Z}^* + \frac{1}{2}\gamma)) \cap \Delta$ are all in the interior of Δ . Furthermore on a neighborhood of each vertex we recover the usual joint spectrum of n harmonic oscillators as follows. For any vertex v denote by F_v the set of one-codimensional faces adjacent to v so that $(f_v, v \in F_v)$ is a basis of $\mathfrak{t}_\mathbb{Z}^*$. Then there exists a neighborhood U of v such that

$$\Delta \cap U = \{v + x / \langle f, x \rangle \geq 0, \forall f \in F_v\} \cap U$$

and

$$(\frac{2\pi}{k}(\mathfrak{t}_\mathbb{Z}^* + \frac{1}{2}\gamma)) \cap \Delta \cap U = \{v + x / \langle f, x \rangle \in \frac{2\pi}{k}(\mathbb{N} + \frac{1}{2}), \forall f \in F_v\} \cap U.$$

Proof. – Let us adapt the proof of Theorem 4.1. First we introduce the quantization of \mathbb{C}^F with metaplectic correction. Choose an ordering of F and define $v \in \Omega(\mathbb{C}^F)$ as the wedge product of the dz_f 's. v is a non vanishing section of the canonical bundle of \mathbb{C}^F . The action of $t \in \mathbb{R}^F/\mathbb{Z}^F$ on the canonical bundle sends v into $\exp(-2i\pi \sum t_f)v$.

Let δ_F be the trivial complex line bundle with base \mathbb{C}^F and φ_F be the isomorphism from δ_F^2 to the canonical bundle of \mathbb{C}^F given by $\varphi(z, 1) = v(z)$. By Proposition 7.1, there exist $d \in \mathbb{Z}^F$ and $\gamma \in \mathfrak{t}_{\mathbb{Z}}^*$ such that

$$\pi^*\gamma = \mathbb{I} + 2d$$

where \mathbb{I} is the vector in $(\mathbb{R}^F)^*$ with all components equal to 1. Consider the action of $\mathbb{R}^F/\mathbb{Z}^F$ on δ_F given by

$$t \cdot (z, u) = (t \cdot z, e^{2i\pi \sum t_f d_f} u)$$

Then the isomorphism φ intertwines the action of the subtorus N of $\mathbb{R}^F/\mathbb{Z}^F$ on δ_F with the action of N on the canonical bundle. This condition has the consequence that the quotient of δ_F^2 by N defines a half-form bundle δ_{Δ} on the Delzant space M_{Δ} , cf. Section 8.3 of [12]. Furthermore one has an isomorphism

$$\Phi_k : (\mathcal{B}_{m,k})^N \rightarrow \mathcal{H}_{m,k}^{\Delta}$$

from the N -invariant part of the space $\mathcal{B}_{m,k}$ of holomorphic sections of $\mathcal{L}_F^k \otimes \delta_F$ to the quantum space $\mathcal{H}_{m,k}^{\Delta}$.

Introduce the rescaled Kostant-Souriau operators

$$S_{X,k} := \langle \tilde{\mu}, X \rangle + \frac{1}{ik} (\nabla_{X^{\sharp}}^{\mathcal{L}_F^k} \otimes \text{Id} + \text{Id} \otimes L_{X^{\sharp}}^{\delta}).$$

Then $(\mathcal{B}_{m,k})^N$ consists of the sections $\Psi \in \mathcal{B}_{m,k}$ such that $S_{X,k}\Psi = 0$ for any $X \in \mathfrak{n}$. Furthermore

$$\Phi_k(S_{X,k}\Psi) = T_{\pi(X),k}\Phi_k(\Psi), \quad \text{for all } \Psi \in (\mathcal{B}_{m,k})^N.$$

To conclude the proof let us compute the action of the rescaled Kostant-Souriau operators. We have for $X = e_f$

$$S_{X,k}(e^{-\frac{k}{8\pi}|z|^2}g(z)) = e^{-\frac{k}{8\pi}|z|^2} \left(\frac{2\pi}{k} (z_f \partial_{z_f} g + \frac{1}{2}g) - \lambda_f g \right).$$

For any $\alpha \in \mathbb{N}^F$ set $\psi_{\alpha} = e^{-\frac{k}{2}|z|^2} z^{\alpha}$ so that

$$(20) \quad S_{X,k}(\psi_{\alpha}) = \langle X, \frac{2\pi}{k}(\alpha + \frac{1}{2}\mathbb{I}) - \lambda \rangle \psi_{\alpha}.$$

Hence the space $(\mathcal{B}_{m,k})^N$ admits as a basis the family (Ψ_{α}) where α runs over $(\mathbb{Z}^F + \frac{1}{2}\mathbb{I}) \cap \ker \iota_{\mathfrak{n}}^*$. To conclude we prove as in Lemma 4.2 that $\pi^* + \lambda$ restricts to a bijection

$$\frac{2\pi}{k}(\mathfrak{t}_{\mathbb{Z}}^* + \frac{1}{2}\gamma) \longrightarrow \frac{2\pi}{k}(\mathbb{Z}^F + \frac{1}{2}\mathbb{I}) \cap \ker \iota_{\mathfrak{n}}^*. \quad \square$$

Let us consider now the generalization of Theorem 5.2. Let $(M, \omega, \mathbb{R}^n/\mathbb{Z}^n, \mu)$ be a symplectic toric manifold equipped with a prequantum bundle \mathcal{L} , a compatible complex structure j and a half-form bundle δ . Denote by Δ the momentum polytope $\mu(M) \subset \mathbb{R}^n$. Let T_1, \dots, T_n be commuting self-adjoint Toeplitz operators of $\mathcal{H}_{m,k}$ whose principal symbols are the components of μ . Denote by f_1^i the subprincipal symbol of T_i .




FIGURE 5. Spectrum in Figure 2 with metaplectic correction.

THEOREM 7.3. – *There exists $k_0 > 0$, there exists a sequence $(g(\cdot; k))_{k \geq k_0}$ of smooth maps $\mathbb{R}^n \rightarrow \mathbb{R}^n$, and there exists an operator $U = (U_k : \mathcal{H}_k \rightarrow \mathcal{H}_k^\Delta)_{k \geq k_0}$ with U_k unitary for any k , such that*

$$U_k(T_{1,k}, \dots, T_{n,k})U_k^{-1} = g(T_{1,k}^\Delta, \dots, T_{n,k}^\Delta; k) + \mathcal{O}(k^{-\infty}).$$

Moreover, g admits an asymptotic expansion in the C^∞ topology of the form $\text{Id} + k^{-1}g_1 + k^{-2}g_2 + \dots$ where the subprincipal term g_1 is given by

$$g_1^i(E) = \int_1^0 f_1^i(\varphi_{\mu_i}^t(x)) dt, \quad \text{for all } E \in \Delta, x \in \mu^{-1}(E).$$

Here $\varphi_{\mu_i}^t$ is the Hamiltonian flow of μ_i .

Proof. – The proof is the same as the one of Theorem 5.2. Because of the metaplectic correction, we can choose in the first step the operator U_k quantizing φ in such a way that for any Toeplitz operator (S_k) , (S_k) and $(U_k S_k U_k^*)$ have the same principal and subprincipal symbols. This follows from Theorem 5.1 in [11]. From this we can extract g_1 from the cohomological equation

$$\{\mu_i, a\} = f_1^i - g_1^i(\mu_\Delta). \quad \square$$

Given the statement of Theorem 7.3 above, the proof of Theorem 1.3 is identical to the proof of Theorem 1.1 given in Section 6.

8. Final Remarks

Isospectrality in geometry

In the present paper we have dealt with isospectrality in the context of integrable systems and symplectic geometry. The paper settles the *Spectral Goal for Quantum Systems* for the case of toric systems outlined in the last two authors' article [60]: to prove that large classes of integrable systems are determined by their semiclassical joint spectrum.

Corollary 1.2 says that the joint spectrum does indeed determine the system. This type of conclusion often has a negative answer, at least if one considers it in Riemannian geometry. In Riemannian geometry the operator whose spectrum is considered is the Laplace operator. Two compact Riemannian manifolds are said to be *isospectral* if the associated Laplace operators have the same spectrum.

Bochner and Kac's question has a negative answer in this case, even for planar domains with Dirichlet boundary conditions (which is the original version posed in [41]). As P. Sarnak

mentioned to us, a much better question to ask is whether the set of isospectral domains is finite. There are many related works, see for instance Milnor [52], Sunada [64], Osgood-Phillips-Sarnak [54, 55, 56], Bérard [2], Buser [4] and Gordon-Webb-Wolpert [32].

As we have mentioned, in symplectic geometry a few positive results are known. These results, and the present paper, give evidence that symplectic invariants seem to be more encodable in the spectrum of a quantum integrable system than Riemannian invariants in the spectrum of the Laplace operator.

Inverse type results in the realm of spectral geometry have been obtained by many other authors, see for instance Brüning-Heintze [3], Colin de Verdière [14, 13, 15], Colin de Verdière-Guillemin [16], Croke-Sharafutdinov [18], Guillemin-Kazhdan [35], McKean-Singer [50], Osgood-Phillips-Sarnak [54, 55, 56], and Zelditch [75], and the references therein. An interesting general problem (for instance in the context of toric geometry) is to what extent information about measures may be recovered from the spectrum, see Guillemin-Sternberg [38, p. 268-271] for a result in this direction.

Isospectral conjecture for semitoric systems

In [61, 59] the last two authors formulated an inverse spectral conjecture for semitoric completely integrable systems (see [57, 58] for a classification of semitoric systems in terms of five symplectic invariants): the semiclassical joint spectrum of a quantum semitoric system determines the corresponding classical system.




FIGURE 6. The quantum coupled spin-oscillator is a fundamental example of quantum semitoric integrable system. Its joint spectrum is depicted in the figure for a fixed value of the spectral parameter.

Semitoric systems are four-dimensional integrable systems with two degrees of freedom for which one component of the system generates a 2π -periodic flow. Semitoric systems lie somewhere in between toric systems and general integrable systems. If both components of the semitoric system are 2π -periodic, i.e., the system is generated by a Hamiltonian 2-torus action, then the system is a *toric system* (strictly speaking after a harmless rescaling of the periods).

Theorem 1.2 above solves the conjecture for the class of toric systems. In this class the result is even stronger, since there is no restriction on the dimension, and moreover only the

spectrum modulo $\mathcal{O}(\hbar)$ is needed, whereas in general one expects that an accuracy of order $\mathcal{O}(\hbar^2)$ is necessary.

Microlocal analysis of integrable systems

The notion of a quantum integrable system, as a maximal set of commuting quantum observables, dates back to the early quantum mechanics, to the works of Bohr, Sommerfeld and Einstein [29]. However, the most basic results in the symplectic theory of classical integrable systems like Darboux's theorem or action-angle variables could not be used in Schrödinger's quantum setting at that time because they make use of the analysis of differential (or pseudodifferential) operators in phase space, known now as microlocal analysis, which was developed only in the 1960s. The microlocal analysis of action-angle variables starts with the works of Duistermaat [23] and Colin de Verdière [14, 13], followed by the semiclassical theory by Charbonnel [5], and more recently by Vũ Ngọc [69], Zelditch and Toth [65, 66, 67], Charbonnel and Popov [6], Melin-Sjöstrand [51], and many others.

Effective models in quantum mechanics often require a compact phase space, and thus cannot be treated using pseudodifferential calculus. For instance the natural classical limit of a quantum spin is a symplectic sphere. The study of quantum action-angle variables in the case of compact symplectic manifolds treated in this paper was started by Charles [8], using the theory of Toeplitz operators. In the present paper, we present global spectral results for toric integrable systems; we use in a fundamental way Delzant's theorem on symplectic toric manifolds [21].

Toric integrable systems ⁽²⁾ have played an influential role in symplectic and complex algebraic geometry, representation theory, and spectral theory since T. Delzant classified them in terms of combinatorial information (actually, in terms of a convex polytope, see Theorem 2.3). A comparative study of symplectic toric manifolds from the symplectic and complex algebraic viewpoints was given by Duistermaat and Pelayo [28]. A beautiful treatment of the classical theory of toric systems is given in Guillemin's book [34].

Toeplitz operators are a natural generalization of Toeplitz matrices (which correspond to Toeplitz operators on the unit disk). On \mathbb{R}^n , Toeplitz operators correspond to pseudodifferential operators via the Bargmann transform. Of course, such a correspondence cannot hold in the case of a compact phase space, but it turns out that Toeplitz operators always give rise to a semiclassical algebra of operators with a symbolic calculus and microlocalization properties, which is microlocally isomorphic to the algebra of pseudodifferential operators. See the book by Boutet de Monvel and Guillemin for an introduction to the spectral theory of Toeplitz operators [53].

⁽²⁾ Toric integrable systems always have singularities of elliptic and transversally elliptic type, but do not have singularities of hyperbolic or focus-focus type. The local and semiglobal theory for regular points, elliptic and transversally elliptic singularities is now well understood both at the symplectic level (action-angle theorem of Liouville-Arnold-Mineur, Eliasson's linearization theorems), as well as the quantum level, see Charles [8] and Vũ Ngọc [70]. This gives the foundation of the modern theory of integrable systems, in the spirit of Duistermaat's article [24], but also of KAM-type perturbation theorems (see de la Llave's survey article [48]).

Acknowledgements

We are very grateful to a referee for many insightful comments which have improved the paper, in particular for pointing out an omission of a constant in one of the lemmas (Lemma 6.1) which affected the constants in Lemma 6.3.

We thank Jochen Brüning, Helmut Hofer, Peter Sarnak, and Thomas Spencer for fruitful discussions. The authors are grateful to Helmut Hofer for his essential support, which made it possible for LC and VNS to visit AP at the Institute for Advanced Study during the Winter and Summer of 2011, where a part of this paper was written. Additional financial support for the visits was provided by Washington University and NSF.

AP was partly supported by an NSF Postdoctoral Fellowship, a MSRI membership, an IAS membership, NSF Grants DMS-0965738 and DMS-0635607, an NSF CAREER Award, a Leibniz Fellowship from the Mathematisches Forschungsinstitut Oberwolfach, Spanish Ministry of Science Grant MTM 2010-21186-C02-01, and by the Spanish National Research Council (CSIC). VNS was partly supported by the NONA, a grant from the French ANR and the Institut Universitaire de France.

REFERENCES

- [1] M. F. ATIYAH, Convexity and commuting Hamiltonians, *Bull. London Math. Soc.* **14** (1982), 1–15.
- [2] P. BÉRARD, Transplantation et isospectralité. I, *Math. Ann.* **292** (1992), 547–559.
- [3] J. BRÜNING, E. HEINTZE, Spektrale Starrheit gewisser Drehflächen, *Math. Ann.* **269** (1984), 95–101.
- [4] P. BUSER, Isospectral Riemann surfaces, *Ann. Inst. Fourier (Grenoble)* **36** (1986), 167–192.
- [5] A.-M. CHARBONNEL, Comportement semi-classique du spectre conjoint d’opérateurs pseudodifférentiels qui commutent, *Asymptotic Anal.* **1** (1988), 227–261.
- [6] A.-M. CHARBONNEL, G. POPOV, A semi-classical trace formula for several commuting operators, *Comm. Partial Differential Equations* **24** (1999), 283–323.
- [7] L. CHARLES, Berezin-Toeplitz operators, a semi-classical approach, *Comm. Math. Phys.* **239** (2003), 1–28.
- [8] L. CHARLES, Quasimodes and Bohr-Sommerfeld conditions for the Toeplitz operators, *Comm. Partial Differential Equations* **28** (2003), 1527–1566.
- [9] L. CHARLES, Symbolic calculus for Toeplitz operators with half-form, *J. Symplectic Geom.* **4** (2006), 171–198.
- [10] L. CHARLES, Toeplitz operators and Hamiltonian torus actions, *J. Funct. Anal.* **236** (2006), 299–350.
- [11] L. CHARLES, Semi-classical properties of geometric quantization with metaplectic correction, *Comm. Math. Phys.* **270** (2007), 445–480.

- [12] L. CHARLES, On the quantization of polygon spaces, *Asian J. Math.* **14** (2010), 109–152.
- [13] Y. COLIN DE VERDIÈRE, Spectre conjoint d'opérateurs pseudo-différentiels qui commutent. I. Le cas non intégrable, *Duke Math. J.* **46** (1979), 169–182.
- [14] Y. COLIN DE VERDIÈRE, Spectre conjoint d'opérateurs pseudo-différentiels qui commutent. II. Le cas intégrable, *Math. Z.* **171** (1980), 51–73.
- [15] Y. COLIN DE VERDIÈRE, A semi-classical inverse problem II: reconstruction of the potential, in *Geometric aspects of analysis and mechanics*, Progr. Math. **292**, Birkhäuser, 2011, 97–119.
- [16] Y. COLIN DE VERDIÈRE, V. GUILLEMIN, A semi-classical inverse problem I: Taylor expansions, in *Geometric aspects of analysis and mechanics*, Progr. Math. **292**, Birkhäuser, 2011, 81–95.
- [17] D. A. COX, J. B. LITTLE, H. K. SCHENCK, *Toric varieties*, Graduate Studies in Math. **124**, Amer. Math. Soc., 2011.
- [18] C. B. CROKE, V. A. SHARAFUTDINOV, Spectral rigidity of a compact negatively curved manifold, *Topology* **37** (1998), 1265–1273.
- [19] V. I. DANILOV, The geometry of toric varieties, *Uspekhi Mat. Nauk* **33** (1978), 85–134, 247.
- [20] K. DATCHEV, H. HEZARI, I. VENTURA, Spectral uniqueness of radial semiclassical Schrödinger operators, *Math. Res. Lett.* **18** (2011), 521–529.
- [21] T. DELZANT, Hamiltoniens périodiques et images convexes de l'application moment, *Bull. Soc. Math. France* **116** (1988), 315–339.
- [22] E. B. DRYDEN, V. GUILLEMIN, R. SENA-DIAS, Hearing Delzant polytopes from the equivariant spectrum, to appear in *Trans. Amer. Math. Soc.*
- [23] J. J. DUISTERMAAT, Oscillatory integrals, Lagrange immersions and unfolding of singularities, *Comm. Pure Appl. Math.* **27** (1974), 207–281.
- [24] J. J. DUISTERMAAT, On global action-angle coordinates, *Comm. Pure Appl. Math.* **33** (1980), 687–706.
- [25] J. J. DUISTERMAAT, Principal fiber bundles, in *Notes for the Utrecht Spring School*, 2004, <http://www.projects.science.uu.nl/Duistermaat/www/homepageHD/pf.pdf>.
- [26] J. J. DUISTERMAAT, *The heat kernel Lefschetz fixed point formula for the spin-c Dirac operator*, Modern Birkhäuser Classics, Birkhäuser, 2011.
- [27] J. J. DUISTERMAAT, L. HÖRMANDER, Fourier integral operators. II, *Acta Math.* **128** (1972), 183–269.
- [28] J. J. DUISTERMAAT, Á. PELAYO, Reduced phase space and toric variety coordinatizations of Delzant spaces, *Math. Proc. Cambridge Philos. Soc.* **146** (2009), 695–718.
- [29] A. EINSTEIN, Zum Quantensatz von Sommerfeld und Epstein, *Verhandlungen der Deutschen Physikalischen Gesellschaft* **19** (1917), 82–92.
- [30] L. H. ELIASSON, Normal forms for Hamiltonian systems with Poisson commuting integrals—elliptic case, *Comment. Math. Helv.* **65** (1990), 4–35.
- [31] M. D. GARAY, D. VAN STRATEN, Classical and quantum integrability, *Mosc. Math. J.* **10** (2010), 519–545, 661.

- [32] C. GORDON, D. L. WEBB, S. WOLPERT, Isospectral plane domains and surfaces via Riemannian orbifolds, *Invent. Math.* **110** (1992), 1–22.
- [33] C. GORDON, D. L. WEBB, S. WOLPERT, One cannot hear the shape of a drum, *Bull. Amer. Math. Soc. (N.S.)* **27** (1992), 134–138.
- [34] V. GUILLEMIN, *Moment maps and combinatorial invariants of Hamiltonian T^n -spaces*, Progress in Math. **122**, Birkhäuser, 1994.
- [35] V. GUILLEMIN, D. KAZHDAN, Some inverse spectral results for negatively curved 2-manifolds, *Topology* **19** (1980), 301–312.
- [36] V. GUILLEMIN, S. STERNBERG, Convexity properties of the moment mapping, *Invent. Math.* **67** (1982), 491–513.
- [37] V. GUILLEMIN, S. STERNBERG, Geometric quantization and multiplicities of group representations, *Invent. Math.* **67** (1982), 515–538.
- [38] V. GUILLEMIN, S. STERNBERG, *Symplectic techniques in physics*, second ed., Cambridge Univ. Press, 1990.
- [39] L. HÖRMANDER, Fourier integral operators. I, *Acta Math.* **127** (1971), 79–183.
- [40] A. IANTCHENKO, J. SJÖSTRAND, M. ZWORSKI, Birkhoff normal forms in semi-classical inverse problems, *Math. Res. Lett.* **9** (2002), 337–362.
- [41] M. KAC, Can one hear the shape of a drum?, *Amer. Math. Monthly* **73** (1966), 1–23.
- [42] Y. KARSHON, L. KESSLER, M. PINSONNAULT, A compact symplectic four-manifold admits only finitely many inequivalent toric actions, *J. Symplectic Geom.* **5** (2007), 139–166.
- [43] B. KOSTANT, Orbits, symplectic structures and representation theory, in *Proc. U.S.-Japan Seminar in Differential Geometry (Kyoto, 1965)*, Nippon Hyoronsha, Tokyo, 1966, p. 71.
- [44] B. KOSTANT, Quantization and unitary representations. I. Prequantization, in *Lectures in modern analysis and applications, III*, Springer, 1970, 87–208. Lecture Notes in Math., Vol. 170.
- [45] B. KOSTANT, Symplectic spinors, in *Symposia Mathematica, Vol. XIV (Convegno di Geometria Simplettica e Fisica Matematica, INDAM, Rome, 1973)*, Academic Press, 1974, 139–152.
- [46] B. KOSTANT, On the definition of quantization, in *Géométrie symplectique et physique mathématique (Colloq. Internat. CNRS, No. 237, Aix-en-Provence, 1974)*, Éditions du CNRS, Paris, 1975, 187–210.
- [47] B. KOSTANT, Á. PELAYO, Geometric quantization, a Lie theory approach, to appear as *Springer Universitext*.
- [48] R. DE LA LLAVE, A tutorial on KAM theory, in *Smooth ergodic theory and its applications (Seattle, WA, 1999)*, Proc. Sympos. Pure Math. **69**, Amer. Math. Soc., 2001, 175–292.
- [49] I. MADSEN, J. TORNEHAVE, *From calculus to cohomology*, Cambridge Univ. Press, 1997.
- [50] H. P. J. MCKEAN, I. M. SINGER, Curvature and the eigenvalues of the Laplacian, *J. Differential Geometry* **1** (1967), 43–69.
- [51] A. MELIN, J. SJÖSTRAND, Bohr-Sommerfeld quantization condition for non-selfadjoint operators in dimension 2, *Astérisque* **284** (2003), 181–244.

- [52] J. MILNOR, Eigenvalues of the Laplace operator on certain manifolds, *Proc. Nat. Acad. Sci. U.S.A.* **51** (1964), 542.
- [53] L. BOUTET DE MONVEL, V. GUILLEMIN, *The spectral theory of Toeplitz operators*, Annals of Math. Studies **99**, Princeton Univ. Press, 1981.
- [54] B. OSGOOD, R. PHILLIPS, P. SARNAK, Compact isospectral sets of plane domains, *Proc. Nat. Acad. Sci. U.S.A.* **85** (1988), 5359–5361.
- [55] B. OSGOOD, R. PHILLIPS, P. SARNAK, Compact isospectral sets of surfaces, *J. Funct. Anal.* **80** (1988), 212–234.
- [56] B. OSGOOD, R. PHILLIPS, P. SARNAK, Moduli space, heights and isospectral sets of plane domains, *Ann. of Math.* **129** (1989), 293–362.
- [57] Á. PELAYO, S. VŨ NGỌC, Semitoric integrable systems on symplectic 4-manifolds, *Invent. Math.* **177** (2009), 571–597.
- [58] Á. PELAYO, S. VŨ NGỌC, Constructing integrable systems of semitoric type, *Acta Math.* **206** (2011), 93–125.
- [59] Á. PELAYO, S. VŨ NGỌC, Symplectic theory of completely integrable Hamiltonian systems, *Bull. Amer. Math. Soc. (N.S.)* **48** (2011), 409–455.
- [60] Á. PELAYO, S. VŨ NGỌC, First steps in symplectic and spectral theory of integrable systems, *Discrete Contin. Dyn. Syst.* **32** (2012), 3325–3377.
- [61] Á. PELAYO, S. VŨ NGỌC, Hamiltonian dynamics and spectral theory for spin-oscillators, *Comm. Math. Phys.* **309** (2012), 123–154.
- [62] J.-M. SOURIAU, Quantification géométrique, *Comm. Math. Phys.* **1** (1966), 374–398.
- [63] J.-M. SOURIAU, *Structure des systèmes dynamiques*, Maîtrise de mathématiques, Dunod, 1970.
- [64] T. SUNADA, Riemannian coverings and isospectral manifolds, *Ann. of Math.* **121** (1985), 169–186.
- [65] J. A. TOTH, On the quantum expected values of integrable metric forms, *J. Differential Geom.* **52** (1999), 327–374.
- [66] J. A. TOTH, S. ZELDITCH, Riemannian manifolds with uniformly bounded eigenfunctions, *Duke Math. J.* **111** (2002), 97–132.
- [67] J. A. TOTH, S. ZELDITCH, L^p norms of eigenfunctions in the completely integrable case, *Ann. Henri Poincaré* **4** (2003), 343–368.
- [68] G. M. TUYNMAN, Quantization: towards a comparison between methods, *J. Math. Phys.* **28** (1987), 2829–2840.
- [69] S. VŨ NGỌC, Bohr-Sommerfeld conditions for integrable systems with critical manifolds of focus-focus type, *Comm. Pure Appl. Math.* **53** (2000), 143–217.
- [70] S. VŨ NGỌC, Systèmes intégrables semi-classiques : du local au global, *Panoramas et Synthèses* **22** (2006).
- [71] S. VŨ NGỌC, Symplectic inverse spectral theory for pseudodifferential operators, in *Geometric aspects of analysis and mechanics*, Progr. Math. **292**, Birkhäuser, 2011, 353–372.
- [72] H. WEYL, Über die asymptotische Verteilung der Eigenwerte, *Göttinger Nachrichten* (1911), 110–117.

- [73] H. WEYL, Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung), *Math. Ann.* **71** (1912), 441–479.
- [74] S. ZELDITCH, Index and dynamics of quantized contact transformations, *Ann. Inst. Fourier (Grenoble)* **47** (1997), 305–363.
- [75] S. ZELDITCH, Inverse spectral problem for analytic domains. II. \mathbb{Z}_2 -symmetric domains, *Ann. of Math.* **170** (2009), 205–269.

(Manuscrit reçu le 9 mars 2012 ;
 accepté, après révision, le 13 novembre 2012.)

Laurent CHARLES
 Institut de Mathématiques de Jussieu
 Université Pierre et Marie Curie (Paris VI)
 Case 247
 4, place Jussieu
 F-75252 Paris Cedex 05
 E-mail: charles@math.jussieu.fr

Álvaro PELAYO
 School of Mathematics
 Institute for Advanced Study
 Einstein Drive
 Princeton, NJ 08540 USA.
 and
 Washington University
 Mathematics Department
 One Brookings Drive, Campus Box 1146
 St Louis, MO 63130-4899, USA
 E-mail: apelayo@math.wustl.edu

San Vũ NGỌC
 Institut Universitaire de France
 and
 Institut de Recherches Mathématiques de Rennes
 Université de Rennes 1
 Campus de Beaulieu
 F-35042 Rennes cedex, France
 E-mail: san.vu-ngoc@univ-rennes1.fr

