

**369**

**ASTÉRISQUE**

**2015**

DE LA GÉOMÉTRIE ALGÈBRIQUE  
AUX FORMES AUTOMORPHES (I)

J.-B. BOST, P. BOYER, A. GENESTIER,  
L. LAFFORGUE, S. LYSENKO, S. MOREL, B.C. NGÔ, eds.

---

FOURIER TRANSFORM OF ALGEBRAIC MEASURES

Vladimir DRINFELD

**SOCIÉTÉ MATHÉMATIQUE DE FRANCE**

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

---

Astérisque est un périodique de la Société Mathématique de France.

Numéro 369, 2015

---

***Comité de rédaction***

Ahmed ABBES	Damien GABORIAU
Viviane BALADI	Michael HARRIS
Gérard BESSON	Fabrice PLANCHON
Laurent BERGER	Pierre SCHAPIRA
Philippe BIANE	Bertrand TOËN
Hélène ESNAULT	
Éric VASSEROT (dir.)	

***Diffusion***

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

***Tarifs***

*Vente au numéro* : 82 € (\$ 123)

*Abonnement* Europe : 650 €, hors Europe : 689 € (\$ 1033)

Des conditions spéciales sont accordées aux membres de la SMF.

***Secrétariat : Nathalie Christiaën***

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2015

*Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.*

ISSN 0303-1179

ISBN 978-2-85629-805-3

Directeur de la publication : Marc Peigné

---

## FOURIER TRANSFORM OF ALGEBRAIC MEASURES

by

Vladimir Drinfeld

---

*To Gérard Laumon on his 60th birthday*

**Abstract.** — These are notes of a talk based on the work [AD] joint with A. Aizenbud.

Let  $V$  be a finite-dimensional vector space over a local field  $F$  of characteristic 0. Let  $f$  be a function on  $V$  of the form  $x \mapsto \psi(P(x))$ , where  $P$  is a polynomial on  $V$  and  $\psi$  is a nontrivial additive character of  $F$ . Then it is clear that the Fourier transform  $\text{Four}(f)$  is well-defined as a distribution on  $V^*$ . Due to J. Bernstein, Hrushovski-Kazhdan, and Cluckers-Loeser, it is known that  $\text{Four}(f)$  is smooth on a non-empty Zariski-open conic subset of  $V^*$ . The goal of these notes is to sketch a proof of this result (and some related ones), which is very simple modulo resolution of singularities (the existing proofs use D-module theory in the Archimedean case and model theory in the non-Archimedean one).

**Résumé (Transformation de Fourier de mesures algébriques).** — Ce sont les notes d'un exposé basé sur le travail [AD] commun avec A. Aizenbud.

Soit  $V$  un espace vectoriel de dimension finie sur un corps local  $F$  de caractéristique 0. Soit  $f$  une fonction sur  $V$  de la forme  $x \mapsto \psi(P(x))$ , où  $P$  est un polynôme sur  $V$  et  $\psi$  est un caractère additif non trivial de  $F$ . Alors il est clair que la transformée de Fourier  $\text{Four}(f)$  est bien définie comme distribution sur  $V^*$ . D'après J. Bernstein, Hrushovski-Kazhdan et Cluckers-Loeser, il est connu que  $\text{Four}(f)$  est lisse sur un sous-ensemble ouvert de Zariski conique de  $V^*$ . Le but de ces notes est d'esquisser une démonstration de ce résultat (et de résultats liés), qui est très simple modulo la résolution des singularités (les preuves existantes utilisent la théorie des D-modules dans le cas archimédien et la théorie des modèles dans le cas non archimédien).

These are notes of a talk based on the work [AD] joint with A. Aizenbud. The results from [AD] are formulated in §§1–3, the proofs are sketched in §§4–5.

In Appendix A we discuss some “baby examples”; this material is not contained in [AD].

---

**2010 Mathematics Subject Classification.** — 46F, 46F10.

**Key words and phrases.** — Wave front set, Fourier transform, distributions, oscillating integrals, resolution of singularities, local fields.

Partially supported by NSF grant DMS-1063470.

I thank D. Kazhdan for drawing my attention to the questions considered in these notes. I also thank M. Kashiwara for communicating to me Example A.1 from Appendix A.

## 1. A theorem on Fourier transform

Let  $F$  be a local field of characteristic 0 (Archimedean or not). Let  $\psi : F \rightarrow \mathbb{C}^\times$  be a nontrivial additive character.

Let  $V$  be a finite-dimensional vector space over  $F$  and  $P : V \rightarrow F$  a polynomial. Then  $\psi(P(x))$  is a smooth<sup>(1)</sup>  $\mathbb{C}$ -valued function on  $V$ .

Consider  $\widehat{\psi(P(x))}$ , *i.e.*, the Fourier transform of the function  $\psi(P(x))$ .

Note that in the naive sense the Fourier transform is not defined because  $\psi(P(x))$  doesn't decay as  $x \rightarrow \infty$ , rather it oscillates. However,  $\widehat{\psi(P(x))}$  is well-defined as a distribution<sup>(2)</sup> on  $V^*$ .

In the non-Archimedean case this is clear because the Fourier transform in the sense of distributions is well-defined for *any* generalized function, in particular, for any smooth function.

In the Archimedean case the Fourier transform is well-defined for generalized functions of *moderate growth*, and of course,  $\psi(P(x))$  has moderate growth.

**Theorem 1.1.** — *There exists a Zariski-open  $U \subset V^*$ ,  $U \neq \emptyset$ , such that the distribution  $\widehat{\psi(P(x))}$  is smooth on  $U$ .*

If  $F$  is Archimedean Theorem 1.1 was proved by J. Bernstein [Ber1] using *D-module theory*. Moreover, he proved that  $\widehat{\psi(P(x))}$  satisfies a holonomic system of linear p.d.e.'s with polynomial coefficients.

If  $F$  is non-Archimedean Theorem 1.1 was proved by Kazhdan-Hrushovski [HK] and Cluckers-Loeser [CL] using *model theory*. In both articles Theorem 1.1 appears as one of many corollaries of a general theory, and this general theory is quite different from D-module theory used by Bernstein.

The goal of these notes is to explain another proof of Theorem 1.1 and its refinements, namely the one from [AD]. It works equally well in the Archimedean and non-Archimedean case. Unlike the older proofs, it uses resolution of singularities<sup>(3)</sup>.

1. In the non-Archimedean case “smooth” means “locally constant”, in the non-Archimedean case the word “smooth” is understood literally.

2. Our conventions are as follows: a distribution on a manifold  $M$  is a generalized measure (*i.e.*, a linear functional on the space of smooth *functions* with compact support), while a generalized function on  $M$  is a linear functional on the space of smooth *measures* with compact support. If  $M$  is a vector space then sometimes (but not here) we do not distinguish functions from measures.

3. In my talk I said that a variant of “local uniformization” (see [Za], [ILO]) would suffice. But the argument that I had in mind contained a gap.

On the other hand, once you believe in resolution of singularities, the rest is an exercise in elementary analysis (with a bit of elementary symplectic geometry).

Here are some refinements of Theorem 1.1.

Refinement A. The open subset  $U$  can be chosen to be independent of  $\psi$ .

Refinement B. The open subset  $U$  can be chosen to be defined over the field  $K$  generated by the coefficients of  $P$ .

Refinement B'. The open subset  $U$  can be chosen to work for all embeddings of  $K$  into all possible local fields. (This makes sense by virtue of A.)

## 2. A theorem that implies Theorem 1.1

Let  $W$  be a finite-dimensional vector space over  $F$ . Let  $X$  be a smooth algebraic variety over  $F$  and  $\varphi : X \rightarrow W$  a proper morphism. Let  $\omega$  be a regular top differential form on  $X$ . Then we have a measure  $|\omega|$  on  $X(F)$ . Set  $\mu := \varphi_*|\omega|$  (note that  $\varphi_*$  is well-defined because  $\varphi$  is proper);  $\mu$  is a measure on the vector space  $W$ , in particular, it is a distribution. Its Fourier transform,  $\hat{\mu}$ , is a well-defined<sup>(4)</sup> generalized function on  $W^*$ ; it depends on the choice of  $\psi$ . The next theorem is an analog of Theorem 1.1 and its Refinements A,B,B'.

**Theorem 2.1.** — *There exists a non-empty Zariski-open  $U \subset W^*$ , independent of  $\psi$ , such that  $\hat{\mu}$  is smooth on  $U$ . Moreover, if  $(X, \varphi, \omega)$  is defined over a subfield  $K \subset F$  then one can choose  $U$  to be defined over  $K$  and to have the required property for all embeddings of  $K$  into all possible local fields.*

**Remark 2.2.** — In Theorem 2.1 independence of  $U$  on  $\psi$  is equivalent to *stability of  $U$  under homotheties* of  $W$ . (This was not the case in the situation of Theorem 1.1 because  $\psi$  occurred there twice: in the definition of Fourier transform and in the expression  $\psi(P(x))$ .)

Let us show that Theorem 2.1 implies Theorem 1.1 and its refinements formulated at the end of §1. To prove this, apply Theorem 2.1 as follows. Set  $W := V \oplus F = V \times F$ ,  $X := V$ . Define  $\varphi : V \rightarrow V \times F$  by  $\varphi(v) := (v, P(v))$ . Take  $\omega$  to be an invariant differential form on  $V = X$ .

Then the generalized function  $\hat{\mu}$  on  $W^* = V^* \times F$  is equal to the continuous map

$$F \longrightarrow \{\text{generalized functions on } V^*\}$$

that takes  $\eta \in F$  to the Fourier transform of  $\psi(\eta \cdot P(x))$  with respect to  $x \in V$ .

So Theorem 2.1 says that the Fourier transform of  $\psi(\eta \cdot P(x))$  with respect to *both*  $x$  and  $\eta$  (which is *a priori* a *generalized* function on  $V^* \times F$ ) is, in fact, *smooth* on some non-empty open subset  $U \subset V^* \times F$ ,  $U \neq \emptyset$ , which can be chosen to be stable

4. In the Archimedean case one has to check that  $\mu$  has moderate growth. This is not hard and well known.

with respect to homotheties (see Remark 2.2). It remains to note that such  $U$  has non-empty intersection with the hyperplane  $V^* \times \{1\} \subset V^* \times F$ .

Theorem 2.1 is deduced from Theorem 3.3, which is formulated in the next section.

### 3. The theorem on wave front sets

**3.1. Isotropic subsets of symplectic varieties.** — Let  $M$  be a symplectic algebraic manifold<sup>(5)</sup> over a field of characteristic 0. An algebraic subvariety (or more generally, a constructible subset) of  $M$  is said to be *isotropic* if each of its smooth subvarieties is. (As usual, a smooth subvariety of  $M$  is said to be isotropic if each of its tangent spaces is). It is known that the closure of an isotropic subvariety is isotropic (e.g., see [CG, Proposition 1.3.30] and [CG, §1.5.16]). So a subvariety of a symplectic variety is isotropic if and only if its smooth locus is.

Now suppose that  $M$  is the cotangent bundle of an algebraic manifold  $Y$ . A subset of  $M = T^*Y$  is said to be *conic* if it is stable with respect to the action of  $\mathbb{G}_m$  on  $T^*Y$ . The following two statements are easy and standard.

**Lemma 3.1.** — *A conic algebraic subvariety  $Z \subset T^*Y$  is isotropic if and only if there exists a finite collection of smooth locally closed subvarieties of  $Y$  such that  $Z$  is contained in the union of their conormal bundles.*

**Corollary 3.2.** — *Let  $Z \subset T^*Y$  be an isotropic conic algebraic subvariety. Then there exists a dense open subset  $U \subset T^*Y$  such that  $Z \cap T^*U$  is contained in the zero section.*

**3.2. The theorem on wave fronts.** — Let  $F$  be a local field of characteristic 0. Let  $Y$  be an analytic manifold<sup>(6)</sup> over  $F$ . (If  $F = \mathbb{R}$  one can assume  $Y$  to be a  $C^\infty$  manifold.) Following L. Hörmander [Hör] and D. Heifetz [Hef], to each distribution or generalized function  $\nu$  on  $Y$  one associates a conic closed subset of  $T^*Y$  called the *wave front set* of  $\nu$ . The precise definition will be recalled in §3.3 below.

In the situation of Theorem 2.1  $\hat{\mu}$  is a generalized function on  $W^*$ , so its wave front is a subset of  $T^*W^*$ . Note that  $T^*W^* = W^* \times W$  carries an action of  $\mathbb{G}_m \times \mathbb{G}_m$ .

**Theorem 3.3.** — *In the situation of Theorem 2.1 the wave front of  $\hat{\mu}$  is contained in  $I(F)$ , where  $I \subset T^*W^*$  is some isotropic algebraic subvariety stable with respect to  $\mathbb{G}_m \times \mathbb{G}_m$  and defined over  $K$  (as before,  $K$  is a field of definition of our data). Moreover, one can choose  $I$  to have the required property for all embeddings of  $K$  into all local fields.*

---

5. “Manifold”=“smooth variety”.

6. The words “analytic manifold” are understood in the most naive sense (not rigid-analytic or Berkovich-analytic).

Note that Theorem 3.3 implies Theorem 2.1. This is a consequence of Corollary 3.2 and the following key property of the wave front of a generalized function  $\nu$  on a manifold  $Y$ : the restriction of  $\nu$  to an open subset  $U \subset Y$  is smooth if and only if the intersection of the wave front with  $T^*U$  is contained in the zero section.

The next subsection can be skipped by the reader.

**3.3. Definition of wave front set.** — The definition that we are using slightly differs from the classical definition of L. Hörmander [Hör].

Let  $V$  be a finite-dimensional  $F$ -vector space and let  $\nu$  be a distribution on an open subset  $U \subset V$ . The *wave front* of  $\nu$  is a certain closed subset in  $T^*U = U \times V^*$ . Namely, a point  $(x_0, w_0) \in U \times V^*$  is *not* in the wave front of  $\nu$  if there exists a smooth compactly supported function  $\rho$  on  $V$  with  $\rho(x_0) \neq 0$  such that the Fourier transform<sup>(7)</sup> of  $\rho \cdot \nu$  “vanishes asymptotically” in the direction of  $w_0$ .

The precise meaning of these words is as follows: we say that a function  $f$  on  $V^*$  *vanishes asymptotically in the direction of*  $w_0$  if there exists a smooth compactly supported function  $\sigma$  on  $V^*$  with  $\sigma(w_0) \neq 0$  such that the function  $\phi$  on  $V^* \times F$  defined by  $\phi(w, \lambda) := f(\lambda w) \cdot \sigma(w)$  is a Schwartz function<sup>(8)</sup>.

In other words, if  $w_0 \neq 0$  and  $F$  is non-Archimedean we require the Fourier transform of  $\rho \cdot \nu$  to become compactly supported after restricting to a small open  $F^\times$ -stable neighborhood of  $w_0$ . In the Archimedean case compact support is replaced by rapid decay. If  $w_0 = 0$  we require that  $\rho \cdot \nu = 0$ .

According to [Hör] and [Hef], the above notion of wave front set is invariant with respect to changes of variables, so it makes sense for distributions (or generalized functions) on manifolds.

For more details, see [AD, Appendix A], [Hör], and [Hef].

**3.4. Warning.** — It is well known that the characteristic variety of a coherent  $\mathcal{D}$ -module is always coisotropic (in particular, if it is isotropic then it is Lagrangian). On the other hand, a similar statement for wave front sets is *false*, see Lemma A.3 and Example A.4 from Appendix A.

#### 4. Construction of $U \subset W^*$ and $I \subset T^*W^*$

In this section we construct the sets  $U$  and  $I$  whose existence is claimed in Theorems 2.1 and 3.3.

Let  $\overline{W}$  denote the space of lines in  $W \oplus F$ ; in other words,  $\overline{W}$  is the projective space containing  $W$  as an open subspace. Set  $W_\infty := \overline{W} \setminus W$ ; this is the hyperplane at infinity.

7. The Fourier transform is a smooth function on  $V^*$ .

8. Thus we understand the above word “direction” as a line. Hörmander considers  $F = \mathbb{R}$  and understands “direction” as a half-line (*i.e.*, a ray).

By assumption, the differential form  $\omega$  has no poles on  $X$ , but it may have zeros. Set  $X^\circ := \{x \in X \mid \omega(x) \neq 0\}$ . In what follows we assume that  $X^\circ$  is dense in  $X$  (so  $X \setminus X^\circ$  is a divisor).

By Hironaka's theorem, after replacing the variety  $X$  by a certain modification<sup>(9)</sup> of it,  $\varphi : X \rightarrow W$  admits a compactification to  $\bar{\varphi} : \bar{X} \rightarrow \bar{W}$  with  $\bar{X}$  smooth and  $\bar{X} \setminus X^\circ$  being a NC divisor (or even an SNC divisor). As usual, "NC" (resp. "SNC") stands for "normal crossings" (resp. "strict normal crossings"); the word "strict" means here that each irreducible component of the divisor is smooth.

So one has a commutative diagram

$$(4.1) \quad \begin{array}{ccccc} X & \hookrightarrow & \bar{X} & \supset & X_\infty := \bar{\varphi}^{-1}(W_\infty) \\ \varphi \downarrow & & \bar{\varphi} \downarrow & & \downarrow \\ W & \hookrightarrow & \bar{W} & \supset & W_\infty \end{array}$$

Set  $D := \bar{X} \setminus X^\circ$ . By assumption,  $D$  is a NC divisor. Note that  $X_\infty$  is a divisor contained in  $\bar{X} \setminus X^\circ$ , so  $X_\infty$  is also NC.

We choose the modification of the original  $X$  and its compactification to be defined over the small field  $K$ .

The open subset  $U \subset W^*$  from Theorem 2.1 and the isotropic subvariety  $I$  from Theorem 3.3 are constructed very explicitly in terms of the above choices, and it will be clear that  $U$  and  $I$  are defined over  $K$ . To construct  $U$  and  $I$ , we need some notation.

Let  $\nu : \hat{D} \rightarrow D$  be the normalization. For each  $r \geq 1$  let  $Z'_r$  be the normalization of

$$\{x \in D \mid \text{Card } \nu^{-1}(x) \geq r\}.$$

In particular,  $Z'_1 = \hat{D}$  and  $Z'_r = \emptyset$  if  $r > \dim X$ . It is easy to see that  $Z'_r$  is smooth. If  $D$  is an SNC divisor with irreducible components  $D_j$ ,  $j \in J$ , then  $Z'_r$  is just the disjoint union of the intersections  $D_S := \bigcap_{j \in S} D_j$  corresponding to all subsets  $S \subset J$  of order  $r$ .

Let  $Z_r \subset Z'_r$  denote the disjoint union of those connected components of  $Z_r$  whose image in  $\bar{X}$  is contained in  $X_\infty$ . The map  $\bar{\varphi} : \bar{X} \rightarrow \bar{W}$  induces a map  $\varphi_r : Z_r \rightarrow W_\infty$ .

*Definition of  $U$ .* — Recall that the projective space  $W_\infty$  is the space of 1-dimensional subspaces in  $W$ . The dual projective space  $W_\infty^*$  equals  $(W^* \setminus \{0\})/\mathbb{G}_m$ ; on the other hand, points of  $W_\infty^*$  can be considered as projective hyperplanes  $H \subset W_\infty$ . Let  $\tilde{U} \subset W_\infty^*$  denote the set of those projective hyperplanes  $H \subset W_\infty$  that are transversal<sup>(10)</sup> to  $\varphi_r : Z_r \rightarrow W_\infty$  for each  $r$ . Finally, define  $U \subset W^* \setminus \{0\}$  to be the preimage of  $\tilde{U}$ .

9. A modification of  $X$  is a variety equipped with a proper morphism to  $X$  which is a birational isomorphism.

10. Transversality means that  $Z_r \times_{W_\infty} H$  is a smooth divisor in  $Z_r$ .



To describe the isotropic subset  $I$ , let us recall a general construction: to a morphism  $f : X \rightarrow Y$  between smooth algebraic varieties one associates a conic subset  $\text{Crit}_f \subset T^*Y$ , namely

$$(4.2) \quad \text{Crit}_f = \{(y, \xi) \in T^*Y \mid \exists x \in f^{-1}(y) \text{ such that } (d_x f)^*(\xi) = 0\}.$$

Here we understand the quantifier  $\exists x$  in the sense of algebraic geometry (i.e., the field of definition of  $x$  may be larger than the ground field).

It is well known that  $\text{Crit}_f$  is isotropic<sup>(11)</sup> (e.g., this follows from [CG, Prop. 2.7.51] or [G, Lemma 1]). If  $f$  is proper then  $\text{Crit}_f$  is closed.

*Definition of  $I$ .* — Applying the above construction to  $\varphi_r : Z_r \rightarrow W_\infty$  we get a conic isotropic subvariety  $\text{Crit}_{\varphi_r} \subset T^*W_\infty$ . Set  $I' := \bigcup_r \text{Crit}_{\varphi_r} \subset T^*W_\infty$ . Note that  $W_\infty = (W \setminus \{0\})/\mathbb{G}_m$ , so  $T^*W_\infty$  is a subquotient of  $T^*(W \setminus \{0\}) = (W \setminus \{0\}) \times W^*$ . So  $I'$  defines a subset  $\tilde{I} \subset (W \setminus \{0\}) \times W^*$ . Now set  $I = \tilde{I} \cup (W \times \{0\}) \cup (\{0\} \times W^*)$ .

**Theorem 4.1.** — *The sets  $U$  and  $I$  defined above have the properties required in Theorems 2.1 and 3.3: namely, the generalized function  $\hat{\mu}$  has smooth restriction to  $U$  and the wave front of  $\hat{\mu}$  is contained in  $I$ .*

The proof will be sketched in the next section. A complete proof (under the assumption that  $\overline{X} \setminus X^\circ$  is a divisor with strict normal crossings) is given in [AD].

Note that the claim about  $U$  from Theorem 4.1 follows from the claim about  $I$ . This is clear from the next remark.

**Remark 4.2.** — The subset  $I' \subset T^*W_\infty$  from the definition of  $I$  and the subset  $\tilde{U} \subset W_\infty^*$  from the definition of  $U$  are related as follows. Identifying  $(T^*W_\infty - 0)/\mathbb{G}_m$  with the incidence relation  $\text{Inc} \subset W_\infty \times W_\infty^*$  we get from  $I'$  a subset of  $\text{Inc}$ . Its image in  $W_\infty^*$  equals  $W_\infty^* \setminus \tilde{U}$ .

## 5. Sketch of the proof of Theorem 4.1

**5.1. Strategy.** — We are studying  $\hat{\mu}$ , where  $\mu := \varphi_*|\omega|$ . Let  $\overline{X}$  be as in §4. Decompose  $\varphi : X \rightarrow W$  as

$$(5.1) \quad \begin{array}{ccc} X & \xrightarrow{i} & \overline{X} \times W \\ & \searrow \varphi & \downarrow \pi \\ & & W \end{array}$$

Note that since  $\varphi$  is proper the map  $i$  is a closed embedding. Thus all three maps in diagram (5.1) are proper.

We have  $\mu := \varphi_*|\omega| = \pi_* i_* |\omega|$ . So  $\hat{\mu} = \pi_* \widehat{i_* |\omega|}$ . Here  $i_* |\omega|$  lives on  $\overline{X} \times W$ , and  $\widehat{i_* |\omega|}$  is the partial Fourier transform of  $i_* |\omega|$  with respect to  $W$ .

11. On the other hand,  $\text{Crit}_f$  is not necessarily Lagrangian, see Example A.1 from Appendix A.

The strategy is to first find an upper bound for the wave front of  $\widehat{i_*|\omega|}$  and then to get from it an upper bound for the wave front of  $\hat{\mu}$ . The latter is straightforward<sup>(12)</sup>. In §§5.2–5.4 below we discuss the problem of finding an upper bound for the wave front of  $\widehat{i_*|\omega|}$ ; this problem is, of course, local with respect to  $\overline{X}$ .

**5.2. A calculus-style formula for  $\widehat{i_*|\omega|}$ .** — Let us get rid of the notation  $i_*$  (which is not a part of a standard calculus course).

Let  $x_0 \in \overline{X}$ . Our  $\varphi : X \rightarrow W$  extends to  $\overline{\varphi} : \overline{X} \rightarrow \overline{W}$ , and on some open subset  $\mathcal{U} \subset \overline{X}$  containing  $x_0$  one can write  $\overline{\varphi} : \mathcal{U} \rightarrow \overline{W}$  as

$$\overline{\varphi}(x) = (f(x) : p(x)),$$

where  $f$  is a  $W$ -valued regular function,  $p$  is a scalar regular function, and the two functions have no common zeros. We claim that on  $\mathcal{U} \times W^*$  one has

$$(5.2) \quad \widehat{i_*|\omega|} = \psi \left( \frac{\langle f(x), \xi \rangle}{p(x)} \right) \cdot |\omega|, \quad (x, \xi) \in \mathcal{U} \times W^*.$$

Here and in similar situations below we allow a slight abuse of notation: we write simply  $\omega$  to denote the pullback of  $\omega$  from  $\overline{X}$  to  $\mathcal{U} \times W^*$ , and we tacitly assume restriction from  $\overline{X} \times W^*$  to  $\mathcal{U} \times W^*$  in the l.h.s. of (5.2).

Let us explain the precise meaning of formula (5.2). To simplify the discussion, let us fix a Haar measure<sup>(13)</sup> on  $W^*$ , then the l.h.s. of (5.2) is a distribution on  $\mathcal{U} \times W^*$ .

The r.h.s. of (5.2) clearly makes sense as a distribution on a smaller set  $(\mathcal{U} \cap X) \times W^*$  (since  $p$  has no zeros on  $\mathcal{U} \cap X$  we just have a smooth function times  $|\omega|$ ). But as explained below, the r.h.s. of (5.2) makes sense as a distribution on the whole set  $\mathcal{U} \times W^*$ , and formula (5.2) is, in fact, an equality of distributions on  $\mathcal{U} \times W^*$ .

Here are two equivalent ways to define the r.h.s. of (5.2) as a distribution on  $\mathcal{U} \times W^*$ . First way: for a smooth compactly supported test function  $h$  on  $\mathcal{U} \times W^*$ , interpret

$$\int_{x, \xi} h(x, \xi) \cdot \psi \left( \frac{\langle f(x), \xi \rangle}{p(x)} \right) \cdot |\omega|$$

as  $\iint_{x, \xi}$  (i.e., integrate along  $\xi$  first<sup>(14)</sup>).

To explain the second way, let us assume, for simplicity, that  $\mathcal{U}$  is so small that there is a regular top form  $\omega_0$  on  $\mathcal{U}$  without zeros. The problem is then to define the expression

$$(5.3) \quad \psi \left( \frac{\langle f(x), \xi \rangle}{p(x)} \right) \cdot \left| \frac{\omega}{\omega_0} \right|$$

12. The calculus of wave fronts was developed by Hörmander precisely to make such computations straightforward.

13. If you do not fix a Haar measure on  $W^*$  then the l.h.s. of (5.2) is a distribution along  $\mathcal{U}$  and a generalized function along  $W^*$ .

14. After integrating along  $\xi$  one gets a smooth compactly supported function on  $\mathcal{U}$ . Moreover, this function vanishes on the locus  $p(x) = 0$ . In the Archimedean case all its derivatives vanish there as well.

as a generalized function on  $\mathcal{U} \times W^*$ . In fact, one defines (5.3) as a map

$$(5.4) \quad \mathcal{U} \longrightarrow \{\text{generalized functions on } W^*\}.$$

Namely, if  $p(x) \neq 0$  then (5.3) is understood literally, and if  $p(x) = 0$  then the corresponding generalized function on  $W^*$  is defined to be zero. One checks that the above map (5.4) is *continuous*, so it defines a generalized function on  $\mathcal{U} \times W^*$ .

**5.3. Using formula (5.2).** — The formula reduces the problem to finding an upper bound for the wave front of the distribution

$$(5.5) \quad \psi \left( \frac{\langle f(x), \xi \rangle}{p(x)} \right) \cdot |\omega|, \quad (x, \xi) \in \mathcal{U} \times W^*.$$

Recall that  $f$  and  $p$  have no common zeros. Note that the wave front of (5.5) does not change after replacing  $f(x)$  by  $f(x) + p(x)w_0$ , where  $w_0 \in W$  is fixed. These facts imply that without loss of generality *we can assume that  $f$  has no zeros on  $\mathcal{U}$ .*

Now consider the expression

$$(5.6) \quad \psi \left( \frac{\eta}{p(x)} \right) \cdot |\omega|, \quad (x, \eta) \in \mathcal{U} \times F,$$

which is simpler than (5.5). Define (5.6) as a distribution on the whole  $\mathcal{U} \times F$  (rather than on the open subset where  $p(x) \neq 0$ ) using the procedure from §5.2. The distribution (5.5) is the pullback of (5.6) with respect to the map  $\mathcal{U} \times W^* \rightarrow \mathcal{U} \times F$  defined by  $\eta = \langle f(x), \xi \rangle$  (the pullback is well-defined because  $f$  has no zeros on  $\mathcal{U}$ , which implies that the map  $\mathcal{U} \times W^* \rightarrow \mathcal{U} \times F$  is a submersion). Thus it remains to find an upper bound for the wave front of (5.6).

**5.4. Using toric symmetry.** — The good news is that  $f$  does not appear in formula (5.6). Because of this, the toric symmetry due to the normal crossings assumption becomes manifest. Let us explain more details.

*5.4.1. The SNC case.* — Let us first assume that the divisor  $\overline{X} \setminus X^\circ$  has *strict* normal crossings (this case is enough to prove Theorem 3.3). In this case we can pretend that  $\mathcal{U} = \mathbb{A}^n$  and that  $p$  and  $\omega$  from formula (5.6) are given by *monomials*. Then the problem is to give an upper bound for the wave front of the generalized function <sup>(15)</sup>

$$(5.7) \quad u(x, \eta) = \psi \left( \frac{\eta}{x^\alpha} \right) \cdot |x^\beta|, \quad \text{where } x = (x_1, \dots, x_n) \in F^n, \eta \in F.$$

As usual,  $\alpha$  and  $\beta$  are multi-indices and  $x^\alpha := x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}$ . We assume that  $\alpha_i \geq 0$  and whenever  $\alpha_i = 0$  we have  $\beta_i \geq 0$ ; the latter assumption ensures that (5.7) is well-defined as a generalized function.

---

15. Again, to define (5.7) as a generalized function on the whole space  $F^n \times F$  (rather than outside of the coordinate hyperplanes) we use the procedure from §5.2.

**Key Lemma.** — The wave front of (5.7) is contained in the union of the zero section of the cotangent bundle of  $F^{n+1}$  and the conormal bundles of the coordinate planes of dimensions  $0, 1, \dots, n$ .

*Proof.* — Let  $\mathcal{W}$  denote the wave front of  $u$ . The generalized function  $u$  is quasi-invariant with respect to the action of the torus  $T := (F^\times)^n$  on  $F^{n+1}$  defined by

$$\tilde{x}_i := \lambda_i \cdot x_i, \quad \tilde{\eta} := \lambda^\alpha \cdot \eta, \quad (\lambda_1, \dots, \lambda_n) \in (F^\times)^n.$$

The quasi-invariance property of  $u$  implies the following property of  $\mathcal{W}$ : suppose that  $(z, \alpha) \in \mathcal{W}$ , where  $z \in \mathbb{A}^{n+1}$  and  $\alpha \in T_z^*(\mathbb{A}^{n+1})$ ; then  $\alpha$  vanishes on the tangent space to the  $T$ -orbit containing  $z$ . Combining this property of  $\mathcal{W}$  with the fact that  $u$  is smooth outside of the union of the coordinate hyperplanes, we get the desired statement.  $\square$

**5.4.2. The general case.** — Now let us drop the strictness assumption on the divisor  $\overline{X} \setminus X^\circ$ . Then instead of (5.7) one has to consider the following generalized function on

$$E_1 \times \dots \times E_n \times F,$$

where  $E_1, \dots, E_n$  are finite extensions of  $F$ :

$$u(x_1, \dots, x_n, \eta) = \psi \left( \frac{\eta}{N_{E_1/F}(x_1)^{\alpha_1} \dots N_{E_n/F}(x_n)^{\alpha_n}} \right) \cdot \prod_i |N_{E_i/F}(x_i)|^{\beta_i},$$

$x_i \in E_i$ ,  $\eta \in F$ . An analog of the Key Lemma from §5.4.1 for this generalized function still holds. The proof from §5.4.1 remains valid with an obvious change: instead of the torus  $(F^\times)^n$  one uses the torus  $E_1^\times \times \dots \times E_n^\times$ .

## Appendix A

### How non-Lagrangian isotropic varieties appear

**A.1. The isotropic subset  $\text{Crit}_f$  is not necessarily Lagrangian.** — To a morphism  $f : X \rightarrow Y$  between smooth algebraic varieties one associates an isotropic conic subset  $\text{Crit}_f \subset T^*Y$ , see formula (4.2). The subset  $\text{Crit}_f \subset T^*Y$  is *not necessarily Lagrangian*, i.e., its closure may have components of dimension less than  $\dim Y$ . I learned the following example of this phenomenon from M. Kashiwara.

**Example A.1.** — Define  $f : \mathbb{A}^2 \rightarrow \mathbb{A}^2$  by

$$f(t, x) = (t, t^n x), \quad n \geq 2.$$

The differential of  $f$  never vanishes; it is degenerate if and only if  $t = 0$ . So far we have used that  $n \geq 1$ . Now using that  $n \geq 2$  we see that  $\text{Crit}_f$  is the union of the zero section and a line in the cotangent space of  $(0, 0) \in Y$ .

**Question A.2.** — Is there a *proper* morphism  $f : X \rightarrow Y$  between smooth algebraic varieties such that  $\text{Crit}_f$  is not Lagrangian?

### A.2. A distribution whose wave front is isotropic but not Lagrangian

Consider the map  $f : \mathbb{A}^2 \rightarrow \mathbb{A}^2$  from Example A.1, *i.e.*,

$$(A.1) \quad f(t, x) = (t, y), \text{ where } y = t^n x, \quad n \geq 2.$$

Let  $F$  be a local field of characteristic 0 and  $\varphi : F \rightarrow \mathbb{C}$  a smooth compactly supported function. Define a distribution  $u$  on  $F^2$  by

$$(A.2) \quad u := f_*(\varphi(x) \cdot |dt \wedge dx|)$$

(the r.h.s. makes sense because the restriction of  $f : F^2 \rightarrow F^2$  to the support of the distribution  $\varphi(x) \cdot |dt \wedge dx|$  is proper). As usual, the distribution  $u$  can be considered as a generalized function. This function is smooth outside of the point  $y = t = 0$ ; in fact, it is easy to see that

$$u(t, y) = |t^{-n}| \cdot \varphi(y/t^n) \quad \text{if } t \neq 0; \quad u(t, y) = 0 \quad \text{if } t = 0, y \neq 0.$$

**Lemma A.3.** — *If  $\varphi \neq 0$  then the wave front of  $u$  is the union of the following two sets:*

- a) *the part of the zero section that corresponds to the support of the function  $\varphi(y/t^n)$ ;*
- b) *a line in the cotangent space of the point  $y = t = 0$ .*

*Proof.* — The calculus of wave fronts<sup>(16)</sup> tells us that the wave front of  $u$  is contained in the set of  $F$ -points of  $\text{Crit}_f$ . So the description of  $\text{Crit}_f$  given in Example A.1 yields an upper bound for the wave front. On the other hand, the wave front set is a conic subset of the cotangent bundle which is not contained in the zero section (because  $u$  is not smooth at the point  $y = t = 0$ ).  $\square$

**Example A.4.** — Here is a variant of the above construction assuming that  $F \neq \mathbb{C}$ . Choose  $a \in F^\times$ ,  $a \notin (F^\times)^2$ . Instead of the map (A.1), consider the map  $f : F^2 \rightarrow F^2$  defined by

$$(A.3) \quad f(t, x) = \left( t, \frac{x^3}{3} - at^2x \right).$$

Consider the distribution (A.2) on  $F^2$  corresponding to the new map  $f : F^2 \rightarrow F^2$ . Then Lemma A.3 holds for this distribution (to see this, note that since  $a \notin (F^\times)^2$  the only critical point of  $f : F^2 \rightarrow F^2$  is  $x = t = 0$ ). Note that the algebraic variety  $\text{Crit}_f$  corresponding to the map (A.3) is Lagrangian, but one of its irreducible components has very few  $F$ -points.

16. *E.g.*, see [AD, Prop. 2.3.8] and [AD, Def. 2.3.6].

## References

- [AD] A. AIZENBUD & V. DRINFELD – “The wave front set of the Fourier transform of algebraic measures”, arXiv: 1212.3630, 2012.
- [Ber1] J. BERNSTEIN – “Modules over the ring of differential operators; the study of fundamental solutions of equations with constant coefficients”, *Funct. Anal. Appl.* **5** (1972), no. 2, p. 89–101.
- [CG] N. CHRISS & V. GINZBURG – *Representation theory and complex geometry*, Birkhäuser, Boston, MA, 1997.
- [CL] R. CLUCKERS & F. LOESER – “Fonctions constructibles exponentielles, transformation de Fourier motivique et principe de transfert”, *C. R. Math. Acad. Sci. Paris* **341** (2005), no. 12, p. 741–746.
- [G] V. GINZBURG – “The global nilpotent variety is Lagrangian”, *Duke Math. J.* **109** (2001), no. 3, p. 511–519.
- [Hef] D. B. HEIFETZ – “ $p$ -adic oscillatory integrals and wave front sets”, *Pacific J. Math.* **116** (1985), no. 2, p. 285–305.
- [Hör] L. HÖRMANDER – *The analysis of linear partial differential operators I. Distribution theory and Fourier analysis*, Grundlehren Math. Wiss., vol. 256, Springer-Verlag, Berlin, 1990.
- [HK] E. HRUSHOVSKI & D. KAZHDAN – “Integration in valued fields”, in *Algebraic geometry and number theory*, Progr. Math., vol. 253, Birkhäuser, Boston, MA, 2006, p. 261–405.
- [ILO] L. ILLUSIE, Y. LASZLO & F. ORGOGOZO – “Travaux de gabber sur l’uniformisation locale et la cohomologie étale des schémas quasi-excellents”, Séminaire à l’École polytechnique 2006–2008, arXiv: 1207.3648.
- [Za] O. ZARISKI – “Local uniformization on algebraic varieties”, *Ann. of Math. (2)* **41** (1940), p. 852–896.

---

V. DRINFELD, Department of Mathematics, University of Chicago, 5734 S. University Avenue,  
Chicago, IL 60637, USA • E-mail : drinfeld@math.uchicago.edu