

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

INCOMPRESSIBLE GRASSMANNIANS OF ISOTROPIC SUBSPACES

Nikita A. Karpenko

**Tome 144
Fascicule 4**

2016

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 763-773

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 4, tome 144, décembre 2016

Comité de rédaction

Emmanuel BREUILLARD

Yann BUGEAUD

Jean-François DAT

Charles FAVRE

Marc HERZLICH

O'Grady KIERAN

Raphaël KRIKORIAN

Julien MARCHÉ

Emmanuel RUSS

Christophe SABOT

Wilhelm SCHLAG

Pascal HUBERT (dir.)

Diffusion

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 9
France
smf@smf.univ-mrs.fr

Hindustan Book Agency
O-131, The Shopping Mall
Arjun Marg, DLF Phase 1
Gurgaon 122002, Haryana
Inde

AMS
P.O. Box 6248
Providence RI 02940
USA
www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 178 €, hors Europe : 194 € (\$ 291)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2016

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Stéphane SEURET

INCOMPRESSIBLE GRASSMANNIANS OF ISOTROPIC SUBSPACES

BY NIKITA A. KARPENKO

ABSTRACT. — We study 2-incompressible Grassmannians of isotropic subspaces of a quadratic form, of a hermitian form over a quadratic extension of the base field, and of a hermitian form over a quaternion algebra.

RÉSUMÉ (*Grassmanniennes incompressibles d'espaces isotropes*)

Nous étudions des grassmanniennes 2-incompressibles de sous-espaces isotropes d'une forme quadratique ainsi que d'une forme hermitienne au-dessus d'une extension quadratique du corps de base ou au-dessus d'une algèbre de quaternions.

1. Introduction

Let F be a field and let p be a prime number. We refer to [6] and [15] for definitions and general discussion of canonical p -dimension and p -incompressibility. We only recall that canonical p -dimension $\text{cdim}_p X$ of a smooth complete variety X is the least dimension of the image of a self-correspondence $X \rightsquigarrow X$

Texte reçu le 10 février 2015, révisé le 10 février 2016 et accepté le 10 mars 2016.

NIKITA A. KARPENKO, Mathematical & Statistical Sciences, University of Alberta, Edmonton, Alberta, Canada • *E-mail* : karpenko@ualberta.ca

2010 Mathematics Subject Classification. — 20G15, 14C25.

Key words and phrases. — Quadratic forms, algebraic groups, projective homogeneous varieties, orthogonal, symplectic, and unitary Grassmannians, Chow groups and motives, canonical dimension and incompressibility.

This work has been supported by a Discovery Grant from the National Science and Engineering Board of Canada.

of multiplicity prime to p ; X is called p -incompressible, if $\mathrm{cdim}_p X = \dim X$, that is, if every self-correspondence of multiplicity prime to p is dominant.

We work with *projective homogeneous varieties*, i.e., twisted flag varieties under semisimple affine algebraic groups. One large class of such varieties, for which the p -incompressibility property is completely understood, is given by the generalized Severi-Brauer varieties of central simple algebras, [8]. In the present paper we study another large class which may be considered as the simplest one for which we do not possess a complete general criterion of p -incompressibility. Namely, we study Grassmannians of totally isotropic spaces of a fixed dimension for:

- a non-degenerate quadratic form (orthogonal case, algebraic groups of types $\mathcal{B}$ and $\mathcal{D}$), or
- a hermitian form over a separable quadratic extension field of F (unitary case, type ${}^2\mathcal{A}$), or
- a hermitian form over a quaternion division F -algebra (symplectic case, type $\mathcal{C}$), where in the characteristic 2 case the hermitian form is supposed to be *alternating*, [14, §4.A].

Note that in the symplectic case, only the Grassmannians of subspaces of *integral* dimension over the quaternion algebra are considered because the others are not interesting from the viewpoint of the canonical dimension. Also note that $p = 2$ is the only interesting prime for the varieties treated here.

Our main result on the orthogonal case is Theorem 3.1. Roughly speaking, it asserts that most of the necessary conditions of 2-incompressibility, established in [7], are actually necessary, see Remark 3.2. Therefore Theorem 3.1 may be considered as a step towards finding a precise criterion of 2-incompressibility for this type of varieties.

Our main results for the unitary and the symplectic case are Theorems 4.1 and 5.2. Although the varieties occurring in the three cases have quite different nature, we treat them in similar ways and the obtained results are parallel.

These results can be used to provide a conceptual proof of the criterion of p -incompressibility for products of two projective homogeneous varieties in the case where one of the varieties involved is a Grassmannian of isotropic subspaces, see [12]. Although the criterion has been recently proved in full in [11], the available general proof is very ad-hoc.

Acknowledgements. — I am grateful to Alexander Merkurjev for asking me the question about possibility of generalization of [10, Theorem 3.1] during my talk at the conference “(A)round forms, cycles and motives” on the occasion of the 80th birthday of Albrecht Pfister in Mainz, Germany. I also thank the two anonymous referees (of the submission of the preprint [12] to *Int. Math. Res. Not. IMRN*) for careful reading and suggestions that improved the exposition.

This work has been mostly done during my stay at the Universität Duisburg-Essen and the Max-Planck-Institut für Mathematik in Bonn; I thank them for hospitality.

2. Canonical p -dimension of a fibration

As shown in [10], canonical p -dimension of the product $X \times Y$ of projective homogeneous F -varieties X and Y has $\mathrm{cdim}_p X + \mathrm{cdim}_p Y_{F(X)}$ as an upper bound. We may view X as the base of the projection (a “trivial fibration”) $X \times Y \rightarrow X$, and $Y_{F(X)}$ is its generic fiber. In this section, we generalize this upper bound relation to the case of a more general fibration and also we sharpen the upper bound, replacing $\mathrm{cdim}_p X$ by an, in general, smaller integer $\mathrm{cdim}'_p X$ defined in terms of X and the function field of the total variety of the fibration.

Here is the type of the fibrations we are interested in. Let G be a quasi-split semisimple affine algebraic group over F becoming split over a finite extension field of a p -power degree, T a G -torsor over F , P a parabolic subgroup of G and P' a parabolic subgroup of G contained in P . We consider the fibration

$$\pi: Z := T/P' \rightarrow T/P =: X$$

of projective homogeneous varieties, and we write Y for its generic fiber. We are using the Chow group Ch with coefficients in $\mathbb{F}_p := \mathbb{Z}/p\mathbb{Z}$. In particular, the degree homomorphism $\deg$ on Ch_0 of a complete variety takes its values in $\mathbb{F}_p$.

Let us first recall

PROPOSITION 2.1 ([7, Corollary 6.2]). — *Canonical p -dimension $\mathrm{cdim}_p X$ of X is the minimal integer d such that there exist a cycle class $\alpha \in \mathrm{Ch}^d X_{F(X)}$ and a cycle class $\beta \in \mathrm{Ch}_d X$ with $\deg(\beta_{F(X)} \cdot \alpha) = 1 \in \mathbb{F}_p$.*

LEMMA 2.2. — *In the above settings, we have*

$$\mathrm{cdim}_p Z \leq \mathrm{cdim}'_p X + \mathrm{cdim}_p Y,$$

where $\mathrm{cdim}'_p X$ is the minimal integer d such that there exist elements $\alpha \in \mathrm{Ch}^d(X_{F(Z)})$ and $\beta \in \mathrm{Ch}_d(X)$ with $\deg(\alpha \cdot \beta_{F(Z)}) = 1$.

REMARK 2.3. — Replacing $F(Z)$ by $F(X)$ in the definition of $\mathrm{cdim}'_p X$, we get $\mathrm{cdim}_p(X)$, see Proposition 2.1. Since $F(Z) \supset F(X)$, we have $\mathrm{cdim}'_p X \leq \mathrm{cdim}_p X$.

REMARK 2.4. — In the case where the parabolic subgroup P' is special, Lemma 2.2 has been proved in [5, Lemma 5.3]. The proof was more complicated (and the statement—more specific) because Proposition 2.1 was not available at the time.

Proof of Lemma 2.2. — By the definition of $\mathrm{cdim}'_p X$, for $x := \mathrm{cdim}'_p X$ we can find

$$\alpha_X \in \mathrm{Ch}^x(X_{F(Z)}) \quad \text{and} \quad \beta_X \in \mathrm{Ch}_x(X)$$

with $\deg(\alpha_X \cdot (\beta_X)_{F(Z)}) = 1$.

Note that the function field $F(X)$ is the field of definition of the variety Y . By Proposition 2.1, for $y := \mathrm{cdim}_p Y$ we can find

$$\alpha_Y \in \mathrm{Ch}^y(Y_{F(X)(Y)}) \quad \text{and} \quad \beta_Y \in \mathrm{Ch}_y(Y)$$

with $\deg(\alpha_Y \cdot (\beta_Y)_{F(X)(Y)}) = 1$. We are going to produce certain

$$\alpha'_Y \in \mathrm{Ch}^y(Z_{F(Z)}) \quad \text{and} \quad \beta'_Y \in \mathrm{Ch}_{y+\dim X}(Z)$$

out of α_Y and β_Y .

By [1, Proposition 57.10], the flat pull-back homomorphisms

$$\mathrm{Ch}_{y+\dim X}(Z) \rightarrow \mathrm{Ch}_y(Y) \quad \text{and} \quad \mathrm{Ch}^y(Z_{F(Z)}) \rightarrow \mathrm{Ch}^y(Y \times_{\mathrm{Spec} F} \mathrm{Spec} F(Z))$$

are surjective. We define β'_Y simply as a preimage of β_Y .

In order to define α'_Y , we notice that $F(Z) = F(X)(Y)$ so that $Y \times_{\mathrm{Spec} F} \mathrm{Spec} F(Z) = Y_{F(X)(Y)(X)}$. We define α'_Y as a preimage of $(\alpha_Y)_{F(X)(Y)(X)}$.

Now we set

$$\alpha := \pi^*(\alpha_X) \cdot \alpha'_Y \in \mathrm{Ch}^{x+y}(Z_{F(Z)}) \quad \text{and} \quad \beta := \pi^*(\beta_X) \cdot \beta'_Y \in \mathrm{Ch}_{x+y}(Z).$$

We claim that $\deg(\alpha \cdot \beta_{F(Z)}) = 1$. According to Proposition 2.1, the claim implies that

$$\mathrm{cdim}_p Z \leq x + y = \mathrm{cdim}'_p X + \mathrm{cdim}_p Y.$$

It remains to prove the claim. The claim is about cycles over $F(Z)$ so that we replace the base field F by $F(Z)$.

The product in $\mathrm{Ch}(Z)$, whose degree we are interested in, can be rewritten as the product of $\pi^*(\delta)$, where δ is a 0-cycle class on X of degree 1, by an element $\gamma \in \mathrm{Ch}(Z)$ whose image under $\mathrm{Ch}(Z) \rightarrow \mathrm{Ch}(Y_{F(X)})$ is a 0-cycle class Y of degree 1. We have

$$\deg(\pi^*(\delta) \cdot \gamma) = \deg \pi_*(\pi^*(\delta) \cdot \gamma) = \deg(\delta \cdot \pi_*(\gamma)).$$

To prove the claim it suffices to check that $\pi_*(\gamma) = [X]$. For this, it suffices to check that the pull-back in $\mathrm{Ch}(\mathrm{Spec} F(X)) = \mathbb{F}_p$ of $\pi_*(\gamma)$ along the generic point morphism $\mathrm{Spec} F(X) \rightarrow X$ is equal to 1. By [2, Proposition 1.7 of Chapter 1] applied to the square

$$\begin{array}{ccc} Y & \longrightarrow & \mathrm{Spec} F(X) \\ \downarrow & & \downarrow \\ Z & \longrightarrow & X \end{array}$$

the pull-back of $\pi_*(\gamma)$ coincides with the degree of the image of γ in $\text{Ch}(Y)$ which is indeed 1. The claim and therefore Lemma 2.2 as well are proved. $\square$

3. Incompressible orthogonal Grassmannians

In this section, n is a positive integer, φ is a non-degenerate n -dimensional quadratic form over F , and X_i with $0 \leq i < n/2$ is the i th orthogonal Grassmannian of φ .

We recall that $\dim X_i = i(i-1)/2 + i(n-2i)$. (This formula is valid for $i = n/2$ as well.)

Now assume that $n \geq 3$. The variety X_1 is the projective quadric of φ and the quadratic form $\varphi_{F(X_1)}$ is isotropic. Let φ' be an $(n-2)$ -dimensional quadratic form Witt-equivalent to $\varphi_{F(X_1)}$. For i with $0 \leq i < (n-2)/2$, we write X'_i for the i th orthogonal Grassmannian of φ' . *Higher Witt indexes* appearing below are the *relative higher Witt indexes* of [1, §25].

THEOREM 3.1. — *For some $n \geq 3$ and some $i \geq 1$, assume that the variety X_i is 2-incompressible. Then φ is anisotropic, all the first i higher Witt indexes $i_1(\varphi), \dots, i_i(\varphi)$ of φ are equal to 1, and the variety X'_{i-1} is also 2-incompressible.*

REMARK 3.2. — In [7], the following sufficient condition for 2-incompressibility of X_i has been given: $i_i(\varphi) = 1$ and the degree of every closed point on X_i is divisible by 2^i . The degree condition implies that φ is anisotropic and $i_1(\varphi) = \dots = i_{i-1}(\varphi) = 1$. So, Theorem 3.1 shows that a part of the sufficient condition for 2-incompressibility of X_i is actually necessary.

REMARK 3.3. — For $i = 1$, Theorem 3.1 says that the 2-incompressibility of the projective quadric X_1 implies that φ is anisotropic and its first Witt index is 1 (the variety X'_{i-1} is just a point and as such is 2-incompressible automatically). This statement is easy to check (as in the proof below), Lemma 2.2 is not used. A known old result due to A. Vishik actually says that X_1 is 2-incompressible if and only if φ is anisotropic of first Witt index 1. For a proof (based on ideas of [17]) see [1, Theorem 90.2].

REMARK 3.4. — For odd n , the statement of Theorem 3.1 on maximal $i = (n-1)/2$ can be easily deduced from the properties of the J -invariant of φ . The original paper introducing J -invariant and establishing its main properties is [18]. The monograph [1, §88], where the J -invariant is replaced by its “opposite”, can also be consulted. In particular, see [1, Theorem 90.3] for the relation between canonical 2-dimension and the J -invariant.

Here is the deduction. For $i = (n - 1)/2$, let $X := X_i$ and let

$$e_1 \in \mathrm{Ch}^1(X_{F(X)}), \quad \dots, \quad e_i \in \mathrm{Ch}^i(X_{F(X)})$$

be the ring generators of $\mathrm{Ch}(X_{F(X)})$ defined as in [1, §86]. (The generators e_i are originally introduced in [18], see also [20], and called *elementary classes* there.)

We call an element of $\mathrm{Ch}(X_{F(X)})$ *rational*, if it lies in the image of the change of field homomorphism $\mathrm{Ch}(X) \rightarrow \mathrm{Ch}(X_{F(X)})$. Assuming that X is 2-incompressible, none of $e_1, \dots, e_i$ is rational by [1, Theorem 90.3]. It follows by [1, Proposition 88.8] that the quadratic form φ in question is anisotropic and has $i_1(\varphi) = \dots = i_{i-1}(\varphi) = 1$. Therefore $i_i(\varphi) = 1$ as well. Finally, replacing the base field F by the function field $F(X_1)$ of the quadric X_1 , the element e_i becomes rational, but none of $e_1, \dots, e_{i-1}$, see [1, Corollary 88.6].⁽¹⁾ This shows that

$$\mathrm{cdim}_2 X = 1 + \dots + (i - 1) = \dim X'_{i-1}$$

for the now *equivalent* (in the sense of existence of rational maps in both directions) varieties $X = X_i$ and X'_{i-1} , meaning that X'_{i-1} is 2-incompressible.

Proof of Theorem 3.1. — If φ is isotropic, the variety X_i is equivalent (in the above sense of existence of rational maps in both directions) to the variety X'_{i-1} . Therefore $\mathrm{cdim}_2 X_i = \mathrm{cdim}_2 X'_{i-1}$ (see, e.g., [3, Lemma 3.3]). Since $\dim X_i > \dim X'_{i-1}$, we get a contradiction with the 2-incompressibility of X_i .

We have shown that φ is anisotropic. Next, assuming that $i \geq 2$, we are going to check the 2-incompressibility of X'_{i-1} . For this, we apply Lemma 2.2 to the fibration $\pi : X_{1 \subset i} \rightarrow X_1$, where $X_{1 \subset i}$ is the 2-flag variety projecting onto X_1 and X_i . Note that the variety X'_{i-1} is the generic fiber of π . The $(i - 1)$ th power $h^{i-1} \in \mathrm{Ch}^{i-1}(X_1)$ of the hyperplane section class $h \in \mathrm{Ch}^1(X_1)$ and the class $l_{i-1} \in \mathrm{Ch}_{i-1}(X_1)_{F(X_{1 \subset i})}$ of an i -dimensional totally isotropic subspace satisfy the relation

$$\deg(l_{i-1} \cdot h_{F(X_{1 \subset i})}^{i-1}) = 1.$$

Therefore $\mathrm{cdim}'_2 X_1 \leq \dim X_1 - (i - 1) = (n - 2) - (i - 1) = n - i - 1$. It follows by Lemma 2.2 that

$$\mathrm{cdim}_2 X_{1 \subset i} \leq (n - i - 1) + \mathrm{cdim}_2 X'_{i-1}.$$

Since the flag variety $X_{1 \subset i}$ is equivalent to X_i , we may replace $\mathrm{cdim}_2 X_{1 \subset i}$ by $\mathrm{cdim}_2 X_i$ in this inequality. Since $\dim X_i = (n - i - 1) + \dim X'_{i-1}$, the upper bound on $\mathrm{cdim}_2 X_i$, we obtained, shows that the 2-incompressibility of X_i implies that of X'_{i-1} .

⁽¹⁾ In characteristic $\neq 2$, the statement also follows from A. Vishik's Main Tool Lemma [19, Theorem 3.1], see also [9, Theorem 1.1].

We have shown that X'_{i-1} is 2-incompressible. It follows by the preceding part that the quadratic form φ' is anisotropic. This means that $i_1(\varphi) = 1$. Continuing this induction procedure, we get that $i_1(\varphi) = \cdots = i_{i-1}(\varphi) = 1$.

To finish the proof of Theorem 3.1, it remains to show that $i_i(\varphi) = 1$. Assume that $i_i(\varphi) \geq 2$. In particular, $i < [n/2]$. Let ψ be a $(n-1)$ -dimensional non-degenerate subform of φ . (Note that the notion of non-degeneracy for quadratic forms in characteristic 2 we are using is that of [1] so that non-degenerate quadratic forms exist in any dimension.) Let Y_i be the i th orthogonal Grassmannian of ψ . The condition $i_i(\varphi) \geq 2$ ensures that the varieties X_i and Y_i are equivalent. Indeed, we always have a rational map $Y_i \dashrightarrow X_i$; on the other hand, the intersection of an $(i+1)$ -dimensional totally isotropic subspace of $\varphi_{F(X_i)}$ with the hyperplane of definition of $\psi_{F(X_i)}$ has dimension at least i , showing that $Y_i(F(X_i)) \neq \emptyset$, so that a rational map $X_i \dashrightarrow Y_i$ exists as well. But $\dim Y_i < \dim X_i$ contradicting 2-incompressibility of X_i . $\square$

4. Unitary Grassmannians

Let K/F be a separable quadratic field extension, $n \geq 0$ an integer, V an n -dimensional vector space over K , and φ a K/F -hermitian form on V . For any integer i with $0 \leq i \leq n/2$, we write X_i for the unitary Grassmannian of totally isotropic i -dimensional K -subspaces in V .

We recall that $\dim X_i = i(2n-3i)$.

Assume that $n \geq 2$. Then the variety X_1 is defined and the hermitian form $\varphi_{F(X_1)}$ is isotropic. Let φ' be an $(n-2)$ -dimensional $K(X_1)/F(X_1)$ -hermitian form Witt-equivalent to φ . For i with $0 \leq i \leq (n-2)/2$, we write X'_i for the corresponding unitary Grassmannian of totally isotropic i -dimensional subspaces.

THEOREM 4.1. — *For some integers n and i with $1 \leq i \leq n/2$, assume that the variety X_i is 2-incompressible. Then φ is anisotropic, all the first i higher Witt indexes $i_1(\varphi), \dots, i_i(\varphi)$ of φ are equal to 1, and the variety X'_{i-1} is also 2-incompressible.*

Proof. — We start by repeating word by word the proof of Theorem 3.1. The first change that occurs in the unitary case compared to the orthogonal one is in the formula for $\dim X_i$. But we still have $\dim X_i > \dim X'_{i-1}$ in the setting, so that anisotropy of φ is proved.

As the next step, we apply Lemma 2.2 to the projection $\pi : X_{1 \subset i} \rightarrow X_1$ in order to prove the 2-incompressibility of X'_{i-1} , which is again the generic fiber of π . The base X_1 of the projection, which was a usual quadric in the orthogonal case, is now a sort of *unitary quadric*, and we need information on its Chow group analogous to the information we have and have used for usual quadrics.

Such an information is provided in [4, §3]. It is shown there that there are elements $h \in \text{Ch}^2(X_1)$ and $l_{i-1} \in \text{Ch}_{2(i-1)}(X_1)_{F(X_i)}$ with $\deg(l_{i-1} \cdot h^{i-1}) = 1$. (The notation for the elements we use is similar to the quadric case and differs from [4].) Therefore

$$\text{cdim}'_2 X_1 \leq \dim X_1 - 2(i-1) = (2n-3) - 2(i-1) = 2n - 2i - 1.$$

It follows by Lemma 2.2 that

$$\text{cdim}_2 X_{1 \subset i} \leq (2n - 2i - 1) + \text{cdim}_2 X'_{i-1}.$$

Since the flag variety $X_{1 \subset i}$ is equivalent to X_i , we may replace $\text{cdim}_2 X_{1 \subset i}$ by $\text{cdim}_2 X_i$ in this inequality. Since $\dim X_i = (2n - 2i - 1) + \dim X'_{i-1}$, the upper bound on $\text{cdim}_2 X_i$, we obtained, shows that 2-incompressibility of X_i implies that of X'_{i-1} .

We have shown that X'_{i-1} is 2-incompressible. It follows by the preceding part that the quadratic form φ' is anisotropic. This means that $i_1(\varphi) = 1$. Continuing this induction procedure, we get that $i_1(\varphi) = \dots = i_{i-1}(\varphi) = 1$.

To finish the proof of Theorem 4.1, it remains to show that $i_i(\varphi) = 1$. Assume that $i_i(\varphi) \geq 2$. In particular, $i < [n/2]$. Let ψ be a $(n-1)$ -dimensional non-degenerate subform of φ . The form ψ exists because φ can be diagonalized, [16, Theorem 6.3 of Chapter 7]. Let Y_i be the i th unitary Grassmannian of ψ . By precisely the same argument as in the orthogonal case, the condition $i_i(\varphi) \geq 2$ ensures that the varieties X_i and Y_i are equivalent. But again $\dim Y_i < \dim X_i$, contradicting 2-incompressibility of X_i . $\square$

5. Symplectic Grassmannians

Let Q be a quaternion division F -algebra, $n \geq 0$ an integer, V a right n -dimensional vector space over Q , φ a hermitian (with respect to the canonical symplectic involution on Q) form on V . In the case of $\text{char } F = 2$, we require φ to be alternating.

For any integer i with $0 \leq i \leq n/2$, we write X_i for the symplectic Grassmannian of totally isotropic i -dimensional Q -subspaces in V . We refer to [13] for a general discussion on these varieties and recall that

$$\dim X_i = i(4n - 6i + 1).$$

Note that besides the symplectic Grassmannians of φ introduced right above, for any odd integer j with $1 \leq j \leq n$ there is the variety of totally isotropic subspaces in V of “dimension” $j/2$ over Q . Any such variety however is equivalent to the conic of Q and therefore is not interesting regarding the questions we consider in the paper.

LEMMA 5.1. — Assume that $n \geq 2$. There exists an element $h \in \text{Ch}^4(X_1)$ and for any integer i with $1 \leq i \leq n/2$ there exists an element $l_{i-1} \in \text{Ch}^{4(i-1)}(X_1)_{F(X_i)}$ such that $\deg(l_{i-1} \cdot h_{F(X_i)}^{i-1}) = 1$.

Proof. — The variety X_1 is a closed hypersurface (a sort of “symplectic quadric”) in the “projective space” $Q\mathbb{P}(V)$ —the variety of 1-dimensional Q -subspaces of V , cf. [13]. Note that $\dim Q\mathbb{P}(V) = 4(n-1)$, where $n = \dim V$. Picking up a hyperplane W in V , we consider the class

$$H := [Q\mathbb{P}(W)] \in \text{Ch}^4(Q\mathbb{P}(V))$$

and define $h \in \text{Ch}^4(X_1)$ as the pull-back of H .

Now we replace the base field F by the function field $F(X_i)$, pick up a totally isotropic i -dimensional Q -subspace in V , and let l_{i-1} be its class in $\text{Ch}_{4(i-1)}(X_1)$. For the closed imbedding $in : X_1 \hookrightarrow Q\mathbb{P}(V)$ we have $in_*(l_{i-1}) = H^{n-i}$. Since $\deg(H^{n-1}) = 1$, we get, applying the projection formula for in :

$$\deg(l_{i-1} \cdot h^{i-1}) = \deg in_*(l_{i-1} \cdot h^{i-1}) = \deg(in_*(l_{i-1}) \cdot H^{i-1}) = \deg(H^{n-1}) = 1. \quad \square$$

As in Lemma 5.1, assume that $n \geq 2$. Then the variety X_1 is defined and the hermitian form $\varphi_{F(X_1)}$ is isotropic. Let φ' be an $(n-2)$ -dimensional hermitian form Witt-equivalent to φ . For i with $0 \leq i \leq (n-2)/2$, we write X'_i for the corresponding Grassmannian of totally isotropic i -dimensional subspaces.

THEOREM 5.2. — For some $n \geq 2$ and some $i \geq 1$, assume that the variety X_i is 2-incompressible. Then φ is anisotropic, all the first i higher Witt indexes $i_1(\varphi), \dots, i_i(\varphi)$ of φ are equal to 1, and the variety X'_{i-1} is also 2-incompressible.

Proof. — Again everything goes through as in the proof of Theorem 3.1 (or also Theorem 4.1), although the symplectic Grassmannians we are working with now satisfy different dimension formulas compared to the varieties we had in the previous sections. Nevertheless we still have $\dim X_i > \dim X'_{i-1}$ in the current setting, so that we obtain the anisotropy of φ .

The next step is, as before, the proof of 2-incompressibility of X'_{i-1} , involving Lemma 2.2 and the projection $\pi : X_{1 \subset i} \rightarrow X_1$. The information on the Chow group of X_1 that we need here now is provided by Lemma 5.1. It shows that

$$\text{cdim}'_2 X_1 \leq \dim X_1 - 4(i-1) = (4n-5) - 4(i-1) = 4n - 4i - 1.$$

It follows by Lemma 2.2 that

$$\text{cdim}_2 X_{1 \subset i} \leq (4n - 4i - 1) + \text{cdim}_2 X'_{i-1}.$$

Since again the flag variety $X_{1\subset i}$ is equivalent to X_i , we may replace $\text{cdim}_2 X_{1\subset i}$ by $\text{cdim}_2 X_i$ in this inequality. Since $\dim X_i = (4n - 4i - 1) + \dim X'_{i-1}$, the upper bound on $\text{cdim}_2 X_i$, we obtained, shows that 2-incompressibility of X_i implies that of X'_{i-1} .

We have shown that X'_{i-1} is 2-incompressible. It follows as in the orthogonal case that $i_1(\varphi) = \dots = i_{i-1}(\varphi) = 1$.

To finish the proof of Theorem 5.2, it remains to show that $i_i(\varphi) = 1$. Assume that $i_i(\varphi) \geq 2$. In particular, $i < [n/2]$. Let ψ be a $(n - 1)$ -dimensional non-degenerate subform of φ . Let Y_i be the i th unitary Grassmannian of ψ . The condition $i_i(\varphi) \geq 2$ ensures that the varieties X_i and Y_i are equivalent. But now again $\dim Y_i < \dim X_i$ contradicting the 2-incompressibility of X_i . $\square$

BIBLIOGRAPHY

- [1] R. ELMAN, N. KARPENKO & A. MERKURJEV – *The algebraic and geometric theory of quadratic forms*, American Mathematical Society Colloquium Publications, vol. 56, Amer. Math. Soc., Providence, RI, 2008.
- [2] W. FULTON – *Intersection theory*, second ed., *Ergebn. Math. Grenzgeb. 3. Folge. A Series of Modern Surveys in Mathematics*, vol. 2, Springer, Berlin, 1998.
- [3] N. KARPENKO & Z. REICHSTEIN – “A numerical invariant for linear representations of finite groups”, *Comment. Math. Helv.* **90** (2015), p. 667–701.
- [4] N. KARPENKO & M. ZHYKHOVICH – “Isotropy of unitary involutions”, *Acta Math.* **211** (2013), p. 227–253.
- [5] N. A. KARPENKO – “A bound for canonical dimension of the (semi)spinor groups”, *Duke Math. J.* **133** (2006), p. 391–404.
- [6] ———, “Canonical dimension”, in *Proceedings of the International Congress of Mathematicians. Volume II*, Hindustan Book Agency, New Delhi, 2010, p. 146–161.
- [7] ———, “Sufficiently generic orthogonal Grassmannians”, *J. Algebra* **372** (2012), p. 365–375.
- [8] ———, “Upper motives of algebraic groups and incompressibility of Severi-Brauer varieties”, *J. reine angew. Math.* **677** (2013), p. 179–198.
- [9] ———, “Variations on a theme of rationality of cycles”, *Cent. Eur. J. Math.* **11** (2013), p. 1056–1067.
- [10] ———, “Incompressibility of products of Weil transfers of generalized Severi-Brauer varieties”, *Math. Z.* **279** (2015), p. 767–777.
- [11] ———, “Incompressibility of products of pseudo-homogeneous varieties”, *Canadian Math. Bull.* (2016), <http://dx.doi.org/10.4153/CMB-2016-024-4>.

- [12] ———, “Incompressibility of products by Grassmannians of isotropic subspaces”, preprint <https://www.math.uni-bielefeld.de/lag/man/541.pdf>.
- [13] N. A. KARPENKO & A. S. MERKURJEV – “Hermitian forms over quaternion algebras”, *Compos. Math.* **150** (2014), p. 2073–2094.
- [14] M.-A. KNUS, A. MERKURJEV, M. ROST & J.-P. TIGNOL – *The book of involutions*, American Mathematical Society Colloquium Publications, vol. 44, Amer. Math. Soc., Providence, RI, 1998.
- [15] A. S. MERKURJEV – “Essential dimension: a survey”, *Transform. Groups* **18** (2013), p. 415–481.
- [16] W. SCHARLAU – *Quadratic and Hermitian forms*, Grundlehren der Math. Wiss., vol. 270, Springer, Berlin, 1985.
- [17] A. VISHIK – “Motives of quadrics with applications to the theory of quadratic forms”, in *Geometric methods in the algebraic theory of quadratic forms*, Lecture Notes in Math., vol. 1835, Springer, Berlin, 2004, p. 25–101.
- [18] ———, “On the Chow groups of quadratic Grassmannians”, *Doc. Math.* **10** (2005), p. 111–130.
- [19] ———, “Generic points of quadrics and Chow groups”, *Manuscripta Math.* **122** (2007), p. 365–374.
- [20] ———, “Fields of u -invariant $2^r + 1$ ”, in *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, Progr. Math., vol. 270, Birkhäuser, 2009, p. 661–685.

