

quatrième série - tome 46 fascicule 5 septembre-octobre 2013

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Chloé PERIN

Erratum to:

Elementary embeddings in torsion-free hyperbolic groups

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ERRATUM TO: ELEMENTARY EMBEDDINGS IN TORSION-FREE HYPERBOLIC GROUPS

BY CHLOÉ PERIN

The notations adopted are those of [1].

Proposition 5.11 of [1] states that if a torsion-free hyperbolic group A admits a cyclic JSJ-like decomposition Λ , and a non injective morphism $f : A \rightarrow A$ which restricts to conjugation on each non surface type vertex group, and sends surface type vertex groups to non abelian images, then there is a retraction $r : A \rightarrow A'$ which gives A a structure of hyperbolic floor over A' .

Unfortunately, we realised that Proposition 5.11 fails to hold in a few exceptional low complexity cases. The natural modification to overcome this mistake is to proceed to a slight generalization of the notion of hyperbolic floors and hyperbolic towers, which we present in Section 1.

As we will see in Section 2, however, this does not affect Theorem 1.2, the main result of the paper. Moreover, Theorem 1.2 is the only result which directly uses Proposition 5.11 in its proof.

For a corrected version of the paper, see [2]. We sincerely apologize for any confusion caused by this mistake.

1. Modification of Proposition 5.11: extended hyperbolic towers

To be made correct, the statement of 5.11 needs to be slightly weakened to

PROPOSITION 1. – *Let A be a torsion-free hyperbolic group. Let Λ be a cyclic JSJ-like decomposition of A which does not consist of a single surface type vertex.*

Assume that there exists a non injective preretraction $A \rightarrow A$ with respect to Λ . Then there exists a subgroup A' of A and a retraction $r : A \rightarrow A'$ such that (A, A', r) is an extended hyperbolic floor. Moreover given a non surface type vertex group R_0 of Λ we can choose A' to contain R_0 .

where extended hyperbolic floors are given by

DEFINITION 2 (extended hyperbolic floor). – Consider a triple (G, G', r) where G is a group, G' is a subgroup of G , and r is a retraction from G onto G' .

We say that (G, G', r) is an extended hyperbolic floor if there exist a non trivial decomposition Γ of G as a graph of groups with surfaces, and a Bass-Serre presentation (T^1, T^0) of Γ such that:

- the surfaces of Γ which are not once punctured tori have Euler characteristic at most -2 ;
- G' is the free product of the stabilizers of the non surface type vertices of T^0 ;
- every edge of Γ joins a surface type vertex to a non surface type vertex (bipartism);
- either the retraction r sends surface type vertex groups of Γ to non abelian images; or G' is cyclic and there exists a retraction $r' : G * \mathbb{Z} \rightarrow G' * \mathbb{Z}$ which sends surface type vertex groups of Γ to non abelian images.

We can then define extended hyperbolic towers from extended hyperbolic floors in exactly the same way that we defined hyperbolic towers from hyperbolic floors in Definition 5.4 of [1].

REMARK 3. – Note that all the extended hyperbolic floors (G^k, G^{k+1}, r_k) of an extended hyperbolic tower must in fact be hyperbolic floors, except possibly for the bottom floor (G^{m-1}, G^m, r_{m-1}) . Moreover if this is the case, then G^m must be cyclic, so H is cyclic or trivial. In particular every extended hyperbolic tower structure over a non trivial non cyclic group is in fact a hyperbolic tower.

Note also that in parallel to Remark 5.5 of [1], the structure of extended hyperbolic tower is stable under free products and is transitive.

2. Proving the main result

We now show Theorem 1.2 can still be proved using Proposition 1 above instead of Proposition 5.11 of [1].

THEOREM 4 (Theorem 1.2 of [1]). – If H is an elementary subgroup of a torsion-free hyperbolic group G , then G admits a structure of hyperbolic tower over H .

Proof. – The case where G is abelian is trivial, so we may assume that G , and thus H , is not abelian. The proof given in [1] proceeds by uncovering successive hyperbolic floors structures using Proposition 5.11, and combining them using stability under free products and transitivity of hyperbolic tower structure. Applying Proposition 1 instead will produce extended hyperbolic floors, which we noted we can also combine. We thus get that G admits a structure of extended hyperbolic tower over H . However since H is not abelian, Remark 3 implies that G is in fact a hyperbolic tower over H . \square

3. Proof of Proposition 1

The bulk of the proof of Proposition 5.11 in [1] consists in building hyperbolic floors from preretractions which satisfy either one of two sets of (weak) extra hypotheses: those of Lemma 7.25, or those of Lemma 7.26. The proof in these cases is not affected. However Proposition 7.28, which claims that the conclusions of Proposition 5.11 still hold in the special case when neither set of hypotheses is satisfied, is erroneous. Note that the corrections to be made thus affect Section 7.8 of [1] only.

Proposition 7.28 should be replaced by the following weaker statement

PROPOSITION 5. – *Let A be a torsion-free hyperbolic group which admits a cyclic JSJ-like decomposition Λ which does not consist of a single surface type vertex. Let $f : A \rightarrow A$ be a non injective preretraction with respect to Λ .*

Suppose that Λ and f satisfy neither the hypotheses of Lemma 7.25 of [1], nor those of Lemma 7.26 of [1]. Then there exists a retraction $r : A \rightarrow A'$ to a proper subgroup of A such that (A, A', r) is an extended hyperbolic floor. If moreover Λ has at least two surface type vertices, (A, A', r) can be assumed to be a hyperbolic floor.

Recall that the hypotheses of Proposition 5 imply in particular that Λ has a single non surface type vertex, whose corresponding group is a cyclic group $Z = \langle z \rangle$. Any boundary subgroup of a surface group of Λ is thus generated by a conjugate of z^k for some k , we call k the boundary exponent corresponding to this boundary component. We first prove the following stronger version of Lemma 7.29.

LEMMA 6. – *Under the hypotheses of Proposition 5, there exists a surface Σ of Λ with boundary exponents k_1, \dots, k_m such that the sum $k_1 + \dots + k_m$ is even. If Σ is in fact orientable, we have $k_1 + \dots + k_m = 0$.*

Moreover, if Σ is (i) a 4-punctured sphere, (ii) a 3-punctured projective plane, (iii) a 2-punctured Klein bottle, we must have (up to reordering) in case (i) $k_1 = -k_2$ and $k_3 = -k_4$; in case (ii) $k_1 = -k_2$ and k_3 even; in case (iii) $k_1 = -k_2$, or k_1 and k_2 both even.

Proof. – Since the hypotheses of Lemma 7.26 of [1] are not satisfied, there exists a surface type vertex group S of Λ (with corresponding surface Σ) all of whose exterior surface type vertex group have abelian images by f . Lemma 7.29 of [1] shows that the first part of the statement holds for this choice of Σ . To prove the moreover part, note that $f(S)$ is not abelian, though $f(\hat{S})$ is for any exterior surface type vertex group \hat{S} : the morphism f must kill an element corresponding to a two-sided simple closed curve on Σ which does not bound a Möbius band. Since Σ is small, there are few possibilities and the result follows easily. \square

We then have the following analogue of Lemma 7.30.

LEMMA 7. – *Suppose Λ admits a surface Σ which satisfies the conclusion of Lemma 7.29. If Σ has only one boundary component, then A admits a structure of extended hyperbolic floor, which can be assumed to be a hyperbolic floor if $\chi(\Sigma) < -2$ or if Λ has surfaces other than Σ .*

Proof. – Note that $k_1 \neq 0$, so Σ is not orientable and k_1 is even, say $k_1 = 2k$. If S is the vertex group corresponding to Σ , the subgroup $S' = \langle z^k, S \rangle_A$ is the fundamental group of the closed non orientable surface Σ' obtained by gluing a Möbius band M along the boundary component of Σ . We have $S' = \langle d_1, \dots, d_p \mid d_1^2 \dots d_p^2 = 1 \rangle$.

If $p > 5$ there is a retraction $S' \rightarrow \langle d_1, d_2 \rangle_{S'}$ given by $r(d_3) = d_2^{-1}, r(d_4) = d_1^{-1}$ and $r(d_j) = 1$ for all $j > 4$. It extends to a retraction which gives $(A, r(A), r)$ a structure of hyperbolic floor (with a single surface which is the once punctured connected sum of $p - 2$ projective planes).

If $p = 4$, let s be the generator of an infinite cyclic group \mathbb{Z} . We define a retraction r on $A * \mathbb{Z}$ by setting $r(d_1) = d_1, r(d_2) = d_1^{-1}$, and $r(d_3) = r(d_4^{-1}) = s$, and extending it to $A * \mathbb{Z}$ by the identity on the other surface type vertex groups of Λ and on \mathbb{Z} . Thus A has a structure of extended hyperbolic floor over a proper subgroup. If Λ has surfaces other than Σ , we can further retract $A * \mathbb{Z}$ onto \mathbb{Z} by sending s to a non boundary element in one of the other surface groups: this gives a structure of hyperbolic floor. □

The proof of Proposition 5 then goes as follows

Proof. – Let Σ be the surface of Λ which satisfies the conclusion of Lemma 7.29. By Lemma 7, we may assume Σ has at least two boundary components.

If Σ is an m -punctured surface of genus at least 1 ($m \geq 2$), with the exception of a twice punctured torus, the retraction r given in the proof of Proposition 7.28 does make $(A, r(A), r)$ into a hyperbolic floor. In the other cases, the surface retracted by r is not complex enough to appear as the surface of a hyperbolic floor, or the image of the retraction is a cyclic subgroup (see Figures 1 and 2): this is where the mistake lies.




FIGURE 1. The proof of Proposition 7.28 proposes to retract the fundamental group of the surface Σ_2 on that of Σ_1 (in the even and odd genus cases respectively)...

We consider now the case of a twice punctured torus. The corresponding vertex group admits a presentation as $\langle x, y, b_1, b_2 \mid [x, y] = b_1 b_2 \rangle$, and we must have $b_1 = z^k$ and $b_2 = tz^{-k}t^{-1}$ for t a Bass-Serre element corresponding to the extra edge joining the surface vertex to the rigid vertex. The retraction which sends x to z^k, y to t , and is the identity on \mathbb{Z} gives a structure of hyperbolic floor for A over the (free) subgroup generated by z and t .

If Σ has genus 0, it is an m -punctured (i) sphere, (ii) projective plane or (iii) Klein bottle. If m is at least (i) 5, (ii) 4, and (iii) 3 respectively, then A admits a structure of hyperbolic floor with one surface group corresponding to the complement of a thrice punctured sphere in Σ .