

*quatrième série - tome 47    fascicule 4    juillet-août 2014*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Bhargav BHATT & Aise Johan DE JONG

*Lefschetz for Local Picard groups*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

## LEFSCHETZ FOR LOCAL PICARD GROUPS

BY BHARGAV BHATT AND AISE JOHAN DE JONG

---

**ABSTRACT.** – We prove a strengthening of the Grothendieck-Lefschetz hyperplane theorem for local Picard groups conjectured by Kollár. Our approach, which relies on acyclicity results for absolute integral closures, also leads to a restriction theorem for higher rank bundles on projective varieties in positive characteristic.

**RÉSUMÉ.** – Nous prouvons un renforcement du théorème de l’hyperplan de Grothendieck-Lefschetz pour les groupes locaux de Picard conjecturés par Kollár. Notre approche, qui s’appuie sur des résultats en fermetures absolues, conduit également à un théorème de restriction pour les faisceaux de rang supérieur sur les variétés projectives en caractéristique positive.

A classical theorem of Lefschetz asserts that non-trivial line bundles on a smooth projective variety of dimension  $\geq 3$  remain non-trivial upon restriction to an ample divisor, and plays a fundamental role in understanding the topology of algebraic varieties. In [6], Grothendieck recast this result in more general terms using the machinery of formal geometry and deformation theory, and also stated a local version. With a view towards moduli of higher dimensional varieties, especially the deformation theory of log canonical singularities, Kollár recently conjectured [15] that Grothendieck’s local formulation remains true under weaker hypotheses than those imposed in [6]. Our goal in this paper is to prove Kollár’s conjecture for rings containing a field.

### Statement of results

Let  $(A, \mathfrak{m})$  be an excellent normal local ring containing a field. Fix some  $0 \neq f \in \mathfrak{m}$ . Let  $V = \text{Spec}(A) - \{\mathfrak{m}\}$ , and  $V_0 = \text{Spec}(A/f) - \{\mathfrak{m}\}$ . The following result is the key theorem in this paper; it solves [15, Problem 1.3] completely, and [15, Problem 1.2] in characteristic 0:

**THEOREM 0.1.** – *Assume  $\dim(A) \geq 4$ . The restriction map  $\text{Pic}(V) \rightarrow \text{Pic}(V_0)$  is:*

1. *injective if  $\text{depth}_{\mathfrak{m}}(A/f) \geq 2$  and  $A$  has characteristic 0;*
2. *injective up to  $p^\infty$ -torsion if  $A$  has characteristic  $p > 0$ .*

This result is sharp: surjectivity fails in general, while injectivity fails in general if  $\dim(A) \leq 3$ , in characteristic 0 if  $\text{depth}_{\mathfrak{m}}(A/f) < 2$ , and in characteristic  $p$  if one includes  $p$ -torsion. Theorem 0.1 leads to a fibral criterion for a Weil divisor to be Cartier in a family, see Theorem 1.30. A stronger analogue of Theorem 0.1, including the mixed characteristic case, is due to Grothendieck [6, Expose XI] under the stronger condition  $\text{depth}_{\mathfrak{m}}(A/f) \geq 3$ ; complex analytic variants of Grothendieck's theorem are proven in [7], while topological analogues are discussed in [9]. Without this depth constraint, a previously known case of Theorem 0.1 was when  $A$  has log canonical singularities in characteristic 0, and  $\{\mathfrak{m}\} \subset \text{Spec}(A)$  is not an lc center (see [15, Theorem 19]).

Our approach to Theorem 0.1 relies on formal geometry over absolute integral closures [2, 11], and applies to higher rank bundles as well as projective varieties. This technique then leads to a short proof of the following result:

**THEOREM 0.2.** – *Let  $X$  be a normal projective variety of dimension  $d \geq 3$  over an algebraically closed field of characteristic  $p > 0$ . If a vector bundle  $E$  on  $X$  is trivial over an ample divisor, then  $(\text{Frob}_X^e)^* E \simeq \mathcal{O}_X^{\oplus r}$  for  $e \gg 0$ .*

The numerical version of Theorem 0.2 for line bundles is due to Kleiman [13, Corollary 2, page 305]. The non-numerical version of the rank 1 case, with stronger assumptions on the singularities, is studied in [8]. This result may also be deduced from the boundedness [16] of semistable sheaves. We do not know the correct characteristic 0 analogue of this result.

### An outline of the proof

Both theorems are similar in spirit, so we only discuss Theorem 0.1 here. We first prove the characteristic  $p$  result, and then deduce the characteristic 0 one by reduction modulo  $p$  and an approximation argument; the reduction necessitates the (unavoidable) depth assumption in characteristic 0. The characteristic  $p$  proof follows Grothendieck's strategy of decoupling the problem into two pieces: one in formal  $f$ -adic geometry, and the other an algebraization question. Our main new idea is to replace (thanks entirely to the Hochster-Huneke vanishing theorem [11]) our ring  $A$  with a very large extension  $\bar{A}$  with better depth properties; Grothendieck's deformation-theoretic approach then immediately solves the formal geometry problem over  $\bar{A}$ . Next, we algebraize the solution over  $\bar{A}$  by algebraically approximating formal sections of line bundles; the key here is to identify the cohomology of the formal completion of a scheme as the *derived* completion of the cohomology of the original scheme, i.e., a weak analogue of the formal functions theorem devoid of the usual finiteness constraints. Finally, we descend from  $\bar{A}$  to  $A$ ; this step is trivial in our context, but witnesses the torsion in the kernel.

### Acknowledgements

We thank János Kollár for many helpful discussions and email exchanges concerning Theorem 0.1, Adrian Langer for sharing with us the alternative proof of Theorem 0.2 after a first version of this paper was posted, and Brian Lehmann for bringing to our attention the question answered in Theorem 2.9.

## 1. Local Picard groups

The goal of this section is to prove Theorem 0.1. In §1.1, we study formal geometry along a divisor on a (punctured) local scheme abstractly, and establish certain criteria for restriction map on Picard groups to be injective. These are applied in §1.2 to prove the characteristic  $p$  part of Theorem 0.1. Using the principle of “reduction modulo  $p$ ” and a standard approximation argument (sketched in §1.4), we prove the characteristic 0 part of Theorem 0.1 in §1.3. The afore-mentioned fibral criterion is recorded in §1.5. Finally, in §1.6, we give examples illustrating the necessity of the assumptions in Theorem 0.1.

### 1.1. Formal geometry over a punctured local scheme

We establish some notation that will be used in this section.

NOTATION 1.1. – Let  $(A, \mathfrak{m})$  be a local ring, and fix a regular element  $f \in \mathfrak{m}$ . Let  $X = \text{Spec}(A)$ ,  $V = \text{Spec}(A) - \{\mathfrak{m}\}$ . For an  $X$ -scheme  $Y$ , write  $Y_n$  for the reduction of  $Y$  modulo  $f^{n+1}$ , and  $\widehat{Y}$  for the formal completion<sup>(1)</sup> of  $Y$  along  $Y_0$ . Let  $\text{Vect}(Y)$  be the category of vector bundles (i.e., finite rank locally free sheaves) on  $Y$ , and write  $\text{Pic}(Y)$  and  $\underline{\text{Pic}}(Y)$  for the set and groupoid of line bundles respectively. Set  $\underline{\text{Pic}}(\widehat{Y}) := \lim \underline{\text{Pic}}(Y_n)$  (where the limit is in the sense of groupoids), and  $\text{Pic}(\widehat{Y}) := \pi_0(\underline{\text{Pic}}(\widehat{Y}))$ . For any  $A$ -module  $M$  with associated quasi-coherent sheaf  $\widetilde{M}$  on  $\text{Spec}(A)$ , we define  $H_{\mathfrak{m}}^i(M)$  as cohomology supported along  $\{\mathfrak{m}\} \subset X$  of  $\widetilde{M}$ , i.e., as the  $i$ th cohomology of the complex  $\text{R}\Gamma_{\mathfrak{m}}(M)$  defined as the homotopy-kernel of the map  $\text{R}\Gamma(\text{Spec}(A), \widetilde{M}) \rightarrow \text{R}\Gamma(V, \widetilde{M})$ .

We will use formal schemes associated to certain non-Noetherian  $X$ -schemes later in this paper. Rather than developing the general theory of such schemes, we simply define the concept that will be most relevant: cohomology.

DEFINITION 1.2. – Fix an  $X$ -scheme  $Y$ . For  $F \in D(\mathcal{O}_Y)$ , set  $\widehat{F} := \text{R}\lim(F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n})$ ; we view  $\widehat{F}$  as an  $\mathcal{O}_{\widehat{Y}} := \lim_n \mathcal{O}_{Y_n}$ -complex on  $|\widehat{Y}| := Y_0$ , so  $\text{R}\Gamma(\widehat{Y}, \widehat{F}) := \text{R}\Gamma(Y_0, \widehat{F}) \simeq \text{R}\lim \text{R}\Gamma(Y_0, F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n})$ .

The following two examples help explain the meaning of this definition:

EXAMPLE 1.3. – If  $F$  is a quasicoherent  $\mathcal{O}_X$ -module associated to an  $A$ -module  $M$ , then  $\text{R}\Gamma(\widehat{X}, \widehat{F}) \simeq \text{R}\lim(M \otimes_A^L A/(f^n))$ . In particular, if  $M$  is  $A$ -flat, then  $\text{R}\Gamma(\widehat{X}, \widehat{F})$  is the  $f$ -adic completion of  $M$  in the usual sense. Note that if  $M$  is not  $A$ -flat, then  $\text{R}\Gamma(\widehat{X}, \widehat{F})$  could have cohomology in negative degrees.

EXAMPLE 1.4. – Fix a quasicoherent flat  $\mathcal{O}_V$ -module  $F$ , assumed to be obtained from an  $A$ -module  $M$  via localization. Then  $\text{R}\Gamma(\widehat{V}, \widehat{F})$  is computed as follows. Fix an ideal  $(g_1, \dots, g_r) \subset A$  with  $V(g_1, \dots, g_r) = \{\mathfrak{m}\}$  set-theoretically (assumed to exist). Let  $C(M; g_1, \dots, g_r) := \bigotimes_{i=1}^r (M \xrightarrow{1} M_{g_i})$  be the displayed Čech complex, and let  $K(M)$  be the cone of the natural map  $C(M; g_1, \dots, g_r) \rightarrow M$ . Then the (termwise)  $f$ -adic completion of  $K$  computes  $\text{R}\Gamma(\widehat{V}, \widehat{F})$ . To see this, observe first that  $K(M)/f^n K(M)$  computes

<sup>(1)</sup> The formal scheme  $\widehat{Y}$  is used as a purely linguistic device to talk about compatible systems of sheaves on each  $Y_n$ , and not in a deeper manner.

$\mathrm{R}\Gamma(V_n, F \otimes_{\mathcal{O}_V}^L \mathcal{O}_{V_n})$ . It follows that the term-wise  $f$ -adic completion of  $K$  computes  $\mathrm{R}\lim \mathrm{R}\Gamma(V_n, F \otimes_{\mathcal{O}_V} \mathcal{O}_{V_n}) \simeq \mathrm{R}\Gamma(\widehat{V}, \widehat{F})$ .

The derived completion functor  $K \mapsto \mathrm{R}\lim(K \otimes_A^L A/f^n)$  already appears implicitly in the above definition. To access its values, recall the following definition:

**DEFINITION 1.5.** – Given an  $A$ -module  $M$ , we define the  $f$ -adic Tate module as  $T_f(M) := \lim M[f^n]$  with transition maps given by powers of  $f$ ; note that  $T_f(M) = 0$  if  $f^N \cdot M = 0$  for some  $N > 0$ .

The Tate module leads to the second of the following two descriptions of the cohomology of a formal completion:

**LEMMA 1.6.** – Let  $Y$  be an  $X$ -scheme such that  $\mathcal{O}_Y$  has bounded  $f^\infty$ -torsion. For  $F \in D(\mathcal{O}_Y)$ , there are exact sequences

$$1 \rightarrow \mathrm{R}^1 \lim H^{i-1}(Y_n, F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n}) \rightarrow H^i(\widehat{Y}, \widehat{F}) \rightarrow \lim H^i(Y, F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n}) \rightarrow 1,$$

and

$$1 \rightarrow \lim H^i(Y, F)/f^n \rightarrow H^i(\widehat{Y}, \widehat{F}) \rightarrow T_f(H^{i+1}(Y, F)) \rightarrow 1.$$

*Proof.* – We first give a proof when  $\mathcal{O}_Y$  has no  $f$ -torsion (which will be the only relevant case in the sequel). The first sequence is then obtained from the formula

$$\mathrm{R}\Gamma(\widehat{Y}, \widehat{F}) \simeq \mathrm{R}\lim \mathrm{R}\Gamma(Y, F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n})$$

and Milnor's exact sequence for  $\mathrm{R}\lim$  (see [18]). Applying the projection formula (since  $A/f^n$  is  $A$ -perfect) to the above gives

$$\mathrm{R}\Gamma(\widehat{Y}, \widehat{F}) \simeq \mathrm{R}\lim (\mathrm{R}\Gamma(Y, F) \otimes_A^L A/f^n).$$

The second sequence is now obtained by applying the derived  $f$ -adic completion functor  $\mathrm{R}\lim(- \otimes_A^L A/f^n)$  to the canonical filtration on  $\mathrm{R}\Gamma(Y, F)$ , which proves the claim. In general, the boundedness of  $f$ -torsion in  $\mathcal{O}_Y$  shows that the map  $\{\mathcal{O}_Y \xrightarrow{f^n} \mathcal{O}_Y\} \rightarrow \{\mathcal{O}_{Y_n}\}$  of projective systems is a (strict) pro-isomorphism, and hence  $\{F \xrightarrow{f^n} F\} \rightarrow \{F \otimes_{\mathcal{O}_Y}^L \mathcal{O}_{Y_n}\}$  is also a pro-isomorphism. Now the previous argument applies.  $\square$

The following conditions on the data  $(A, f)$  will be assumed throughout this subsection; we do *not* assume  $A$  is Noetherian as this will not be true in applications.

**ASSUMPTION 1.7.** – Assume that the data from Notation 1.1 satisfies the following:

- $X$  is integral, i.e.,  $A$  is a domain;
- $j : V \hookrightarrow X$  is a quasi-compact open immersion, i.e.,  $\mathfrak{m}$  is the radical of a finitely generated ideal;
- $H^0(V, \mathcal{O}_V)$  is a finite  $A$ -module;
- $f^N \cdot H^1(V, \mathcal{O}_V) = 0$  for  $N \gg 0$ .