

quatrième série - tome 48 fascicule 4 juillet-août 2015

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Arend BAYER & Brendan HASSETT & Yuri TSCHINKEL

Mori cones of holomorphic symplectic varieties of K3 type

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2015

N. ANANTHARAMAN B. KLEINER
E. BREUILLARD E. KOWALSKI
R. CERF P. LE CALVEZ
A. CHAMBERT-LOIR M. MUSTAȚĂ
I. GALLAGHER L. SALOFF-COSTE

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition / *Publication*

Société Mathématique de France
Institut Henri Poincaré
11, rue Pierre et Marie Curie
75231 Paris Cedex 05
Tél. : (33) 01 44 27 67 99
Fax : (33) 01 40 46 90 96

Abonnements / *Subscriptions*

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 09
Fax : (33) 04 91 41 17 51
email : smf@smf.univ-mrs.fr

Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

© 2015 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593

Directeur de la publication : Marc Peigné
Périodicité : 6 n^{os} / an

MORI CONES OF HOLOMORPHIC SYMPLECTIC VARIETIES OF K3 TYPE

BY AREND BAYER, BRENDAN HASSETT AND YURI TSCHINKEL

ABSTRACT. – We determine the Mori cone of holomorphic symplectic varieties deformation equivalent to the punctual Hilbert scheme on a K3 surface. Our description is given in terms of Markman’s extended Hodge lattice.

RÉSUMÉ. – On détermine les cônes de Mori des variétés symplectiques holomorphes qui se déforment au schéma de Hilbert de points sur une surface K3. Notre description est donnée en termes de structure de Hodge élargie de Markman.

Introduction

Let X be an irreducible holomorphic symplectic manifold. Let $(,)$ denote the Beauville-Bogomolov form on $H^2(X, \mathbb{Z})$; we may embed $H^2(X, \mathbb{Z})$ in $H_2(X, \mathbb{Z})$ via this form. Fix a polarization h on X ; by a fundamental result of Huybrechts [17], X is projective if it admits a divisor class H with $(H, H) > 0$. It is expected that finer birational properties of X are also encoded by the Beauville-Bogomolov form and the Hodge structure on $H^2(X)$, along with appropriate extension data. In particular, natural cones appearing in the minimal model program—the moving cone, the nef cone, the pseudo-effective cone—should have a description in terms of this form.

Now assume X is deformation equivalent to the punctual Hilbert scheme $S^{[n]}$ of a K3 surface S with $n > 1$. Recall that

$$(1) \quad H^2(S^{[n]}, \mathbb{Z})_{(,)} = H^2(S, \mathbb{Z}) \oplus_{\perp} \mathbb{Z}\delta, \quad (\delta, \delta) = -2(n-1)$$

where the restriction of the Beauville-Bogomolov form to the first factor is just the intersection form on S , and 2δ is the class of the locus of non-reduced subschemes. Recall from [20] that for K3 surfaces S , the cone of (pseudo-)effective divisors is the closed cone generated by

$$\{D \in \text{Pic}(S) : (D, D) \geq -2, (D, h) > 0\}.$$

The first attempt to extend this to higher dimensions was [13]. Further work on moving cones was presented in [14, 24], which built on Markman’s analysis of monodromy groups. The

characterization of extremal rays arising from Lagrangian projective spaces $\mathbb{P}^n \hookrightarrow X$ has been addressed in [14, 12] and [3]. The paper [15] proposed a general framework describing all types of extremal rays; however, Markman found counterexamples in dimensions ≥ 10 , presented in [5].

The formalism of Bridgeland stability conditions [7, 8] has led to breakthroughs in the birational geometry of moduli spaces of sheaves on surfaces. The case of punctual Hilbert schemes of \mathbb{P}^2 and del Pezzo surfaces was investigated by Arcara, Bertram, Coskun, and Huizenga [2, 16, 6, 10]. The effective cone on $(\mathbb{P}^2)^{[n]}$ has a beautiful and complex structure as n increases, which only becomes transparent in the language of stability conditions. Bayer and Macri resolved the case of punctual Hilbert schemes and more general moduli spaces of sheaves on K3 surfaces [5, 4]. Abelian surfaces, whose moduli spaces of sheaves include generalized Kummer varieties, have been studied as well [31, 32].

In this note, we extend the results obtained for moduli spaces of sheaves over K3 surfaces to all holomorphic symplectic manifolds arising as deformations of punctual Hilbert schemes of K3 surfaces. Our principal result is Theorem 1 below, providing a description of the Mori cone (and thus dually of the nef cone).

In any given situation, this also leads to an effective method to determine the list of marked minimal models (i.e., birational maps $f: X \dashrightarrow Y$ where Y is also a holomorphic symplectic manifold): the movable cone has been described by Markman [23, Lemma 6.22]; by [14], it admits a wall-and-chamber decomposition whose walls are the orthogonal complements of extremal curves on birational models, and whose closed chambers correspond one-to-one to marked minimal model, as the pull-backs of the corresponding nef cones.

Acknowledgments:

The first author was supported by NSF grant 1101377; the second author was supported by NSF grants 0901645, 0968349, and 1148609; the third author was supported by NSF grants 0968318 and 1160859. We are grateful to Emanuele Macri for helpful conversations, to Eyal Markman for constructive criticism and correspondence, to Claire Voisin for helpful comments on deformation-theoretic arguments in a draft of this paper, and to Ekatarina Amerik for discussions on holomorphic symplectic contractions. We are indebted to the referees for their careful reading of our manuscript. The first author would also like to thank Giovanni Mongardi for discussions and a preliminary version of [25]. Related questions for general hyperkähler manifolds have been treated in [1].

1. Statement of results

Let X be deformation equivalent to the Hilbert scheme of length- n subschemes of a K3 surface. Markman, see [22, Theorem 1.10] and [23, Cor. 9.5], describes an extension of lattices

$$H^2(X, \mathbb{Z}) \subset \tilde{\Lambda}$$

and weight-two Hodge filtrations

$$H^2(X, \mathbb{C}) \subset \tilde{\Lambda}_{\mathbb{C}}$$

with the properties listed below. We will write

$$\theta_X : H^2(X) \subset \tilde{\Lambda}_X$$

to denote the extension of Hodge structures with pairing; here θ_X is defined canonically up to a choice of sign.

- The orthogonal complement of $\theta_X(H^2(X, \mathbb{Z}))$ has rank one, and is generated by a primitive vector of square $2n - 2$ and type $(1, 1)$;
- as a lattice

$$\tilde{\Lambda} \simeq U^4 \oplus (-E_8)^2$$

where U is the hyperbolic lattice and E_8 is the positive definite lattice associated with the corresponding Dynkin diagram;

- any parallel transport operator $\phi : H^2(X, \mathbb{Z}) \rightarrow H^2(X', \mathbb{Z})$ naturally lifts to an isometry of lattices $\tilde{\phi} : \tilde{\Lambda}_X \rightarrow \tilde{\Lambda}_{X'}$ such that

$$\tilde{\phi} \circ \theta_X = \theta_{X'} \circ \phi;$$

the induced action of the monodromy group on $\tilde{\Lambda}/H^2(X, \mathbb{Z})$ is encoded by a character *cov* (see [21, Sec. 4.1]);

- we have the following Torelli-type statement: X_1 and X_2 are birational if and only if there is Hodge isometry

$$\tilde{\Lambda}_{X_1} \simeq \tilde{\Lambda}_{X_2}$$

taking $H^2(X_1, \mathbb{Z})$ isomorphically to $H^2(X_2, \mathbb{Z})$;

- if X is a moduli space $M_v(S)$ of sheaves (or of Bridgeland-stable complexes) over a K3 surface S with Mukai vector v then there is an isomorphism from $\tilde{\Lambda}$ to the Mukai lattice of S taking $H^2(X, \mathbb{Z})$ to v^\perp .

Generally, we use v to denote a primitive generator for the orthogonal complement of $H^2(X, \mathbb{Z})$ in $\tilde{\Lambda}$. Note that $v^2 = (v, v) = 2n - 2$. When $X \simeq M_v(S)$ we may take the Mukai vector v as the generator.

As the dual of θ_X we obtain a homomorphism⁽¹⁾

$$\theta_X^\vee : \tilde{\Lambda}_X \rightarrow H_2(X, \mathbb{Z})$$

which restricts to an inclusion

$$H^2(X, \mathbb{Z}) \subset H_2(X, \mathbb{Z})$$

of finite index. By extension, it induces a \mathbb{Q} -valued Beauville-Bogomolov form on $H_2(X, \mathbb{Z})$.

Assume X is projective. Let $H^2(X)_{\text{alg}} \subset H^2(X, \mathbb{Z})$ and $\tilde{\Lambda}_{\text{alg}} \subset \tilde{\Lambda}_X$ denote the algebraic classes, i.e., the integral classes of type $(1, 1)$. Since the orthogonal complement of $i_X(H^2(X))$ is generated by an algebraic class, it follows dually that $a \in \tilde{\Lambda}_X$ is of type $(1, 1)$ if and only if $\theta^\vee(a)$ is. The Beauville-Bogomolov form on $H^2(X)_{\text{alg}}$ has signature $(1, \rho(X) - 1)$, where $\rho(X) = \dim(H_{\text{alg}}^2(X))$. The *Mori cone* of X is defined as the closed cone in $H_2(X, \mathbb{R})_{\text{alg}}$ containing the classes of algebraic curves in X . The *positive cone* (or more accurately, non-negative cone) in $H^2(X, \mathbb{R})_{\text{alg}}$ is the closure of the connected component of the cone

$$\{D \in H^2(X, \mathbb{R})_{\text{alg}} : D^2 > 0\}$$

⁽¹⁾ We will often drop the subscript X from the notation when the context is clear.