

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

A NEW TWO-PARAMETER FAMILY OF ISOMONODROMIC DEFORMATIONS OVER THE FIVE PUNCTURED SPHERE

Arnaud Girand

**Tome 144
Fascicule 2**

2016

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 339-368

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 2, tome 144, juin 2016

Comité de rédaction

Valérie BERTHÉ	Marc HERZLICH
Gérard BESSON	O'Grady KIERAN
Emmanuel BREUILLARD	Julien MARCHÉ
Yann BUGEAUD	Emmanuel RUSS
Jean-François DAT	Christophe SABOT
Charles FAVRE	Wilhelm SCHLAG
Raphaël KRIKORIAN (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 178 €, hors Europe : 194 € (\$ 291)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© *Société Mathématique de France* 2016

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

A NEW TWO-PARAMETER FAMILY OF ISOMONODROMIC DEFORMATIONS OVER THE FIVE PUNCTURED SPHERE

BY ARNAUD GIRAND

ABSTRACT. — The object of this paper is to describe an explicit two-parameter family of logarithmic flat connections over the complex projective plane. These connections have dihedral monodromy and their polar locus is a prescribed quintic composed of a conic and three tangent lines. By restricting them to generic lines we get an algebraic family of isomonodromic deformations of the five-punctured sphere. This yields new algebraic solutions of a Garnier system. Finally, we use the associated Riccati one-forms to construct and prove the integrability (in the transversally projective sense) of a subfamily of Lotka-Volterra foliations.

RÉSUMÉ (*Une nouvelle famille à deux paramètres de déformations isomonodromiques sur la sphère à cinq trous*)

Le but de cet article est de décrire une famille explicite à deux paramètres de connexions logarithmiques plates au dessus du plan projectif complexe. Ces connexions sont à monodromie diédrale et leur lieu polaire est une quintique prescrite, composée d'une conique et de trois droites tangentes. Par restriction aux droites génériques, on obtient alors une famille algébrique de déformations isomonodromiques de la sphère à cinq trous. Ceci livre de nouvelles solutions algébriques d'un système de Garnier. Enfin, nous utilisons les formes de Riccati associées à ces connexions pour construire et montrer l'intégrabilité (au sens transversalement projectif) d'une sous-famille de feuilletages de Lotka-Volterra.

Texte reçu le 23 juillet 2015, révisé le 30 septembre 2015, accepté le 13 octobre 2015.

ARNAUD GIRAND, IRMAR (UMR 6625 du CNRS), Université de Rennes 1, campus de Beaulieu. Bât. 22-23. 35042 Rennes Cedex • *E-mail* : arnaud.girand@univ-rennes1.fr
2010 Mathematics Subject Classification. — 14E22, 20G05, 20G20, 32D15, 32G08, 32G34, 34M50, 34M56, 51N15, 55R10.

Key words and phrases. — Isomonodromic deformations, Garnier systems, logarithmic flat connections, transversally projective foliations, Lotka-Volterra systems, Riccati forms, dihedral monodromy groups.

1. Introduction

In this section we describe the main result of this paper, namely an explicit construction of a two-parameter family of logarithmic flat connections over the complement of a particular quintic curve in $\mathbb{P}^2_{\mathbb{C}}$. The restriction of any element of this family to generic lines in the projective plane gives an isomonodromic deformation over the five punctured sphere, to which we can associate an algebraic solution of some Hamiltonian system of partial differential equations, namely the Garnier-2 system.

1.1. Topology of the complement of a particular plane quintic. — In this paper, we concern ourselves with setting up a two-parameter family of logarithmic flat $\mathfrak{sl}_2(\mathbb{C})$ -connections over \mathbb{P}^2 with a specific polar locus, namely a quintic curve Q composed of a circle and three tangent lines. More precisely, in homogeneous coordinates $[x : y : t]$, Q is defined, up to $PGL_3(\mathbb{C})$ action, by the equation

$$xyt(x^2 + y^2 + t^2 - 2(xy + xt + yt)) = 0 .$$

Before stating our main result, let us specify what we are looking for: we want to find a family of rank two logarithmic flat connections over \mathbb{P}^2 with polar locus equal to some small degree curve and “interesting monodromy”. We will show that it is possible to do so with the quintic Q define above.

DEFINITION 1.1. — We say that the monodromy representation associated with a rank two logarithmic flat $\mathfrak{sl}_2(\mathbb{C})$ -connection over $\mathbb{P}^2 - Q$ is non-degenerate if

- its image forms an irreducible subgroup of $SL_2(\mathbb{C})$;
- its local monodromy (see Definition 2.3) around any irreducible component of Q is projectively non-trivial (i.e is non-trivial in $PSL_2(\mathbb{C})$).

In order to establish the existence of such representations, we use the following result by Degtyarev.

PROPOSITION 1.2 (Degtyarev, 1999 [5]). — *The fundamental group Γ of the complement of a smooth conic and three tangent lines in \mathbb{P}^2 admits the following presentation:*

$$\Gamma \cong \langle a, b, c \mid (ab)^2(ba)^{-2} = (ac)^2(ca)^{-2} = [b, c] = 1 \rangle .$$

More precisely, Degtyarev proves that we can take a (resp. b, c) to be a loop realising the local monodromy (see Definition 2.3) around the conic $\mathcal{C} := (x^2 + y^2 + t^2 - 2(xy + xt + yt) = 0)$ (resp. the lines $(y = 0), (x = 0)$), as illustrated in the left-hand side of Fig. 1. Also note that the fundamental group of the intersection of $\mathbb{P}^2 - Q$ with any generic line is isomorphic to the free group $\mathbf{F}_4 := \langle d_1, \dots, d_5 \mid d_1 \dots d_5 = 1 \rangle$; the Lefschetz hyperplane theorem (see [15],




FIGURE 1. Fundamental group of the complement of the quintic Q in \mathbb{P}^2 and restriction to a generic line.

Theorem 7.4) tells us that the natural morphism $\tau : F_4 \rightarrow \Gamma$ is onto. Moreover we know from the explicit Van Kampen method given in Subsection 4.1 of [5] that the group Γ can be computed by taking the four free generators of the fundamental group of the intersection of $\mathbb{P}^2 - Q$ with any generic line and adding some braid monodromy relations. Thus, if we chose a line going through the base point used to define a, b and c then τ is given (up to a permutation of the d_i) by (see the right-hand side of Fig. 1):

$$\begin{aligned} d_1 &\mapsto b \\ d_2 &\mapsto a \\ d_3 &\mapsto bab^{-1} \\ d_4 &\mapsto c \\ d_5 &\mapsto (abac)^{-1} . \end{aligned}$$

In particular, any non-degenerate representation ρ of Γ must satisfy

$$\rho(a), \rho(b), \rho(c), \rho(abac) \neq \pm I_2 .$$

PROPOSITION 1.3. — *The only (up to conjugacy) family of non-degenerate representations of Γ into $SL_2(\mathbb{C})$ is as follows:*

$$\rho_{u,v} : a \mapsto \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad b \mapsto \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}, \quad c \mapsto \begin{pmatrix} v & 0 \\ 0 & v^{-1} \end{pmatrix}, \quad \text{for } u, v \in \mathbb{C}^* .$$

1.2. Main results. — The core of this paper will be devoted to proving the following theorem, in which we explicitly construct the announced family of rank two logarithmic flat connections.