

Antoine Ducros

FAMILIES OF BERKOVICH SPACES

ASTÉRISQUE 400

Société Mathématique de France 2018

Astérisque est un périodique de la Société mathématique de France
Numéro 400

Comité de rédaction

Ahmed ABBES
Viviane BALADI
Laurent BERGER
Philippe BIANE
Hélène ESNAULT

Éric VASSEROT (dir.)

Philippe EYSSIDIEUX
Damien GABORIAU
Michael HARRIS
Fabrice PLANCHON
Pierre SCHAPIRA

Diffusion

Maison de la SMF B.P. 67 13274 Marseille Cedex 9 France christian.munusami@smf.emath.fr	AMS P.O. Box 6248 Providence RI 02940 USA www.ams.org
--	--

Tarifs 2018

Vente au numéro : 50 (\$ 75)
Abonnement électronique : 500 (\$ 750)
Abonnement avec supplément papier : 665 , hors Europe : 718 (\$ 1077)
Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris Cedex 05, France
Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96
nathalie.christiaen@smf.emath.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2018

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0303-1179 (print) 2492-5926 (electronic)
ISBN 978-2-85629-885-5

Stéphane SEURET
Directeur de la publication

ASTÉRISQUE 400

FAMILIES OF BERKOVICH SPACES

Antoine Ducros

Société Mathématique de France 2018

Publié avec le concours du Centre National de la Recherche Scientifique

A. DUCROS

Sorbonne Université, Université Paris Diderot, CNRS, Institut de Mathématiques de Jussieu-Paris Rive Gauche, IMJ-PRG, F-75005, Paris, France.

E-mail : antoine.ducros@imj-prg.fr

Url : <http://www.imj-prg.fr/~antoine.ducros>

2010 Mathematics Subject Classification. — 14G22, 14A99.

Key words and phrases. — Berkovich spaces, flatness, relative properties, constructible loci.

While writing this book the author was supported by the french *Agence nationale de la recherche* (ANR), through the projects *Espaces de Berkovich* (ANR 07-JCJC-0004-CSD5), *Valuations, combinatoire et théorie des modèles* (ANR-13-BS01-0006), and *Définissabilité en géométrie non archimédienne* (ANR-15-CE40-0008). He benefited from two ROSI AND MAX VARON professorships at Weizmann Institute (Rehovot, Israel), both with an invitation by Vladimir Berkovich: from January 15th till March 14th, 2009 and from April 19th till July 18th, 2015. He was a junior member of the *Institut universitaire de France* from October 2012 till September 2017.

FAMILIES OF BERKOVICH SPACES

Antoine Ducros

Abstract. — This book is devoted to the systematic study of relative properties in the context of Berkovich analytic spaces. We first develop a theory of flatness in this setting. After showing through a counter-example that naive flatness cannot be the right notion because it is not stable under base change, we define flatness by *requiring* invariance under base change, and we study a first important class of flat morphisms, that of quasi-smooth ones.

We then show the existence of local *dévissages* (in the spirit of Raynaud and Gruson's work) for coherent sheaves, which we use, together with a study of the local rings of "generic fibers" of morphisms, to prove that a *boundaryless*, naively flat morphism is flat.

After that we prove that the image of a compact strictly analytic space by a flat morphism is an analytic domain of the target and that it admits, when the source is strictly analytic, a compact, flat multisection (*i.e.*, a compact, flat cover of relative dimension zero over which there is a section). The image of a compact analytic space by a flat morphism had already been described by Raynaud in the rigid-analytic setting, but our method is completely different.

In the last part of this work we study where various interesting pointwise relative properties are satisfied. We first prove that the flat locus of a given morphism of analytic spaces is a Zariski-open subset of the source (we follow the method that was introduced by Kiehl for the complex analytic analogue of this statement). We then look at the loci at which a point satisfies various commutative algebra properties *on its fiber*: being geometrically regular, geometrically R_m , complete intersection, or Gorenstein; being S_m or Cohen-Macaulay. We prove that the results we could expect actually hold: these loci are (locally) Zariski-constructible, and Zariski-open under suitable extra assumptions (flatness, and also equidimensionality for S_m and geometric R_m); for that purpose, we first study the general properties of the locally Zariski-constructible subsets of an analytic space.

Résumé (Familles d'espaces de Berkovich). — Ce livre est consacré à une étude systématique des propriétés relatives dans le contexte des espaces de Berkovich. Nous commençons par développer une théorie de la platitude dans ce cadre. L'acceptation naïve de cette notion est inadaptée : nous montrons en effet par un contre-exemple qu'elle n'est pas stable par changement de base, ce qui nous conduit à *imposer* cette stabilité dans la définition. Nous étudions une première classe importante de morphismes plats : celle des morphismes *quasi-lisses*.

Nous montrons ensuite l'existence de *dévissages* locaux (dans l'esprit de Raynaud et Gruson) pour les faisceaux cohérents. Joint à une étude des anneaux locaux des fibres « génériques » des morphismes, cela nous permet de montrer qu'un morphisme *sans bord* qui est plat au sens naïf l'est encore au nôtre.

Puis nous démontrons que l'image d'un espace strictement analytique compact par un morphisme plat est un domaine analytique du but, et qu'elle admet, lorsque la source est strictement analytique, une multisection compacte et plate (*i.e.* un revêtement plat, compact, et de dimension relative nulle sur lequel le morphisme considéré possède une section). L'image d'un espace analytique compact par un morphisme plat avait déjà été décrite par Raynaud dans le contexte rigide-analytique, mais notre méthode est complètement différente.

Dans la dernière partie de ce travail, nous étudions les lieux de validité sur la source de certaines propriétés relatives. Nous y démontrons pour commencer que le lieu de platitude d'un morphisme d'espaces analytiques est un ouvert de Zariski de la source (nous suivons la méthode utilisée par Kiehl pour établir l'assertion correspondante en géométrie analytique complexe). Cela nous permet de procéder à l'investigation systématique des ensemble de points satisfaisant *dans leur fibre* les propriétés classiques de l'algèbre commutative : être géométriquement régulier, géométriquement R_m , d'intersection complète ou de Gorenstein ; être S_n ou de Cohen-Macaulay. Nous prouvons que les énoncés auxquels on peut s'attendre sont effectivement vérifiés : ces lieux de validités sont (localement) Zariski-constructibles, et sont des ouverts de Zariski sous certaines hypothèses supplémentaires (platitude, ainsi qu'équidimensionalité pour les propriétés R_m ou S_m) ; dans ce but, nous procédons tout d'abord à une étude générale des parties localement Zariski-constructibles d'un espace analytique.

CONTENTS

Introduction	1
1. First step: flatness in the Berkovich setting	1
2. Loci of validity	5
3. About our proofs	8
Acknowledgements	11
1. Background material	13
1.1. Prerequisites and basic conventions	13
1.2. Analytic geometry: basic definitions	16
1.3. Coherent sheaves, Zariski topology and closed immersions	20
1.4. Dimension theory	25
1.5. Irreducible components	29
2. Algebraic properties in analytic geometry	35
2.1. Analytification of schemes, algebraic properties of analytic rings	36
2.2. A rather abstract categorical framework	39
2.3. Formalisation of algebraic properties	42
2.4. Validity in analytic geometry	47
2.5. Fibers of coherent sheaves	50
2.6. Ground field extension	52
2.7. Complements on analytifications	55
3. Germs, Temkin's reduction and Γ-strictness	63
3.1. Γ -strictness	64
3.2. Analytic germs	66
3.3. Around graded Riemann-Zariski spaces	69
3.4. Temkin's construction	72
3.5. Temkin's reduction and Γ -strictness	75

4. Flatness in analytic geometry	83
4.1. Naive and non-naive flatness	84
4.2. Algebraic flatness <i>versus</i> analytic flatness	88
4.3. The flat, (locally) finite morphisms	91
4.4. Naive flatness is not preserved by base change	93
4.5. Analytic flatness has the expected properties	97
5. Quasi-smooth morphisms	105
5.1. Reminders about the sheaf of relative differentials	106
5.2. Quasi-smoothness: definition and first properties	108
5.3. Quasi-smoothness, flatness and fiberwise geometric regularity	112
5.4. Links with étale and smooth morphisms	116
5.5. Transfer of algebraic properties	121
6. Generic fibers in analytic geometry	125
6.1. Preliminary lemmas	127
6.2. Relative polydiscs inside relative smooth spaces	129
6.3. Local rings of generic fibers	134
7. Images of morphisms: local results	141
7.1. Maps between Riemann-Zariski Spaces: the non-graded case	142
7.2. Maps between Riemann-Zariski spaces: the general case	144
7.3. Applications to analytic geometry	149
7.4. Complement: around quantifier elimination in the theory ACVF	152
8. Dévissages à la Raynaud-Gruson	157
8.1. Universal injectivity and flatness	158
8.2. Dévissages: definition and existence	164
8.3. Flatness can be checked naively in the inner case	170
8.4. The relative CM property	172
9. Quasi-finite multisections and images of maps	175
9.1. Flat, quasi-finite multisections of flat maps	177
9.2. Images of maps	184
10. Constructible loci	187
10.1. Constructibility in analytic spaces	188
10.2. The diagonal trick	193
10.3. The flat locus	200
10.4. The fiberwise closure of a locally constructible set	203
10.5. The fiberwise exactness locus	209
10.6. Regular sequences	215
10.7. The main theorem	217

11. Algebraic properties: target, fibers and source	229
11.1. A weak “dominant factorization theorem”	230
11.2. The axiomatic setting	233
11.3. The main theorem	235
A. Graded commutative algebra	241
A.1. Basic definitions	241
A.2. Graded linear algebra	243
A.3. Graded algebras and graded extensions	245
A.4. Graded valuations	248
Index of notation	253
Index	255
Bibliography	259

INTRODUCTION

This book is, roughly speaking, devoted to a systematic investigation of *families* of objects in Berkovich's non-Archimedean analytic geometry (see [Ber90], [Ber93]). More precisely, let us assume that we are given a morphism $Y \rightarrow X$ between analytic spaces, and an object D in Y of a certain kind (think of the space Y itself, or a coherent sheaf, or a complex of coherent sheaves...). Every point x of X gives then rise to an object D_x , living on the fiber Y_x , and we thus get in some sense an analytic family of objects parametrized by the space X . The quite vague problem we would like to address is the following: *how do the object D_x and its relevant properties vary?* Of course, such questions have been intensively studied for a long time in *algebraic* geometry, especially by Grothendieck and his school, and our guideline has been to establish analytic avatars of their results every time it was possible. Let us now give a quick overview of our work.

1. First step: flatness in the Berkovich setting

1.1. Motivation. — In scheme theory, the key notion upon which the study of families is based is *flatness*. This is a property of families of *coherent sheaves*, which encodes more or less the intuitive idea of a *reasonable* variation (this is why there is almost always a flatness assumption in the description of moduli problems).

The point is that the study of general families is often reduced (typically, through a suitable stratification of the base scheme) to the case where some of the coherent sheaves involved are flat over the parameter space, which is easier to handle.

But we would like to emphasize that flatness is also a crucial technical tool for many other purposes. Let us mention for example:

- descent theory;
- the first occurrence of flatness in algebraic geometry, in the celebrated paper GAGA by Serre [Ser56], where the following plays a major role: if X is a complex algebraic variety and if X^{an} denotes the corresponding analytic space, then for every $x \in X(\mathbf{C})$ the ring $\mathcal{O}_{X^{\text{an}}, x}$ is flat over $\mathcal{O}_{X, x}$.

By analogy, our first step toward the understanding of analytic families has been the development of a theory of flatness in Berkovich geometry, which is the core of this book; and similarly to what happens for schemes, we hope that it will have many applications beyond the study of families.

In fact, flatness in non-Archimedean geometry had already been considered, but in the *rigid-analytic* setting, following ideas of Raynaud; see Bosch [BL93a] and Lütkebohmert [BL93b], as well as the more comprehensive recent study Abbes [Abb10]. We now give a more precise discussion of rigid-analytic flatness, before saying some words concerning our definition in the Berkovich framework.

1.2. Flatness in rigid geometry. — The definition of flatness in the rigid setting is as simple as one may hope: if $f : Y \rightarrow X$ is a morphism between rigid spaces and if \mathcal{F} is a coherent sheaf on Y , it is rig-flat over X at a point $y \in Y$ if it is flat over X at y *in the sense of the theory of locally ringed spaces*; i.e., the stalk \mathcal{F}_y is a flat $\mathcal{O}_{X,f(y)}$ -module (this is the definition by Abbes; the original one by Bosch and Lütkebohmert was slightly different, but both are easily seen to be equivalent using the good properties of the completion of local rings as far as flatness is concerned, cf. [SGA1], Exposé IV, Cor. 5.8).

Flatness in the above sense behaves well: it is stable under base change and ground field extension. But contrary to what happens in scheme theory, this is in no way obvious, because base change and ground field extension are defined using *completed* tensor products. Roughly speaking, the proofs proceed as follows (see [BL93a], [BL93b] and [Abb10]):

- the study of *rigid* flatness is reduced to that of *formal* flatness through formal avatars of Raynaud-Gruson flattening techniques, which are used to build a flat formal model of any given rig-flat coherent sheaf;
- the study of *formal* flatness is reduced to that of *algebraic* flatness, in a more standard way, upon dividing by various ideals of definition and using flatness criteria in the spirit of [SGA1], Exposé IV.

Let us mention that this general strategy (formal flattening and reduction modulo an ideal of definition to replace an analytic problem with an algebraic one) was also used by Raynaud to prove the following fact: if $\varphi : Y \rightarrow X$ is a flat morphism between affinoid rigid spaces, $\varphi(Y)$ is a finite union of affinoid domains of X (cf. [BL93b], Cor. 5.11).

1.3. Flatness in Berkovich geometry. — We fix from now on and for the remaining part of this introduction a complete, non-Archimedean field k . We shall only consider *Berkovich* analytic spaces. Any analytic space X comes with a usual topology, but also with a set-theoretic Grothendieck topology which refines it, the so-called *G-topology* – the corresponding site is denoted by X_G ; the archetypal example of a G-covering is