

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

LOWER BOUNDS FOR RANKS OF MUMFORD-TATE GROUPS

Martin Orr

Tome 143
Fascicule 2

2015

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 229-246

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 2, tome 143, juin 2015

Comité de rédaction

Gérard BESSON	Daniel HUYBRECHTS
Emmanuel BREUILLARD	Julien MARCHÉ
Antoine CHAMBERT-LOIR	Christophe SABOT
Charles FAVRE	Laure SAINT-RAYMOND
Pascal HUBERT	Wilhelm SCHLAG
Marc HERZLICH	

Raphaël KRIKORIAN (dir.)

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 176 €, hors Europe : 193 € (\$ 290)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© *Société Mathématique de France* 2015

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

LOWER BOUNDS FOR RANKS OF MUMFORD-TATE GROUPS

BY MARTIN ORR

ABSTRACT. — Let A be a complex abelian variety and G its Mumford-Tate group. Supposing that the simple abelian subvarieties of A are pairwise non-isogenous, we find a lower bound for the rank $\mathrm{rk} G$ of G , which is a little less than $\log_2 \dim A$. If we suppose furthermore that $\mathrm{End} A$ is commutative, then we can improve this lower bound to $\mathrm{rk} G \geq \log_2 \dim A + 2$ and prove that this is sharp. We also obtain the same results for the rank of the ℓ -adic monodromy group of an abelian variety defined over a number field.

RÉSUMÉ (*Minoration des rangs de groupes de Mumford-Tate*). — Soit A une variété abélienne complexe et G son groupe de Mumford-Tate. En supposant que les sous variétés abéliennes simples de A sont deux à deux non-isogènes, on trouve une minoration du rang $\mathrm{rk} G$ de G , légèrement inférieure à $\log_2 \dim A$. Si de plus on suppose que $\mathrm{End} A$ est commutatif, alors on peut améliorer cette borne en $\mathrm{rk} G \geq \log_2 \dim A + 2$, et montrer que cette borne-ci est optimale. On obtient les mêmes résultats pour le rang du groupe de monodromie ℓ -adique d'une variété abélienne définie sur un corps de nombres.

Texte reçu le 29 juin 2011, révisé le 19 décembre 2012, accepté le 21 juin 2013.

MARTIN ORR, Université Paris-Sud, Bat. 425, 91400 Orsay, France •
E-mail : martin.orr@math.u-psud.fr

2010 Mathematics Subject Classification. — 14K15, 14G25, 22E55.

Key words and phrases. — Abelian varieties, Mumford-Tate groups.

1. Introduction

Let A be a complex abelian variety of dimension g , whose simple abelian subvarieties are pairwise non-isogenous. In this paper we will establish a lower bound for the rank of the Mumford-Tate group of A . The Mumford-Tate group is an algebraic group over \mathbb{Q} defined via the Hodge theory of A (see Section 2 below for the definition). The same argument will also establish a lower bound for the rank of the ℓ -adic monodromy groups G_ℓ , in the case where A is defined over a number field. The ℓ -adic monodromy group is the Zariski closure of the image of the Galois representation on the ℓ -adic Tate module of A . Our main theorems are the following:

THEOREM 1.1. — *Let A be an abelian variety of dimension g such that $\text{End } A$ is commutative. Let G be the Mumford-Tate group or the ℓ -adic monodromy group of A . Then $\text{rk } G \geq \log_2 g + 2$.*

THEOREM 1.2. — *Let A be an abelian variety of dimension g whose simple abelian subvarieties are pairwise non-isogenous. Let G be the Mumford-Tate group or the ℓ -adic monodromy group of A . If $n = \text{rk } G$, then*

$$n + \alpha(n)\sqrt{n \log n} \geq \log_2 g + 2$$

for a function $\alpha : \mathbb{N}_{\geq 2} \rightarrow \mathbb{R}$ satisfying $\alpha(n) < 2$ for all n and $\alpha(n) \rightarrow 1/\log 2 = 1.44\dots$ as $n \rightarrow \infty$.

Each of these theorems is an instance of a more general bound for weak Mumford-Tate triples, which are defined in Section 2. These more general bounds are Theorems 4.1 and 4.4 respectively. These would apply also for example to the analogue of the Mumford-Tate group for a Hodge-Tate module of weights 0 and 1.

Theorem 1.1 was proved by Ribet in the case of an abelian variety with complex multiplication [13]. Our proof is a generalisation of his, relying on the fact that the defining representation of the Mumford-Tate group or ℓ -adic monodromy group has minuscule weights.

The condition on simple subvarieties in Theorem 1.2 is necessary: taking products of copies of the same simple abelian variety increases the dimension without changing the rank of the Mumford-Tate group. Indeed, if A is isogenous to $\prod_i A_i^{m_i}$ where the A_i are simple and pairwise non-isogenous, then according to [4] Lemme 2.2,

$$\text{MT}(A) \cong \text{MT}\left(\prod_i A_i\right).$$

Hence Theorem 1.2 implies that for a general abelian variety A , if n denotes the rank of either the Mumford-Tate group or the ℓ -adic monodromy group of A , then

$$n + \alpha(n)\sqrt{n \log n} \geq \log_2 \left(\sum_i \dim A_i \right) + 2$$

where the A_i are one representative of each isogeny class of simple abelian subvarieties of A .

The condition of having pairwise non-isogenous simple abelian subvarieties can be interpreted via the endomorphism algebra like the condition in Theorem 1.1: it is equivalent to $\text{End } A \otimes_{\mathbb{Z}} \mathbb{Q}$ being a product of division algebras. Note also that $\text{End } A$ being commutative implies the condition of Theorem 1.2. (Throughout this paper, $\text{End } A$ means the endomorphisms of A after extension of scalars to an algebraically closed field.)

Let G be either the Mumford-Tate group or the ℓ -adic monodromy group of A . It is well known that the rank of G is at most $g + 1$, and that this upper bound is achieved for a generic abelian variety. Indeed, if g is odd and $\text{End } A = \mathbb{Z}$, then $\text{rk } G$ is always $g + 1$ [16]. So in this case the bound in Theorem 1.1 is far from sharp.

On the other hand if g is a power of 2, then there are abelian varieties for which the bound in Theorem 1.1 is achieved (even with $\text{End } A = \mathbb{Z}$). We construct such examples in Section 5. The exact bound for a given g is very sensitive to the prime factors of g . Equality can happen only when g is a power of 2 (for the trivial reason that otherwise $\log_2 g \notin \mathbb{Z}$) but even near-equality can only occur when g has many small prime factors. This was made precise by Dodson in the complex multiplication case [2], and it is possible that something similar could be proved in general.

Theorem 1.2 is not sharp. The function $\alpha(n)$ is specified exactly in Section 4, but it is likely that this could be improved on, perhaps to something which goes to 0 as $n \rightarrow \infty$. In Section 5, we construct a family of examples showing that Theorem 1.2 cannot be improved to $n + k \geq \log_2 g$ for any constant k .

We can deduce a lower bound for the growth of the degrees of the division fields $K(A[\ell^n])$ (for ℓ a fixed prime number) as a straightforward consequence of Theorem 1.1.

COROLLARY 1.3. — *Let A be an abelian variety of dimension g over a number field K , and ℓ a prime number. If $\text{End } A$ is commutative, then there is a constant $C(A, K, \ell)$ such that*

$$[K(A[\ell^n]) : K] \geq C(A, K, \ell) \cdot \ell^{n(\log_2 g + 2)}.$$