

*quatrième série - tome 51      fascicule 1      janvier-février 2018*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Anne QUÉGUINER-MATHIEU & Jean-Pierre TIGNOL

*Outer automorphisms of classical algebraic groups*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

**Responsable du comité de rédaction / *Editor-in-chief***

Emmanuel KOWALSKI

**Publication fondée en 1864 par Louis Pasteur**

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

**Comité de rédaction au 1<sup>er</sup> janvier 2018**

P. BERNARD      A. NEVES  
S. BOUCKSOM    J. SZEFTEL  
E. BREUILLARD   S. VŨ NGỌC  
R. CERF          A. WIENHARD  
G. CHENEVIER    G. WILLIAMSON  
E. KOWALSKI

**Rédaction / *Editor***

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

**Édition / *Publication***

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

**Abonnements / *Subscriptions***

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

**Tarifs**

Europe : 540 €. Hors Europe : 595 € (\$893). Vente au numéro : 77 €.

---

© 2018 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret  
Périodicité : 6 n<sup>os</sup> / an

# OUTER AUTOMORPHISMS OF CLASSICAL ALGEBRAIC GROUPS

BY ANNE QUÉGUINER-MATHIEU AND JEAN-PIERRE TIGNOL

---

**ABSTRACT.** – The so-called Tits class associated to an adjoint absolutely almost simple algebraic group provides a cohomological obstruction for this group to admit an outer automorphism. If the group has inner type, this obstruction is the only one. In this paper, we prove this is not the case for classical groups of outer type, except for groups of type  ${}^2A_n$  with  $n$  even, or  $n = 5$ . More precisely, we prove a descent theorem for exponent 2 and degree 6 algebras with unitary involution, which shows that their automorphism groups have outer automorphisms. In all other relevant classical types, namely  ${}^2A_n$  with  $n$  odd,  $n \geq 3$  and  ${}^2D_n$ , we provide explicit examples where the Tits class obstruction vanishes, and yet the group does not have outer automorphisms. As a crucial tool, we use “generic” sums of algebras with involution.

**RÉSUMÉ.** – À tout groupe algébrique absolument presque simple de type adjoint est associée une classe de cohomologie connue sous le nom de «classe de Tits» qui donne une obstruction à l’existence d’automorphismes extérieurs. Si le groupe est de type intérieur, il n’y a pas d’autre obstruction. Dans ce travail, nous montrons qu’il n’en va pas de même pour les groupes classiques de type extérieur, sauf pour les groupes de type  ${}^2A_n$  avec  $n$  pair ou  $n = 5$ . Plus précisément, nous établissons pour les algèbres à involution unitaire de degré 6 et d’exposant 2 un théorème de descente qui montre que les groupes d’automorphismes de ces algèbres ont des automorphismes extérieurs. Pour les types  ${}^2A_n$  avec  $n$  impair,  $n \geq 3$ , et les types  ${}^2D_n$ , nous construisons des exemples explicites où l’obstruction donnée par la classe de Tits est nulle alors que le groupe ne possède pas d’automorphisme extérieur. Un outil crucial de nos constructions est la somme «générique» d’algèbres à involution.

## 1. Introduction

Every automorphism of an absolutely almost simple algebraic group scheme  $G$  of adjoint type over an arbitrary field  $F$  induces an automorphism of its Dynkin diagram  $\Delta$ . Inner

---

The first author acknowledges the support of the French Agence Nationale de la Recherche (ANR) under reference ANR-12-BL01-0005. The second author is grateful to the first author and the Université Paris 13 for their hospitality while the work for this paper was carried out. He acknowledges support from the Fonds de la Recherche Scientifique-FNRS under grant n° J.0014.15.

automorphisms of  $G$  act trivially on  $\Delta$ , and there is an exact sequence of algebraic group schemes

$$(1) \quad 1 \rightarrow G \rightarrow \mathbf{Aut}(G) \rightarrow \mathbf{Aut}(\Delta) \rightarrow 1,$$

see [2, exp. XXIV, 1.3, 3.6]. If  $G$  is split, the corresponding sequence of groups of rational points is exact and split, see [6, (25.16)], [11, §16.3]. Therefore, a split adjoint group  $G$  admits outer automorphisms if and only if its Dynkin diagram admits automorphisms, i.e., if  $G$  has type  $A_n$  with  $n \geq 2$ ,  $D_n$  with  $n \geq 3$  or  $E_6$ . Moreover, in all three cases,  $\mathbf{Aut}(\Delta)(F)$  lifts to an isomorphic subgroup in  $\mathbf{Aut}(G)(F)$ . This property does not hold generally for nonsplit groups. For instance, if  $G$  is the connected component of the identity in the group scheme of automorphisms of a central simple  $F$ -algebra with quadratic pair  $(A, \sigma, f)$ , then  $G$  has no outer automorphisms if  $A$  is not split by the quadratic étale  $F$ -algebra defined by the discriminant of the quadratic pair, see §2.2 below. More generally, Garibaldi identified in [4, §2] a cohomological obstruction to the existence of outer automorphisms of an arbitrary absolutely almost simple algebraic group scheme  $G$ : the group  $\mathbf{Aut}(\Delta)(F)$  acts on  $H^2(F, C)$ , where  $C$  is the center of the simply connected group scheme isogenous to  $G$ , and the Tits class  $t_G \in H^2(F, C)$  is invariant under the action of the image of  $\mathbf{Aut}(G)(F)$  in  $\mathbf{Aut}(\Delta)(F)$ . Therefore, automorphisms of  $\Delta$  that do not leave  $t_G$  invariant do not lift to outer automorphisms of  $G$ . For adjoint or simply connected groups of inner type, Garibaldi showed in [4, §2] that this is the only obstruction to the lifting of automorphisms of  $\Delta$ . As he explains in [4, Thm 1.1] this has interesting consequences in Galois cohomology. In a subsequent paper, Garibaldi-Petersson [5, Conjecture 1.1.2] conjectured that this Tits class obstruction is the only obstruction, also for adjoint or simply connected groups of outer type.

In this paper, we provide a complete answer to the question raised by Garibaldi and Petersson for groups of outer type A and D, leaving aside triality groups (see the Appendix). Thus, in all the cases we consider,  $\mathbf{Aut}(\Delta)(F)$  has order 2. Our main goal is to compare the following three conditions, listed from weaker to stronger:

- (Out 1): The Tits class  $t_G$  is fixed under  $\mathbf{Aut}(\Delta)(F)$ ;
- (Out 2):  $G$  admits an outer automorphism defined over  $F$ ;
- (Out 3):  $G$  admits an outer automorphism of order 2 defined over  $F$ .

Under condition (Out 2), the sequence

$$1 \rightarrow G(F) \rightarrow \mathbf{Aut}(G)(F) \rightarrow \mathbf{Aut}(\Delta)(F) \rightarrow 1$$

is exact, and under condition (Out 3), it is split. In [4], Garibaldi proves that all three conditions are equivalent if  $G$  has inner type A or D (see Remarks 2.3 and 2.7). This is not the case for groups of outer type, and our main result is the following:

**THEOREM 1.1.** – *Let  $G$  be an absolutely almost simple adjoint or simply connected algebraic group scheme of type  ${}^2A_n$ , with  $n \geq 2$ , or  ${}^2D_n$ , with  $n \geq 3$ .*

- (1) *If  $G$  has type  ${}^2A_n$ , with  $n$  even, or  ${}^2A_5$ , then conditions (Out 1), (Out 2) and (Out 3) are equivalent.*
- (2) *In all the other types, there are examples of groups for which (Out 1) holds and (Out 2) does not hold, and examples of groups for which (Out 2) holds and (Out 3) does not hold.*

In other words, assertion (2) says there are examples where the condition on the Tits class is satisfied, and yet  $G$  does not have any outer automorphism, and examples where  $G$  has an outer automorphism, but no outer automorphism of order 2. In particular, this disproves Conjecture 1.1.2 in [5], and provides examples of simply connected absolutely simple algebraic group schemes  $G$  for which the Galois cohomology sequence

$$H^1(F, C) \rightarrow H^1(F, G) \rightarrow H^1(F, \mathbf{Aut}(G))$$

from [4, Thm 11(b)] (where  $C$  is the center of  $G$ ) is not exact.

Every absolutely almost simple algebraic group scheme of adjoint type  ${}^2A_n$  over  $F$  is isomorphic to  $\mathbf{PGU}(B, \tau) = \mathbf{Aut}_K(B, \tau)$  for some central simple algebra  $B$  of degree  $n + 1$  over a separable quadratic field extension  $K$  of  $F$  with a  $K/F$ -unitary involution  $\tau$ . As explained below in §2.1, condition (Out 1) holds for the group  $\mathbf{PGU}(B, \tau)$  if and only if  $B$  has exponent at most 2, and condition (Out 3) holds if and only if  $(B, \tau)$  has a descent, i.e.,  $(B, \tau) = (B_0, \tau_0) \otimes_F (K, \iota)$  for some central simple  $F$ -algebra with  $F$ -linear involution  $(B_0, \tau_0)$ . For  $n$  even, Theorem 1.1(1) can be reformulated in a more precise form:

**THEOREM 1.2.** – *Let  $(B, \tau)$  be a central simple algebra with unitary involution. If  $\deg B$  is odd, then conditions (Out 1), (Out 2), and (Out 3) for  $\mathbf{PGU}(B, \tau)$  are equivalent and hold if and only if  $B$  is split.*

The proof is easy: see Corollary 2.4.

Now, assume  $G = \mathbf{PGU}(B, \tau)$  has type  ${}^2A_5$ , i.e.,  $B$  has degree 6. If the exponent of  $B$  is at most 2, then its index is at most 2. Therefore, Theorem 1.1(1) for such groups follows from the following descent theorem for algebras with unitary involution, proved in §4.1:

**THEOREM 1.3.** – *Let  $(B, \tau)$  be a central simple algebra of degree at most 6 and index at most 2, with a  $K/F$ -unitary involution. There exists a central simple algebra with orthogonal involution  $(B_0, \tau_0)$  over  $F$ , of the same index as  $B$ , such that  $(B, \tau) = (B_0, \tau_0) \otimes (K, \iota)$ , where  $\iota$  is the unique nontrivial  $F$ -automorphism of  $K$ .*

It also follows from this theorem that assertion (1) does hold for groups of type  ${}^2A_3$  when the underlying algebra  $B$  has index at most 2; but this does not apply to all groups of type  ${}^2A_3$ , since a degree 4 central simple algebra of exponent 2 can be of index 4. An example of a degree 4 and exponent 2 algebra with unitary involution that does not have a descent will be provided in §3.3.2 below (see Remark 3.14).

As usual for classical groups, we use as a crucial tool their explicit description in terms of algebras with involution or quadratic pair. How conditions (Out 1), (Out 2) and (Out 3) translate into conditions on these algebraic structures is explained in §2. Section 3 studies in more details the  ${}^2D_n$  case. In particular, we introduce our main tool for proving assertion (2) of Theorem 1.1, namely “generic” orthogonal sums of hermitian forms or involutions. In §4, using the same kind of strategy, we prove Theorem 1.3, and complete the proof of Theorem 1.1 by producing examples of outer type  ${}^2A_n$ .

We refer the reader to [6] for definitions and basic facts on central simple algebras, involutions, and quadratic pairs. Recall that if  $\text{char } F \neq 2$ , then for any quadratic pair  $(\sigma, f)$ ,  $\sigma$  is an involution of orthogonal type, and  $f$  is the map defined on the set  $\text{Sym}(A, \sigma)$  of  $\sigma$ -symmetric