

Guy David
Marcel Filoche
David Jerison
Svitlana Mayboroda

**A FREE BOUNDARY PROBLEM FOR THE
LOCALIZATION OF EIGENFUNCTIONS**

ASTÉRIQUE 392

Société Mathématique de France 2017

Astérisque est un périodique de la Société mathématique de France
Numéro 392

Comité de rédaction

Ahmed ABBES	Philippe EYSSIDIEUX
Viviane BALADI	Damien GABORIAU
Laurent BERGER	Michael HARRIS
Philippe BIANE	Fabrice PLANCHON
Hélène ESNAULT	Pierre SCHAPIRA

Éric VASSEROT (dir.)

Diffusion

Maison de la SMF	AMS
B.P. 67	P.O. Box 6248
13274 Marseille CEDEX 9	Providence RI 02940
France	USA
smf@smf.univ-mrs.fr	www.ams.org

Tarifs 2017

Vente au numéro : 45 € (\$ 67)
Abonnement électronique : 500 € (\$ 750)
Abonnement avec supplément papier : 657 €, hors Europe : 699 € (\$ 1049)
Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris CEDEX 05, France
Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96
astsmf@ihp.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2017

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN : print 0303-1179, electronic 2492-5926
ISBN 978-2-85629-863-3

Directeur de la publication : Stéphane SEURET

ASTÉRIQUE 392

**A FREE BOUNDARY PROBLEM FOR THE
LOCALIZATION OF EIGENFUNCTIONS**

**Guy David
Marcel Filoche
David Jerison
Svitlana Mayboroda**

Société Mathématique de France 2017
Publié avec le concours du Centre National de la Recherche Scientifique

G. David

Univ Paris-Sud, Laboratoire de Mathématiques, UMR 8658 Orsay, F-91405; CNRS, Orsay, F-91405; Institut Universitaire de France.

E-mail : Guy.David@math.u-psud.fr

M. Filoche

Physique de la Matière Condensée, École Polytechnique, CNRS, Palaiseau, France; CMLA, ÉNS Cachan, UniverSud, Cachan, France.

E-mail : marcel.filoche@polytechnique.edu

D. Jerison

Massachusetts Institute of Technology, 77 Massachusetts Ave, Cambridge, MA 02139-4307 USA.

E-mail : jerison@math.mit.edu

S. Mayboroda

School of Mathematics, University of Minnesota, 206 Church St SE, Minneapolis, MN 55455 USA.

E-mail : svitlana@math.umn.edu

2010 Mathematics Subject Classification. --- 49Q20, 35B65.

Key words and phrases. --- Free boundary problem, localization of eigenfunctions.

- The first author acknowledges the generous support of the Institut Universitaire de France, and of the ANR (programme blanc GEOMETRYA, ANR-12-BS01-0014).
- The second author was partially supported by the CNRS grant PEPS-PTI Thorgal.
- The third author was partially supported by the NSF grant DMS 1069225, the Bergman Trust, and wishes to thank Université de Paris Sud (Orsay) for a monthlong visit during June–July 2012.
- The fourth author was partially supported by the NSF grants DMS 1220089 (CAREER), DMS 1344235 (INSPIRE), DMR 0212302 (UMN MRSEC Seed grant), and the the Alfred P. Sloan Fellowship. The fourth author would also like to acknowledge the support of the CNRS and hospitality of Université Paris-Sud, Orsay, France, in Oct.–Dec. of 2012 and April of 2013, when parts of this work were completed.

A FREE BOUNDARY PROBLEM FOR THE LOCALIZATION OF EIGENFUNCTIONS

Guy David, Marcel Filoche, David Jerison, Svitlana Mayboroda

Abstract. --- We study a variant of the Alt, Caffarelli, and Friedman free boundary problem, with many phases and a slightly different volume term, which we originally designed to guess the localization of eigenfunctions of a Schrödinger operator in a domain. We prove Lipschitz bounds for the functions and some nondegeneracy and regularity properties for the domains.

Résumé (Un problème de frontière libre pour la localisation de fonctions propres)

On étudie une variante du problème de frontière libre de Alt, Caffarelli et Friedman, avec plusieurs phases et un terme de volume légèrement différent, que l'on a choisie pour deviner la localisation des fonctions propres d'un opérateur de Schrödinger dans un domaine. On démontre des estimations lipschitziennes pour les fonctions associées à un minimiseur et des propriétés de nondégérescence et de régularité pour les frontières libres.

CONTENTS

1. Introduction	1
2. Motivation for our main functional	11
3. Existence of minimizers	17
4. Poincaré inequalities and restriction to spheres	25
5. Minimizers are bounded	31
6. Two favorite competitors	39
7. Hölder-continuity of u inside Ω	45
8. Hölder-continuity of u on the boundary	55
9. The monotonicity formula	63
10. Interior Lipschitz bounds for u	67
11. Global Lipschitz bounds for u when Ω is smooth	81
12. A sufficient condition for $ u $ to be positive	87
13. Sufficient conditions for minimizers to be nontrivial	97
14. A bound on the number of components	101
15. The main non degeneracy condition; good domains	105
16. The boundary of a good region is rectifiable	123
17. Limits of minimizers	133
18. Blow-up Limits are minimizers	151

19. Blow-up limits with two phases	159
20. Blow-up limits with one phase	175
21. Local regularity when all the indices are good	187
22. First variation and the normal derivative	195
Bibliography	201

CHAPTER 1

INTRODUCTION

The initial motivation for this paper was to describe the localization of eigenfunctions for an operator \mathcal{L} on a domain $\Omega \subset \mathbb{R}^n$. Let us assume that $|\Omega|$, the measure of Ω , is finite. The typical operator that we consider is the positive Laplacian $\mathcal{L} = -\Delta$, or a Schrödinger operator $\mathcal{L} = -\Delta + \mathcal{V}$, with \mathcal{V} bounded and nonnegative.

In [20], a pointwise estimate for eigenfunctions for \mathcal{L} is found, which bounds them in terms of a single function w_0 , namely the solution of $\mathcal{L}w_0 = 1$ on Ω , with the Dirichlet condition $w_0 = 0$ on $\mathbb{R}^n \setminus \Omega$. Our first goal is to derive an automatic way, using w_0 , to find subdomains W_j , $1 \leq j \leq N$, of Ω , where the eigenfunctions of \mathcal{L} are more likely to be supported. The work in [20] indicates that, roughly speaking, one seeks a collection of disjoint $W_j \subset \Omega$, $1 \leq j \leq N$, such that w_0 is small on the boundaries of the W_j , and it is natural to try to measure “smallness” in terms of the operator \mathcal{L} itself. Even though many handmade or numerical decompositions of Ω based on w_0 seem to give very good predictions of the localization of eigenfunctions, we would like to have a more systematic way to realize the decomposition.

The functional described below is intended to give such a good partition of Ω into subdomains, and it turns out to be an interesting variant of functionals introduced by Alt and Caffarelli [1], and studied by many others. In the present paper, we shall mainly study the theoretical properties of our functional (existence and regularity of the minimizers and regularity of the corresponding free boundaries).

Let us now describe the main free boundary problem that we shall study here; the relation with w_0 and our original localization problem will be explained more in Chapter 2.

We are given a domain $\Omega \subset \mathbb{R}^n$, and (for instance) an operator $\mathcal{L} = -\Delta + \mathcal{V}$; assumptions on the potential \mathcal{V} , or other functions associated to a similar problem, will come later. We are also given an integer $N \geq 1$, and we want to cut Ω into subregions W_i , $1 \leq i \leq N$, according to the geometry associated to \mathcal{L} . For this, we want to define and minimize a functional J . But let us first define the set of admissible pairs (\mathbf{u}, W) for which $J(\mathbf{u}, W)$ is defined.

DEFINITION 1.1. — Given the open set $\Omega \subset \mathbb{R}^n$ and the integer $N \geq 1$, we denote by $\mathcal{F} = \mathcal{F}(\Omega)$ the set of admissible pairs (\mathbf{u}, \mathbf{W}) , where $\mathbf{W} = (W_i)_{1 \leq i \leq N}$ is a N -uple of pair-wise disjoint Borel-measurable sets $W_i \subset \Omega$, and $\mathbf{u} = (u_i)_{1 \leq i \leq N}$ is a N -uple of real-valued functions u_i , such that

$$(1.1) \quad u_i \in W^{1,2}(\mathbb{R}^n)$$

and

$$(1.2) \quad u_i(x) = 0 \text{ for almost every } x \in \mathbb{R}^n \setminus W_i.$$

Here $W^{1,p}(\mathbb{R}^n)$, $1 \leq p < +\infty$, denotes the set of functions $f \in L^p_{\text{loc}}(\mathbb{R}^n)$ whose derivative, computed in the distribution sense, lies in $L^p(\mathbb{R}^n)$. We chose a definition for which we do not need to assume any regularity for the sets W_i , nor to give a precise meaning to the Sobolev space $W^{1,2}_0(W_i)$, but under mild assumptions, the functions u_i associated to minimizers will be continuous, and we will be able to take

$$W_i = \{x \in \Omega; u_i(x) > 0\}.$$

For the moment we took real-valued functions, but what we will say will systematically apply when some of the functions u_i are required to be nonnegative. In addition, some of our results will only work under this constraint (that $u_i \geq 0$).

Our functional J will have three main terms. The first one is the energy

$$(1.3) \quad E(\mathbf{u}) = \sum_{i=1}^N \int |\nabla u_i(x)|^2 dx,$$

where we denoted by ∇u_i the distributional gradient of u_i , which is an L^2 function. It does not matter whether we integrate on \mathbb{R}^n , Ω , or W_i , because one can check that if $u_i \in W^{1,2}(\mathbb{R}^n)$ vanishes almost everywhere on $\mathbb{R}^n \setminus W_i$, then $\nabla u_i = 0$ almost everywhere on $\mathbb{R}^n \setminus W_i$. Indeed, the Rademacher-Calderón theorem says that u_i is differentiable (i.e., has a derivative) at almost every point of \mathbb{R}^n , with a differential that coincides almost everywhere with the (density of the) distribution Du_i . See for instance Theorem 6.15 on page 47 of [23]. It is then easy to check that $Du_i(x) = 0$ when x is a point of Lebesgue differentiability of $\mathbb{R}^n \setminus W_i$, hence, for almost every $\mathbb{R}^n \setminus W_i$.

The second term of our functional will be

$$(1.4) \quad M(\mathbf{u}) = \sum_{i=1}^N \int [u_i(x)^2 f_i(x) - u_i(x) g_i(x)] dx,$$

where the f_i and the g_i , $1 \leq i \leq N$, are given functions on Ω that we may choose, depending on our problem. We are slightly abusing notation here, because $M(\mathbf{u})$ also depends on the W_i through the choice of functions we integrate against u_i^2 or u_i , at least if the f_i and g_i depend on i . The convergence of the integrals in (1.4) will follow from our assumptions on the f_i and the g_i . For our initial localization problem, all the f_i will be equal to the potential \mathcal{V} , and $g_i = 2$ for all i .