

380

ASTÉRIQUE

2016

SÉMINAIRE BOURBAKI  
VOLUME 2014/2015  
EXPOSÉ N° 1100

Gil KALAI

*Designs exist!*

**SOCIÉTÉ MATHÉMATIQUE DE FRANCE**

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

*Comité de rédaction*

Ahmed ABBES  
Viviane BALADI  
Laurent BERGER  
Gérard BESSON  
Philippe BIANE  
Hélène ESNAULT

Damien GABORIAU  
Michael HARRIS  
Fabrice PLANCHON  
Pierre SCHAPIRA  
Bertrand TOËN

Éric VASSEROT (dir.)

*Diffusion*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 9  
France  
smf@smf.univ-mrs.fr

Hindustan Book Agency  
O-131, The Shopping Mall  
Arjun Marg, DLF Phase 1  
Gurgaon 122002, Haryana  
Inde

AMS  
P.O. Box 6248  
Providence RI 02940  
USA  
www.ams.org

*Tarifs*

*Vente au numéro* : 75 € (\$ 112)

*Abonnement* Europe : 650 €, hors Europe : 689 € (\$ 1 033)

Des conditions spéciales sont accordées aux membres de la SMF.

*Secrétariat : Nathalie Christiaën*

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2016

*Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.*

ISSN 0303-1179

ISBN 978-85629-836-7

Directeur de la publication : Marc PEIGNÉ

---



**DESIGNS EXIST!**  
**[after Peter Keevash]**

by Gil KALAI

**INTRODUCTION**

A set  $S$  of  $q$ -subsets of an  $n$ -set  $X$  is a *design* with parameters  $(n, q, r, \lambda)$  if every  $r$ -subset of  $X$  belongs to exactly  $\lambda$  elements of  $S$ . (The elements of  $S$  are called *blocks* and designs are also referred to as *block designs*.) There are some necessary *divisibility conditions* for the existence of such a design, namely that

$$(1) \quad \binom{q-i}{r-i} \text{ divides } \lambda \binom{n-i}{r-i}, \quad 0 \leq i \leq r-1.$$

To see that the divisibility conditions are necessary, fix any  $i$ -subset  $I$  of  $X$  and consider the sets in  $S$  that contain  $I$ .

The following result was conjectured in the 19th century and was recently proved by Peter Keevash.

**THEOREM 0.1 ([29]).** — *For fixed  $q, r$ , and  $\lambda$ , there exist  $n_0(q, r, \lambda)$  such that if  $n > n_0(q, r, \lambda)$  satisfies the divisibility conditions (1) then a design with parameters  $(n, q, r, \lambda)$  exists.*

In other words, for fixed  $q, r$ , and  $\lambda$ , the divisibility conditions are sufficient apart from a finite number of exceptional values of  $n$ .

A case of special interest is when  $\lambda = 1$ . A design of parameters  $(n, q, r, 1)$  is called a *Steiner system* of parameters  $(n, q, r)$ . The question if Steiner systems of given parameters exist goes back to works of Plücker, Kirkman, and Steiner. Until Keevash's result not a single Steiner system for  $r > 5$  was known to exist.

The presentation of Keevash's work in this paper is based on Keevash's original paper [29], his Jerusalem videotaped lectures [27], and his subsequent paper [28]. It is also based on lecture notes and personal explanations by Jeff Kahn.

## 1. REGULARITY, SYMMETRY AND RANDOMNESS

### 1.1. Between regularity and symmetry

An object is “regular” if it looks locally the same (for a certain notion of “locality,”), and it is “symmetric” if it admits a transitive group action (on its “local” pieces). The interplay between regularity and symmetry is of interest in several parts of mathematics. For example, a regular graph is a graph where every vertex is adjacent to the same number of neighbors, and every graph whose group of automorphisms act transitively on its vertices is regular. Regular graphs need not be symmetric (most of them have no non-trivial automorphisms), but there are still various connections between general regular graphs and symmetry. Another example: manifolds are regular objects and Lie groups are symmetric types of manifolds; also here there are rich connections between regularity and symmetry.

Designs are regular objects. (The local pieces can be regarded as the  $r$ -subsets of the ground set.) You can get them from groups acting transitively on  $r$ -sets.

**PROPOSITION 1.1.** — *Let  $\Gamma$  be a  $r$ -transitive permutation group. Then the orbit of a set of size  $q$  is a block design so that every set of size  $r$  belongs to the same number of blocks.*

However, it follows from the classification of finite simple groups that

**THEOREM 1.2.** — *Let  $\Gamma$  be a  $r$ -transitive permutation group,  $r > 5$ , then  $G$  is  $A_n$  or  $S_n$ .*

### 1.2. The probabilistic method and quasi-randomness

The proof of the existence of designs is probabilistic. In order to prove the existence of objects of some kind satisfying a property  $\mathbf{P}$ , one proves that for a suitable probability distribution on all objects of this kind there is a positive probability for property  $\mathbf{P}$  to hold. The probabilistic method is of central importance in combinatorics (and other areas) [1]. Keevash defines a complicated combinatorial process with random ingredients for building a design, and shows that with positive probability it leads to the desired construction.

Quasi-randomness refers to deterministic properties of mathematical structures which allow them to behave (for certain restricted purposes) “as if they were random.”<sup>(1)</sup> Quasi random properties of primes are, of course, of much importance. In graph theory, a sequence of graphs  $G_n$  (where  $G_n$  has  $n$  vertices) is quasi-random

---

<sup>(1)</sup> There are even cases that quasi-randomness of some kind can be attributed to arbitrary structures as is the case in Szemerédi regularity lemma which describes quasi-random structure on arbitrary graphs.