

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

THE SCHUR MULTIPLIER OF FINITE SYMPLECTIC GROUPS

Louis Funar & Wolfgang Pitsch

Tome 148
Fascicule 3

2020

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

pages 515-527

Le *Bulletin de la Société Mathématique de France* est un périodique trimestriel
de la Société Mathématique de France.

Fascicule 3, tome 148, septembre 2020

Comité de rédaction

Christine BACHOC	Laurent MANIVEL
Yann BUGEAUD	Julien MARCHÉ
Jean-François DAT	Kieran O'GRADY
Clothilde FERMANIAN	Emmanuel RUSS
Pascal HUBERT	Christophe SABOT

Marc HERZLICH (Dir.)

Diffusion

Maison de la SMF	AMS
Case 916 - Luminy	P.O. Box 6248
13288 Marseille Cedex 9	Providence RI 02940
France	USA
commandes@smf.emath.fr	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Bulletin de la SMF

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 1 44 27 67 99 • Fax : (33) 1 40 46 90 96

bulletin@smf.emath.fr • smf.emath.fr

© Société Mathématique de France 2020

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Fabien DURAND

THE SCHUR MULTIPLIER OF FINITE SYMPLECTIC GROUPS

BY LOUIS FUNAR & WOLFGANG PITSCH

ABSTRACT. — We show that the Schur multiplier of $Sp(2g, \mathbb{Z}/D\mathbb{Z})$ is $\mathbb{Z}/2\mathbb{Z}$, when D is divisible by 4.

RÉSUMÉ (*Multiplicateur de Schur des groupes symplectiques finis*). — Nous montrons que le multiplicateur de Schur de $Sp(2g, \mathbb{Z}/D\mathbb{Z})$ est $\mathbb{Z}/2\mathbb{Z}$ quand D est divisible par 4.

1. Introduction and statements

Let $g \geq 1$ be an integer and denote by $Sp(2g, \mathbb{Z})$ the symplectic group of $2g \times 2g$ matrices with integer coefficients. Deligne's nonresidual finiteness theorem from [5] states that the *universal central extension* $\widetilde{Sp(2g, \mathbb{Z})}$ is not residually finite, since the image of its center under any homomorphism into a finite group has an order of at most 2 when $g \geq 3$. Our first motivation was to understand this result and give a sharp statement, namely to decide whether the

Texte reçu le 30 octobre 2013, modifié le 5 novembre 2019, accepté le 3 mars 2020.

LOUIS FUNAR, Institut Fourier BP 74, UMR 5582, Laboratoire de Mathématiques, Université Grenoble Alpes, CS 40700, 38058 Grenoble cedex 9, France • *E-mail* : louis.funar@univ-grenoble-alpes.fr

WOLFGANG PITSCH, Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra (Cerdanyola del Vallès), Spain • *E-mail* : pitsch@mat.uab.es

Mathematical subject classification (2010). — 57M50, 55N25, 19C09, 20F38.

Key words and phrases. — Symplectic groups, Group homology, Mapping class groups, Central extension, Residually finite group.

The first author was supported by the ANR 2011 BS 01 020 01 ModGroup and the second author by the FEDER/MEC grant MTM2016-80439-P.

image of the central \mathbb{Z} might be of order 2. Since these symplectic groups have the congruence subgroup property, this boils down to understanding the second homology of symplectic groups with coefficients in finite cyclic groups. In the sequel, for simplicity and unless otherwise explicitly stated, all (co)homology groups will be understood to be with trivial integer coefficients. An old theorem of Stein (see [14], Thm. 2.13 and Prop. 3.3.a) is that $H_2(Sp(2g, \mathbb{Z}/D\mathbb{Z})) = 0$, when D is not divisible by 4. The case $D \equiv 0 \pmod{4}$ has remained open since then; this is explicitly mentioned, for instance, in [13] (Remarks after Thm. 3.8). Our main result settles this case.

THEOREM 1.1. — *The second homology group of finite principal congruence quotients of $Sp(2g, \mathbb{Z})$, $g \geq 3$ is*

$$H_2(Sp(2g, \mathbb{Z}/D\mathbb{Z})) = \mathbb{Z}/2\mathbb{Z}, \text{ if } D \equiv 0 \pmod{4}.$$

In comparison, recall that Beyl (see [2]) showed that $H_2(SL(2, \mathbb{Z}/D\mathbb{Z})) = \mathbb{Z}/2\mathbb{Z}$, for $D \equiv 0 \pmod{4}$, and Dennis and Stein proved using K-theoretic methods that for $n \geq 3$, we have $H_2(SL(n, \mathbb{Z}/D\mathbb{Z})) = \mathbb{Z}/2\mathbb{Z}$, for $D \equiv 0 \pmod{4}$, while $H_2(SL(n, \mathbb{Z}/D\mathbb{Z})) = 0$, for $D \not\equiv 0 \pmod{4}$ (see [6], Cor. 10.2 and [11], Section 12).

Our proof also relies on Deligne's nonresidual finiteness theorem from [5] and deep results of Putman in [13], and shows that we can detect this $\mathbb{Z}/2\mathbb{Z}$ factor on $H_2(Sp(2g, \mathbb{Z}/32\mathbb{Z}))$ for $g \geq 4$, providing an explicit extension that detects this homology class.

2. Preliminaries

2.1. Residual finiteness of universal central extensions. — In this section, we collect results about universal central extensions of perfect groups, for the sake of completeness of our arguments. Every perfect group Γ has a universal central extension $\tilde{\Gamma}$; the kernel of the canonical projection map $\tilde{\Gamma} \rightarrow \Gamma$ contains the center $Z(\tilde{\Gamma})$ of $\tilde{\Gamma}$ and is canonically isomorphic to the second integral homology group $H_2(\Gamma)$. We will recall now how the residual finiteness problem for the universal central $\tilde{\Gamma}$ of a perfect and residually finite group Γ translates into an homological problem about $H_2(\Gamma)$. We start with a classical result for maps between universal central extensions of perfect groups.

LEMMA 2.1. — *Let Γ and F be perfect groups, $\tilde{\Gamma}$ and \tilde{F} be their universal central extensions and $p : \Gamma \rightarrow F$ be a group homomorphism. Then, there exists a unique homomorphism $\tilde{p} : \tilde{\Gamma} \rightarrow \tilde{F}$ lifting p such that the following*

diagram is commutative:

$$\begin{array}{ccccccc}
 1 & \rightarrow & H_2(\Gamma) & \rightarrow & \tilde{\Gamma} & \rightarrow & \Gamma & \rightarrow & 1 \\
 & & p_* \downarrow & & \tilde{p} \downarrow & & \downarrow p & & \\
 1 & \rightarrow & H_2(F) & \rightarrow & \tilde{F} & \rightarrow & F & \rightarrow & 1
 \end{array}$$

For a proof we refer the interested reader to ([10], Chap. VIII) or ([4], Chap. IV, Ex. 1, 7). If Γ is a perfect residually finite group, to prove that its universal central extension $\tilde{\Gamma}$ is also residually finite we only have to find enough finite quotients of $\tilde{\Gamma}$ to detect the elements in its center $H_2(\Gamma)$. The following lemma analyzes the situation.

LEMMA 2.2. — *Let Γ be a perfect group and denote by $\tilde{\Gamma}$ its universal central extension.*

1. *Let H be a finite index normal subgroup $H \subset \Gamma$ such that the image of $H_2(H)$ into $H_2(\Gamma)$ contains the subgroup $dH_2(\Gamma)$, for some $d \in \mathbb{Z}$. Let $F = \Gamma/H$ be the corresponding finite quotient of Γ and $p : \Gamma \rightarrow F$ the quotient map. Then, $d \cdot p_*(H_2(\Gamma)) = 0$, where $p_* : H_2(\Gamma) \rightarrow H_2(F)$ is the homomorphism induced by p . In particular, if $p_* : H_2(\Gamma) \rightarrow H_2(F)$ is surjective, then $d \cdot H_2(F) = 0$.*
2. *Assume that F is a finite quotient of Γ satisfying $d \cdot p_*(H_2(\Gamma)) = 0$. Let \tilde{F} denote the universal central extension of F . Then, the homomorphism $p : \Gamma \rightarrow F$ has a unique lift $\tilde{p} : \tilde{\Gamma} \rightarrow \tilde{F}$, and the kernel of \tilde{p} contains $d \cdot H_2(\Gamma)$.*

Observe that in point 2. of Lemma 2.2 the group F being finite, $H_2(F)$ is also finite, and, hence, one can take $d = |H_2(F)|$.

Proof. — The image of H into F is trivial, and, thus, the image of $H_2(H)$ into $H_2(F)$ is trivial. This implies that $p_*(d \cdot H_2(\Gamma)) = 0$, which proves the first part of the lemma.

Further, by Lemma 2.1 there exists a unique lift $\tilde{p} : \tilde{\Gamma} \rightarrow \tilde{F}$. If $d \cdot p_*(H_2(\Gamma)) = 0$, then Lemma 2.1 yields $d \cdot \tilde{p}(c) = d \cdot p_*(c) = 0$, for any $c \in H_2(\Gamma)$. This settles the second part of the lemma. □

REMARK 2.3. — It might be possible that we have $d' \cdot p_*(H_2(\Gamma)) = 0$ for some proper divisor d' of d , so the first part of Lemma 2.2 can only give an upper bound of the orders of the image of the second cohomology. In order to find lower bounds we need additional information concerning the finite quotients F .

LEMMA 2.4. — *Let Γ be a perfect group, $\tilde{\Gamma}$ its universal central extension, $p : \Gamma \rightarrow F$ a surjective homomorphism onto a finite group F , and $\hat{p} : \tilde{\Gamma} \rightarrow G$ be some lift of p to a central extension G of F by some finite abelian group C . Assume that the image of $H_2(\Gamma) \subset \tilde{\Gamma}$ in G by \hat{p} contains an element of order q . Then there exists an element of $p_*(H_2(\Gamma)) \subset H_2(F)$ of order q .*