

COINCIDENCE FOR SUBSTITUTIONS OF PISOT TYPE

BY MARCY BARGE & BEVERLY DIAMOND

ABSTRACT. — Let φ be a substitution of Pisot type on the alphabet $\mathcal{A} = \{1, 2, \dots, d\}$; φ satisfies the *strong coincidence condition* if for every $i, j \in \mathcal{A}$, there are integers k, n such that $\varphi^n(i)$ and $\varphi^n(j)$ have the same k -th letter, and the prefixes of length $k - 1$ of $\varphi^n(i)$ and $\varphi^n(j)$ have the same image under the abelianization map. We prove that the strong coincidence condition is satisfied if $d = 2$ and provide a partial result for $d \geq 2$.

RÉSUMÉ (*Coincidence pour les substitutions de type Pisot*). — Soit φ une substitution de type Pisot sur un alphabet $\mathcal{A} = \{1, 2, \dots, d\}$; on dit que φ satisfait la *condition de coïncidence forte* si pour tout $i, j \in \mathcal{A}$, il existe des entiers k, n tels que $\varphi^n(i)$ et $\varphi^n(j)$ aient la même k -ième lettre et les préfixes de longueur $k - 1$ de $\varphi^n(i)$ et $\varphi^n(j)$ aient la même image par l'application d'abélianisation. Nous montrons que la condition de coïncidence forte est satisfaite pour $d = 2$ et nous donnons un résultat partiel pour $d \geq 2$.

A substitution φ on a finite alphabet $\mathcal{A} = \{1, 2, \dots, d\}$ satisfies the *strong coincidence condition* if for every $i, j \in \mathcal{A}$, there are integers k, n such that
(i) $\varphi^n(i)$ and $\varphi^n(j)$ have the same k -th letter and

Texte reçu le 21 novembre 2001, accepté le 14 mai 2002

MARCY BARGE, Department of Mathematics, Montana State University, Bozeman MT 59717 (USA) • *E-mail* : barge@math.montana.edu

BEVERLY DIAMOND, Department of Mathematics, College of Charleston, Charleston SC 29424 (USA) • *E-mail* : diamondb@cofc.edu

2000 Mathematics Subject Classification. — 37B10.

Key words and phrases. — Substitution, dynamical system, Pisot, coincidence conjecture, pure discrete spectrum.

The second author was partially supported by a research incentive grant from the South Carolina Commission on Higher Education.

(ii) the prefixes of length $k - 1$ of $\varphi^n(i)$ and $\varphi^n(j)$ have the same image under the abelianization map (that is, these prefixes contain the same number of occurrences of each $i \in \mathcal{A}$).

This condition (without the requirement on prefixes) was first introduced for substitutions of constant length by F.M. Dekking [5], who proved that for such substitutions, the condition is satisfied if and only if the associated substitutive dynamical system has pure discrete spectrum (equivalently, the substitutive system is metrically isomorphic with translation on a compact abelian group). In its current form, the strong coincidence condition is due to Arnoux and Ito. The equivalence of the strong coincidence condition and pure discrete spectrum also holds for substitutions of nonconstant length of Pisot type on two letters (unpublished work of B. Host, [8], [9], [6], and [7]). According to [4], Host also proved that if φ is a Pisot type substitution on two letters which is unimodular (its transition matrix has determinant $+1$ or -1) and which satisfies the strong coincidence condition, then the substitutive system associated with φ is metrically isomorphic with both an interval exchange and a one-dimensional toral rotation. Arnoux and Ito [1] have recently shown that every unimodular Pisot substitution on $d \geq 2$ letters satisfying the strong coincidence condition is metrically isomorphic with an exchange of domains in \mathbb{R}^{d-1} and is semi-conjugate to a rotation on the $d-1$ -torus (see also [11]). At present, an additional condition is needed to improve the semi-conjugacy to metric isomorphism. For examples of nonunimodular Pisot substitutive systems that are metrically isomorphic with translations on p -adic groups and p -adic solenoids, see [10].

No examples of Pisot type substitutions which do not satisfy the strong coincidence condition are known, and the statement that all such substitutions satisfy this condition is known as the *Strong Coincidence Conjecture*. In this paper, we prove that the Strong Coincidence Conjecture holds for Pisot substitutions on two letters and provide a partial result for substitutions on d letters where $d > 2$.

THEOREM 1. — *Let φ be a Pisot substitution on an alphabet $\mathcal{A} = \{1, 2, \dots, d\}$. There are distinct letters $i, j \in \mathcal{A}$ for which there are integers k, n such that $\varphi^n(i)$ and $\varphi^n(j)$ have the same k -th letter, and the prefixes of length $k - 1$ of $\varphi^n(i)$ and $\varphi^n(j)$ have the same image under the abelianization map.*

We introduce terminology necessary for the proof of the theorem.

For $j = 1$ to d , I_j will denote the interval of length 1 in \mathbb{R}^d given by

$$I_j = \{0\} \times \{0\} \times \cdots \times [0, 1] \times \cdots \times \{0\},$$

where $[0, 1]$ appears in the j -th position.

Similarly, for $1 \leq j \leq d$, \mathbf{e}_j will denote the unit vector $(0, 0, \dots, 1, \dots, 0)$, where 1 appears in the j -th position. If $x \in \mathbb{Z}^d$, the integer lattice in \mathbb{R}^d , the vector from the origin to x is denoted by \mathbf{x} .

For $\mathbf{v} = (v_1, v_2, \dots, v_d)$,

$$I_j + \mathbf{v} = \{v_1\} \times \{v_2\} \times \cdots \times [v_j, v_j + 1] \times \cdots \times \{v_d\}.$$

The collection \mathcal{M} of line segments joining ‘adjacent’ elements of the integer lattice is then

$$\mathcal{M} = \{I_j + \mathbf{x} : 1 \leq j \leq d, x \in \mathbb{Z}^d\}.$$

The term *segment* will refer to an element of \mathcal{M} .

Let $\mathcal{A} = \{1, 2, \dots, d\}$ be a finite alphabet and \mathcal{A}^* the collection of finite non-empty words formed from the alphabet \mathcal{A} .

A *substitution* φ is a map $\varphi : \mathcal{A} \rightarrow \mathcal{A}^*$; φ has an associated *transition matrix* $A_\varphi = A = (a_{ij})_{i,j \in \mathcal{A}}$ in which a_{ij} is the number of occurrences of i in the word $\varphi(j)$. (In what follows, we will also use A to denote the linear transformation on vectors of \mathbb{R}^d and the linear function acting on points of \mathbb{R}^d .)

The substitution φ is *primitive* if there is n so that for each $i, j \in \mathcal{A}$, j appears in $\varphi^n(i)$; φ is *of Pisot type*, or simply *Pisot*, if all eigenvalues of A other than the Perron-Frobenius eigenvalue have modulus strictly between 0 and 1.

If φ is Pisot, then φ is primitive and the (hyperbolic) linear map on \mathbb{R}^d defined by the matrix A has stable space E^s of dimension $d - 1$ and unstable space E^u of dimension 1 spanned by a positive Perron-Frobenius eigenvector \mathbf{v}_u . Also, neither E^s nor E^u contain elements of the integer lattice other than the origin. (See Chapter 1 of [3], for instance.)

Assume for the remainder of this paper that φ is Pisot.

We define an order on \mathbb{Z}^d as follows. Given $P_1, P_2 \in \mathbb{Z}^d$, $P_1 \leq P_2$ if there are $t_1 \leq t_2 \in \mathbb{R}$ so that $P_1 \in E^s + t_1 \mathbf{v}_u$, $P_2 \in E^s + t_2 \mathbf{v}_u$ (we use $E^s + t \mathbf{v}_u$ to denote the set of points x such that $\mathbf{x} = \mathbf{w} + t \mathbf{v}_u$ for some $\mathbf{w} \in E^s$). Note that if $P_1, P_2 \in \mathbb{Z}^d$, and $P_1 \leq P_2$, then $AP_1 \leq AP_2$.

We employ a geometrical formulation of substitution developed by Arnoux, Ito and Sano [2]. The *inflation and substitution map* F_φ is defined on subsets of the set \mathcal{M} of line segments in the following manner. Suppose that

$$\varphi(j) = a(j, 1)a(j, 2) \cdots a(j, n(j)), \quad \text{for } 1 \leq j \leq d.$$

Define

$$\begin{aligned} \mathbf{v}_\varphi(j, 0) &= \mathbf{0}, \\ \mathbf{v}_\varphi(j, i) &= \mathbf{v}_\varphi(j, i-1) + \mathbf{e}_{a(j, i)}, \quad 1 \leq i \leq n(j). \end{aligned}$$

Then

$$\begin{aligned} F_\varphi(I_j) &= \{I_{a(j, i)} + \mathbf{v}_\varphi(j, i-1) : 1 \leq i \leq n(j)\}, \\ F_\varphi(I_j + \mathbf{v}) &= F_\varphi(I_j) + A\mathbf{v} = \{s + A\mathbf{v} : s \in F_\varphi(I_j)\}. \end{aligned}$$

Finally, if $M \subseteq \mathcal{M}$,

$$F_\varphi(M) = \{F(s) : s \in M\}.$$

(Strictly speaking, in the above, we should write $F_\varphi(\{I_j\})$ rather than $F_\varphi(I_j)$, etc., but we abuse notation and use the latter when convenient.) Note that

$$F_{\varphi^n} = F_\varphi^n.$$

Suppose that

$$S = \{s_i = I_{s(i)} + \mathbf{v}_i : 1 \leq i \leq k, 1 \leq s(i) \leq d\}$$

is a non-empty finite collection of segments, i.e., a subset of \mathcal{M} . The set S is a *strand* if for each $1 \leq i \leq k-1$,

$$\mathbf{v}_{i+1} = \mathbf{v}_i + \mathbf{e}_{s(i)}.$$

The *word associated with S* is $w_S = s(1)s(2) \cdots s(k)$. For such a word w ,

$$\mathbf{l}(w) = (w_1, w_2, \dots, w_d)$$

where for each $i \in \mathcal{A}$, $w_i = \#$ of occurrences of i in W . The *displacement vector for S* is then $\mathbf{l}(S) = \mathbf{l}(w_S)$ and the *length of S* is $|S| = |\mathbf{l}(S)|$, where

$$|(v_1, v_2, \dots, v_d)| = \sum_{1 \leq j \leq d} |v_j|.$$

The endpoints of the segments belonging to S will be called *vertices of S* . A strand S has an *initial vertex* P and a *terminal vertex* Q with the property that $P \leq R \leq Q$ for all vertices R of S ; we say that S is a strand from P to Q . The segments containing P and Q are the *initial* and *terminal* segments of S , respectively. Note that if S is a strand from P to Q , then $F_\varphi(S)$ is a strand from AP to AQ . Two strands S_1 and S_2 are *coincident* if $S_1 \cap S_2 \neq \emptyset$ (that is, they share a segment), and *eventually coincident* if $F_\varphi^m(S_1) \cap F_\varphi^m(S_2) \neq \emptyset$ for some $m \in \mathbb{N}$.

The coincidence conjecture can then be stated as follows:

COINCIDENCE CONJECTURE. — For $1 \leq j, r \leq d$, I_j and I_r are eventually coincident.

We prove the following:

THEOREM 1. — There are $1 \leq j \neq r \leq d$ so that I_j and I_r are eventually coincident.

In the following, $N_B(E^u)$ will denote $\{x \in \mathbb{R}^d : \text{dist}(x, E^u) < B\}$ where $\text{dist}(x, y) = |\mathbf{x} - \mathbf{y}|$, the usual Euclidean distance between x and y .

LEMMA 2. — There are $B > 0$ and $n \in \mathbb{N}$ so that

- (i) if a segment s lies in $N_B(E^u)$, then the strand $F_\varphi^n(s)$ lies in $N_B(E^u)$ and
- (ii) for any segment s , there is $k \in \mathbb{N}$ so that $F_\varphi^{nk}(s)$ lies in $N_B(E^u)$.

Proof. — The fact that φ is Pisot implies that there are λ , $0 < \lambda < 1$, and a constant C so that $|A^n \mathbf{v}| < C\lambda^n |\mathbf{v}|$ for all $\mathbf{v} \in E^s$. It follows that there are n and λ' with $\lambda < \lambda' < 1$ so that $A^n(N_B(E^u)) \subseteq N_{\lambda'B}(E^u)$ for all $B > 0$.

Now suppose that s is any segment lying in $N_B(E^u)$. The initial point of $F_\varphi^n(s)$ lies in $N_{\lambda'B}(E^u)$ and the entire strand $F_\varphi^n(s)$ lies in $N_{\lambda'B+m}(E^u)$, where m is the maximum of the lengths of $\varphi^n(i)$ for $i \in \mathcal{A}$. Choose B large enough so that $\lambda'B + m < \lambda''B$ for some λ'' , $0 < \lambda'' < 1$. For such B and any segment s in $N_B(E^u)$, $F_\varphi^n(s)$ lies in $N_B(E^u)$.

If s is a segment not in $N_B(E^u)$, let B' be large enough so that s lies in $N_{B'}(E^u)$. If $k \geq 1$ is such that $(\lambda'')^k B' \leq B$, then $F_\varphi^{kn}(s)$ lies in $N_B(E^u)$. \square

Since $F_{\varphi^n} = F_\varphi^n$, I_j and I_r are eventually coincident under F_φ if and only if they are eventually coincident under F_{φ^n} . Thus we assume for the remainder of this paper that the n of Lemma 2 is $n = 1$.

A *configuration of segments* is a collection \mathcal{C} of segments with the property that $E^s + \mathbf{v}$ intersects each element of \mathcal{C} in an interior point for some \mathbf{v} . A *configuration of strands* is a collection \mathcal{C} of strands with the property that both the collections of initial segments and final segments form configurations of segments. The *size* of a configuration \mathcal{C} of strands is the number of strands in \mathcal{C} . If P' is the largest of the initial vertices of strands of \mathcal{C} and Q' is the smallest of the terminal vertices of strands of \mathcal{C} , we say that \mathcal{C} is a configuration of strands *from P' to Q'* , or that \mathcal{C} *extends from P' to Q'* , with *length* equal to $|Q' - P'|$.

Given a configuration \mathcal{C} of strands from P' to Q' , there is a unique configuration of strands $\mathcal{C}^{(1)} \subseteq F_\varphi(\mathcal{C})$ with the following properties:

- (i) $\mathcal{C}^{(1)}$ has the same size as \mathcal{C} ;
- (ii) $\mathcal{C}^{(1)}$ extends from AP' to AQ' ; and
- (iii) for each strand $S' \in \mathcal{C}^{(1)}$, there is a strand $S \in \mathcal{C}$ with $S' \subseteq F_\varphi(S)$.

We call $\mathcal{C}^{(1)}$ the first iterate of \mathcal{C} ; higher iterates are defined by

$$\mathcal{C}^{(2)} = (\mathcal{C}^{(1)})^{(1)}, \text{ etc.}$$

A configuration \mathcal{C} of strands is *not eventually coincident* if each distinct pair of strands of \mathcal{C} is not eventually coincident.

The following is a consequence of the definitions and Lemma 2.

LEMMA 3. — *Suppose that \mathcal{C} is a not eventually coincident configuration of strands of size m . The iterates $\mathcal{C}^{(k)}$ of \mathcal{C} satisfy the following:*

- (i) $\mathcal{C}^{(k)}$ is a not eventually coincident configuration of strands;
- (ii) $\text{length}(\mathcal{C}^{(k)}) \rightarrow \infty$ as $k \rightarrow \infty$; and
- (iii) there is a positive integer K so that $\mathcal{C}^{(k)}$ lies in $N_B(E^u)$ for all $k \geq K$.

LEMMA 4. — *There is an integer M with the following properties:*