

ON COVERINGS OF SIMPLE ABELIAN VARIETIES

BY OLIVIER DEBARRE

ABSTRACT. — To any finite covering $f : Y \rightarrow X$ of degree d between smooth complex projective manifolds, one associates a vector bundle E_f of rank $d - 1$ on X whose total space contains Y . It is known that E_f is ample when X is a projective space ([9]), a Grassmannian ([11]), or a Lagrangian Grassmannian ([7]). We show an analogous result when X is a simple abelian variety and f does not factor through any nontrivial isogeny $X' \rightarrow X$. This result is obtained by showing that E_f is M -regular in the sense of Pareschi-Popa, and that any M -regular sheaf is ample.

RÉSUMÉ (*Sur les revêtements des variétés abéliennes simples*). — On associe à tout revêtement fini $f : Y \rightarrow X$ de degré d entre variétés projectives lisses complexes un fibré vectoriel E_f de rang $d - 1$ sur X dont l'espace total contient Y . On sait que E_f est ample lorsque X est un espace projectif ([9]), une grassmannienne ([11]) ou une grassmannienne lagrangienne ([7]). Nous montrons un résultat analogue lorsque X est une variété abélienne simple et que f ne se factorise par aucune isogénie non triviale $X' \rightarrow X$. Ce résultat est obtenu en montrant que E_f est M -régulier au sens de Pareschi-Popa, puis que tout faisceau M -régulier est ample.

Texte reçu le 1er octobre 2004, accepté le 2 février 2005.

OLIVIER DEBARRE, UFR de Mathématique et d'Informatique, 7 rue René Descartes, 67084 Strasbourg Cedex (France). • *E-mail* : debarre@math.u-strasbg.fr

2000 Mathematics Subject Classification. — 14E20, 14J60, 14K02, 14K05, 14K12.

Key words and phrases. — Abelian variety, vector bundle, ample sheaf, M -regular sheaf, continuously generated sheaf, Barth-Lefschetz Theorem, Mukai transform.

1. Introduction

We work over the complex numbers. Let $f : Y \rightarrow X$ be a finite surjective morphism of degree d between smooth projective varieties of the same dimension n . The morphism f is flat, hence the sheaf $f_*\mathcal{O}_Y$ is locally free. We may define a locally free sheaf E_f of rank $d - 1$ on X as the dual of the kernel of the trace map $\mathrm{Tr}_{Y/X} : f_*\mathcal{O}_Y \rightarrow \mathcal{O}_X$, so that

$$f_*\mathcal{O}_Y = \mathcal{O}_X \oplus E_f^*$$

By duality for a finite flat morphism, we have

$$f_*\omega_{Y/X} = \mathcal{O}_X \oplus E_f$$

Our aim is to prove the following statement conjectured in [1].

THEOREM 1.1. — *Let X be a simple abelian variety, let Y be a smooth connected projective variety, and let $f : Y \rightarrow X$ be a finite cover. If f does not factor through any nontrivial isogeny $X' \rightarrow X$, the vector bundle E_f is ample.*

For a more general statement, see Theorem 4.1. See also the remarks at the end of this article for more comments. Even if X is not simple, the vector bundle E_f is known to be nef (see [14, Theorem 1.17], [10, Example 6.3.59]) and its restriction to a general complete intersection curve in X to be ample (see [6, Lemma 2.7]).

The ampleness of E_f has a number of consequences, as explained in [10, Example 6.3.56]. In our case, one new statement beyond the Fulton-Hansen-type results already obtained in [1] is the following: under the hypotheses of the theorem, the induced morphism

$$H^i(f, \mathbb{C}) : H^i(X, \mathbb{C}) \longrightarrow H^i(Y, \mathbb{C})$$

is bijective for $i \leq n - d + 1$ (see [10, Theorem 7.1.16]).

When moreover $d \leq n$, the morphism $\pi_1(f) : \pi_1(Y) \rightarrow \pi_1(X)$ is bijective.⁽¹⁾ In particular, the group $H_1(Y, \mathbb{Z})$ is isomorphic to $H_1(X, \mathbb{Z})$, hence is torsion-free, and so is $H^2(Y, \mathbb{Z})$ by the universal coefficient theorem.

When $d \leq n - 1$, the morphism $H^2(f, \mathbb{Z}) : H^2(X, \mathbb{Z}) \rightarrow H^2(Y, \mathbb{Z})$ is injective with finite cokernel, hence so is $\mathrm{Pic}(f) : \mathrm{Pic}(X) \rightarrow \mathrm{Pic}(Y)$. It seems likely that those two maps are bijective.

The proof is a simple application of the results of [13] about global generation of sheaves on an abelian variety. More precisely, it is based on the remark that any M -regular sheaf (§ 3) on an abelian variety is ample (Corollary 3.2).

⁽¹⁾For algebraic fundamental groups, this is [1, Corollaire 6.2]; for topological fundamental groups, this is [2, Exercice VIII.5], where the hypothesis $d \leq n$ is unfortunately missing.

This work was done while the author was visiting the University of Michigan at Ann Arbor. Many thanks to Bill Fulton and Rob Lazarsfeld for support and many stimulating conversations.

2. Ample sheaves

To any coherent sheaf $\mathcal{F}$ on a scheme X of finite type over $\mathbb{C}$, one associates the X -scheme

$$\mathbf{P}(\mathcal{F}) = \operatorname{Proj} \left(\bigoplus_{m \geq 0} \operatorname{Sym}^m \mathcal{F} \right)$$

and an invertible sheaf $\mathcal{O}_{\mathbf{P}(\mathcal{F})}(1)$ on $\mathbf{P}(\mathcal{F})$. The sheaf $\mathcal{F}$ is said to be ample if $\mathcal{O}_{\mathbf{P}(\mathcal{F})}(1)$ is.

Well-known properties of ampleness for locally free sheaves (see for example [10, Chapter 6]) still hold in this general setting:

- a) the sheaf $\mathcal{F}$ is ample if and only if, for any coherent sheaf $\mathcal{G}$ on X , the sheaf $\mathcal{G} \otimes \operatorname{Sym}^m \mathcal{F}$ is globally generated for all $m \gg 0$ (see [8, Theorem 1]);
- b) any quotient of an ample sheaf is ample (see [8, Proposition 1]);
- c) if $\pi : Y \rightarrow X$ is a finite morphism, $\mathcal{F}$ is ample if and only if $\pi^* \mathcal{F}$ is (this is because $\mathbf{P}(\pi^* \mathcal{F}) = \mathbf{P}(\mathcal{F}) \times_X Y$ and $\mathcal{O}_{\mathbf{P}(\mathcal{F})}(1)$ pulls back, by a finite morphism, to $\mathcal{O}_{\mathbf{P}(\pi^* \mathcal{F})}(1)$);
- d) if X is proper and $\mathcal{F}$ is globally generated, $\mathcal{F}$ is ample if and only if, for any curve C in X , the restriction $\mathcal{F} \otimes \mathcal{O}_C$ has no trivial quotient (Gieseker's Lemma).

3. Continuously generated sheaves

Following [13, Definition 2.10], we say that a coherent sheaf $\mathcal{F}$ on an irreducible projective variety X is *continuously globally generated* if, for any nonempty subset U of $\operatorname{Pic}^0(X)$, the sum of the twisted evaluation maps

$$\bigoplus_{\xi \in U} H^0(X, \mathcal{F} \otimes P_\xi) \otimes P_\xi^\vee \longrightarrow \mathcal{F}$$

is surjective, where, for any element ξ of $\operatorname{Pic}^0(X)$, we denote by P_ξ the corresponding numerically trivial line bundle on X . This property is equivalent to the existence of a positive integer N such that for $(\xi_1, \dots, \xi_N)$ general in $\operatorname{Pic}^0(X)^N$, the analogous map

$$(1) \quad \bigoplus_{i=1}^N H^0(X, \mathcal{F} \otimes P_{\xi_i}) \otimes P_{\xi_i}^\vee \longrightarrow \mathcal{F}$$

is surjective. Being a quotient of a direct sum of numerically trivial line bundles, a continuously globally generated sheaf is nef. Our aim is to show that under certain circumstances, it is ample.

PROPOSITION 3.1. — *A coherent sheaf $\mathcal{F}$ on an irreducible projective variety X is continuously globally generated if and only if there exists a connected abelian Galois étale cover $\pi : Y \rightarrow X$ such that $\pi^*(\mathcal{F} \otimes P_\xi)$ is globally generated for all $\xi \in \text{Pic}^0(X)$.*

Proof. — Assume $\mathcal{F}$ is continuously globally generated and let $\xi \in \text{Pic}^0(X)$. Since torsion points are dense in $\text{Pic}^0(X)^N$, the open subset of $\text{Pic}^0(X)^N$ of points for which the map (1) is surjective and all $h^0(X, \mathcal{F} \otimes P_{\xi_i})$ are minimal contains a point of the type

$$(\xi + \eta_1(\xi), \dots, \xi + \eta_N(\xi))$$

where $(\eta_1(\xi), \dots, \eta_N(\xi))$ is torsion, hence contains also $U_\xi + (\eta_1(\xi), \dots, \eta_N(\xi))$, where U_ξ is a neighborhood of ξ in $\text{Pic}^0(X)$. Since $\text{Pic}^0(X)$ is quasi-compact, it is covered by finitely many such neighborhoods, say $U_{\xi_1}, \dots, U_{\xi_M}$.

Let $\pi : Y \rightarrow X$ be a connected abelian Galois étale cover such that the kernel of $\text{Pic}^0(\pi) : \text{Pic}^0(X) \rightarrow \text{Pic}^0(Y)$ contains all $\eta_i(\xi_j)$, for $i \in \{1, \dots, N\}$ and $j \in \{1, \dots, M\}$. Fix $j \in \{1, \dots, M\}$; the map

$$\bigoplus_{i=1}^N H^0(X, \mathcal{F} \otimes P_\xi \otimes P_{\eta_i(\xi_j)}) \otimes \pi^* P_\xi^\vee \otimes \pi^* P_{\eta_i(\xi_j)}^\vee \longrightarrow \pi^* \mathcal{F}$$

is surjective for all $\xi \in U_{\xi_j}$. But this map is

$$\bigoplus_{i=1}^N H^0(X, \mathcal{F} \otimes P_\xi \otimes P_{\eta_i(\xi_j)}) \otimes \pi^* P_\xi^\vee \longrightarrow \pi^* \mathcal{F}$$

and since each $H^0(X, \mathcal{F} \otimes P_\xi \otimes P_{\eta_i(\xi_j)})$ is a vector subspace of $H^0(Y, \pi^*(\mathcal{F} \otimes P_\xi))$, the sheaf $\pi^*(\mathcal{F} \otimes P_\xi)$ is globally generated for all $\xi \in U_{\xi_j}$, hence for all ξ in $\text{Pic}^0(X)$.

For the converse, assume that there exists a connected abelian Galois étale cover $\pi : Y \rightarrow X$ such that the evaluation map

$$H^0(Y, \pi^*(\mathcal{F} \otimes P_\xi)) \otimes \mathcal{O}_Y \longrightarrow \pi^*(\mathcal{F} \otimes P_\xi)$$

is surjective for all $\xi \in \text{Pic}^0(X)$. Since π is finite, the map

$$H^0(X, \mathcal{F} \otimes P_\xi \otimes \pi_* \mathcal{O}_Y) \otimes \pi_* \mathcal{O}_Y \longrightarrow \mathcal{F} \otimes P_\xi \otimes \pi_* \mathcal{O}_Y$$

is also surjective. If we let $\text{Ker}(\text{Pic}^0(\pi)) = \{\eta_1, \dots, \eta_N\}$, we have $\pi_* \mathcal{O}_Y = \bigoplus_{i=1}^N P_{\eta_i}$, the map

$$\left(\bigoplus_{i=1}^N H^0(X, \mathcal{F} \otimes P_\xi \otimes P_{\eta_i}) \right) \otimes \left(\bigoplus_{i=1}^N P_{\eta_i} \right) \longrightarrow \mathcal{F} \otimes P_\xi \otimes \left(\bigoplus_{i=1}^N P_{\eta_i} \right)$$

is surjective, and so is

$$\bigoplus_{i=1}^N H^0(X, \mathcal{F} \otimes P_\xi \otimes P_{\eta_i}) \otimes P_{\eta_i}^\vee \longrightarrow \mathcal{F} \otimes P_\xi.$$

In other words, the map (1) is surjective for $(\xi_1, \dots, \xi_N) = (\xi + \eta_1, \dots, \xi + \eta_N)$, for all $\xi \in \text{Pic}^0(X)$. Choosing ξ_0 such that $h^0(X, \mathcal{F} \otimes P_{\xi_0 + \eta_i})$ takes the general (minimal) value for each i in $\{1, \dots, N\}$, we obtain that the map (1) is still surjective for $(\xi_1, \dots, \xi_N)$ in a neighborhood of $(\xi_0 + \eta_1, \dots, \xi_0 + \eta_N)$. This proves that $\mathcal{F}$ is continuously globally generated. $\square$

COROLLARY 3.2. — *Let X an irreducible projective variety with a finite map to an abelian variety. Any continuously globally generated coherent sheaf on X is ample.*

Proof. — Let $\mathcal{F}$ be a continuously globally generated coherent sheaf on X . By Proposition 3.1, there exists a connected abelian Galois étale cover $\pi : Y \rightarrow X$ such that $\pi^*(\mathcal{F} \otimes P_\xi)$ is globally generated for all $\xi \in \text{Pic}^0(X)$.

Let C be a curve in Y . If there is a trivial quotient $\pi^*\mathcal{F}|_C \twoheadrightarrow \mathcal{O}_C$, we have also surjections $\pi^*(\mathcal{F} \otimes P_\xi)|_C \twoheadrightarrow \pi^*P_\xi|_C$ for each $\xi \in \text{Pic}^0(X)$. Since $\pi^*(\mathcal{F} \otimes P_\xi)$ is globally generated, so is $\pi^*P_\xi|_C$. This implies that the composition $\text{Pic}^0(X) \rightarrow \text{Pic}^0(Y) \rightarrow \text{Pic}^0(C)$ is zero, hence that $\pi(C)$ is contracted by any map from X to an abelian variety. This contradicts our hypothesis, hence $\pi^*\mathcal{F}|_C$ has no trivial quotient.

By Gieseker's Lemma, $\pi^*\mathcal{F}$ is ample, and so is $\mathcal{F}$ (§ 2). $\square$

4. The main theorem

Following [13, Definition 2.1], we say that a coherent sheaf $\mathcal{F}$ on an abelian variety A is *M-regular* if

$$\text{codim}_{\text{Pic}^0(A)} \text{Supp}(R^i \widehat{\mathcal{S}}(\mathcal{F})) > i$$

for all $i > 0$ ($R^i \widehat{\mathcal{S}}$ is the i th Fourier-Mukai functor). This is the case if

$$\text{codim}_{\text{Pic}^0(A)} \{\xi \in \text{Pic}^0(A) \mid H^i(A, \mathcal{F} \otimes P_\xi) \neq 0\} > i$$

for all $i > 0$. We refer to [12] and [13] for more details. For our purposes, the main result of [13] (Proposition 2.13) is that *an M-regular coherent sheaf on an abelian variety is continuously globally generated.*

THEOREM 4.1. — *Let X be a smooth connected projective variety with a finite map to a simple abelian variety, let Y be a smooth connected projective variety with a finite surjective map $f : Y \rightarrow X$. If f factors through no nontrivial connected abelian Galois étale covering of X , the vector bundle $E_f \otimes \omega_X$ is ample.*