

## NON-COMPACT FORM OF THE ELEMENTARY DISCRETE INVARIANT

BY RAPHAËL FINO

---

ABSTRACT. — We determine the non-compact form of Vishik's elementary discrete invariant for quadrics. As an application, we obtain new restrictions on the possible values of the elementary discrete invariant by studying the action of Steenrod operations on the algebraic cycles defining the non-compact form.

RÉSUMÉ (*Forme non-compacte de l'invariant discret élémentaire*). — On détermine la forme non-compacte de l'invariant discret élémentaire de Vishik pour quadriques. Comme application, on obtient de nouvelles restrictions sur les valeurs possibles de l'invariant discret élémentaire en étudiant l'action des opérations de Steenrod sur les cycles algébriques définissant la forme non-compacte.

### 1. Introduction

Let  $X$  be a smooth projective quadric of dimension  $n$  over a field  $F$  associated with a non-degenerate  $F$ -quadratic form  $q$ . The *splitting pattern* of  $X$  is a discrete invariant that measures the possible Witt indices of  $q_E$  over all field extensions  $E/F$  (see [4] and [5]). The *motivic decomposition type* of  $X$  is a discrete invariant which measures in what pieces the Chow motive of  $X$  can be

---

*Texte reçu le 19 décembre 2017, modifié le 21 août 2018, accepté le 4 septembre 2018.*

RAPHAËL FINO, Instituto de Matemáticas, Ciudad Universitaria, UNAM, DF 04510, México  
• *E-mail* : [fino@im.unam.mx](mailto:fino@im.unam.mx) • *Url* : <http://www.matem.unam.mx/fino>

Mathematical subject classification (2010). — 14C25, 11E39.

Key words and phrases. — Chow groups, quadratic forms, grassmannians, Steenrod operations.

decomposed. Moreover, Alexander Vishik noticed in [6] that the study of the interaction between these two invariants provides further information about both of them.

For this reason, he introduced the *generic discrete invariant* of quadrics, a bigger discrete invariant containing the splitting pattern and the motivic decomposition type invariants as faces, see [7] and [9]. The Generic Discrete Invariant  $GDI(X)$  is defined as follows. Let  $K/F$  be a splitting field extension of  $q$ . Let us denote  $[n/2]$  as  $d$ . For any  $i \in \{0, \dots, d\}$ , we write  $G_i$  for the grassmannian of  $i$ -dimensional totally  $q$ -isotropic subspaces (in particular  $G_0$  is the quadric  $X$ ). Then  $GDI(X)$  is the collection of the subalgebras of rational elements

$$\overline{\text{Ch}}^*(G_i) := \text{Image}(\text{Ch}^*(G_i) \rightarrow \text{Ch}^*(G_{iK}))$$

for  $i \in \{0, \dots, d\}$ , where  $\text{Ch}$  stands for the Chow ring with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients (an algebraic cycle already defined at the level of the base field  $F$  is called *rational*).

In his paper [10] dedicated to the Kaplansky’s conjecture on the  $u$ -invariant of a field, A. Vishik used the *elementary discrete invariant* of quadrics, a handier invariant than the  $GDI$  as it only deals with some particular cycles in  $\text{Ch}^*(G_{iK})$ . More precisely, for any  $i \in \{0, \dots, d\}$ , we denote by  $\mathcal{F}(0, i)$  the partial orthogonal flag variety of  $q$ -isotropic lines contained in  $i$ -dimensional totally  $q$ -isotropic subspaces. One can consider the diagram

$$X \xleftarrow{\pi_{(0,i)}} \mathcal{F}(0, i) \xrightarrow{\pi_{(0,i)}} G_i,$$

given by the natural projections and, for  $0 \leq j \leq d$ , we set

$$Z_{n-i-j}^i := \pi_{(0,i)*} \circ \pi_{(0,i)}^*(l_j) \in \text{CH}^{n-i-j}(G_{iK}),$$

where  $\text{CH}$  stands for the Chow ring with  $\mathbb{Z}$ -coefficients and  $l_j$  is the class in  $\text{CH}_j(X_K)$  of a  $j$ -dimensional totally isotropic subspace of  $\mathbb{P}((V_q)_K)$  (with  $V_q$  the  $F$ -vector space associated with  $q$ ). We set  $z_{n-i-j}^i := Z_{n-i-j}^i \pmod{2} \in \text{Ch}^{n-i-j}(G_{iK})$ , with  $\text{Ch}$  being the Chow ring with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients. The cycles  $z_{n-i-j}^i$  are the elementary classes defining the elementary discrete invariant  $EDI(X)$ :

DEFINITION 1.1. — The *elementary discrete invariant*  $EDI(X)$  is the collection of subsets  $EDI(X, i)$  consisting of those integers  $m$  such that  $z_m^i$  is rational.

Furthermore, for any  $r \geq 1$ , the Chow motive of  $X^r$  with  $\mathbb{Z}/2\mathbb{Z}$ -coefficients decomposes into a direct sum of shifts of the motive of some  $G_i$ , see [2, Corollary 91.8]. Therefore, knowing  $GDI(X)$  is the equivalent to knowing

$$\overline{\text{Ch}}^*(X^r) := \text{Image}(\text{Ch}^*(X^r) \rightarrow \text{Ch}^*(X_K^r))$$

for all  $r \geq 1$ . Hence, the collection of the latter subalgebras constitutes a *non-compact* (in the sense that one has to consider infinitely many objects) form of  $GDI(X)$ . For the same reason, there exists a non-compact form of  $EDI(X)$  (with defining cycles living in  $\text{Ch}^*(X_K^r)$ ), which we determine in this work: for any  $i \in \{0, \dots, d\}$ , let us denote by  $\text{sym} : \text{CH}^*(X^{i+1}) \rightarrow \text{CH}^*(X^{i+1})$  the homomorphism  $\sum_{s \in S_{i+1}} s_*$ , where  $s : X^{i+1} \rightarrow X^{i+1}$  is the isomorphism associated with a permutation  $s$ . For  $0 \leq j \leq d$ , we set

$$\rho_{i,j} := \text{sym} \left( \left( \times_{k=0}^{i-1} h^k \right) \times l_j \right) \in \text{CH}^{n-j+i(i-1)/2} (X_K^{i+1}),$$

where  $\times$  is the external product and  $h^k$  is the  $k$ -th power of the hyperplane section class  $h \in \text{CH}^1(X)$  (always rational). Note that  $\rho_{0,j} = Z_{n-j}^0 = l_j$ . The symmetric cycles  $\rho_{i,j} \pmod{2}$  are the classes defining the non-compact form of  $EDI(X)$ :

**THEOREM 1.2.** — *Let  $1 \leq i \leq d$  and  $0 \leq j \leq d$ . The cycle  $z_{n-i-j}^i$  is rational if and only if the cycle  $\rho_{i,j} \pmod{2}$  is rational.*

Because of the stability of rational cycles under pull-backs of diagonal morphisms and the possibility of a refined use of Steenrod operations of cohomological type, studying the non-compact form provides new restrictions on the possible values of  $EDI(X)$ , as illustrated by Sections 4 and 5.

Moreover, Theorem 1.2 reduces certain questions about the rationality of algebraic cycles on orthogonal grassmannians to the sole level of quadrics. For example, it allows one to reformulate both Vishik’s conjecture [8, Conjecture 3.11] and the conjecture [9, Conjecture 0.13] on the *dimensions of Bruno Kahn*.

In Section 2, we introduce some basic tools which are required in Section 3, where we prove Theorem 1.2, using mainly compositions of correspondences and Chern classes of vector bundles over orthogonal grassmannians.

## 2. Preliminaries

In this section, we continue to use the notation introduced in Section 1.

**2.1. Rational cycles on powers of quadrics.** — We refer to [2, § 68] for an introduction to cycles on powers of quadrics. For any  $1 \leq i \leq d$  and  $0 \leq j \leq i - 1$ , we set

$$\Delta_{i,j} := \text{sym} \left( \left( \times_{k=0}^{i-1} h^k \right) \times l_j \right) + \sum_{m=i}^d \text{sym} \left( \left( \times_{\substack{k=0 \\ k \neq j}}^{i-1} h^k \right) \times h^m \times l_m \right)$$

in  $\text{Ch}^{n-j+i(i-1)/2}(X_K^{i+1})$ . If  $n = 2d$ , we choose an orientation  $l_d$  of the quadric.

**LEMMA 2.1.** — *For any  $1 \leq i \leq d$  and  $0 \leq j \leq i - 1$ , the cycle  $\Delta_{i,j}$  is rational.*

*Proof.* — We proceed by induction on  $i$ . In  $\text{Ch}^n(X_K^2)$ , the cycle  $\Delta_{1,0}$  or  $\Delta_{1,0} + h^d \times h^d$ , depending on whether  $l_d^2 = 0$  or not, is the class of the diagonal. Therefore, the cycle  $\Delta_{1,0}$  is rational. Let  $\sigma \in S_{i+1}$  be a cyclic permutation (with  $i \geq 2$ ). For  $0 \leq j \leq i - 2$ , the induction hypothesis step is provided by the identity

$$\Delta_{i,j} = \sum_{l=0}^i \sigma_*^l (\Delta_{i-1,j} \times h^{i-1}) \quad \text{in } \text{Ch}(X_K^{i+1}).$$

It just remains to show that the cycle  $\Delta_{i,i-1}$  is rational to complete the proof. In  $\text{Ch}(X_K^{i+1})$ , one has

$$\begin{aligned} \Delta_{i,i-1} &= \sum_{m=i-1}^d \text{sym} \left( (\times_{k=0}^{i-2} h^k) \times l_m \times h^m \right) \\ &= \sum_{m=0}^d \text{sym} \left( (\times_{k=0}^{i-2} h^k) \times l_m \times h^m \right) \end{aligned}$$

and the latter sum can be rewritten as

$$\sum_{s \in A_{i+1}} s_* \left( (\times_{k=0}^{i-2} h^k) \times \Delta_{1,0} \right).$$

Thus, the cycle  $\Delta_{i,i-1}$  is rational. □

**2.2. Correspondences.** — We refer to [2, § 62] for an introduction to Chow correspondences.

For any  $1 \leq i \leq d$ , we denote by  $\theta_i$  the class of the subvariety

$$\{(y, x_1, \dots, x_{i+1}) \mid x_1, \dots, x_{i+1} \in y\} \subset G_i \times X^{i+1}$$

in  $\text{CH}(G_i \times X^{i+1})$  and we view the cycle  $\theta_i$  as a correspondence  $G_i \rightsquigarrow X^{i+1}$ .

We set

$$(1) \quad \eta_i := \prod_{k=1}^i \left( \text{Id}_{G_i} \times p_{X_k^i} \right)^* ([\mathcal{F}(i, 0)]) \in \text{CH}(G_i \times X^i),$$

with  $p_{X_k^i}$  the projection from  $X^i$  to the  $k$ -th coordinate. For any integer  $i \leq s \leq d$ , we write

$$W_{s-i}^i := \pi_{(0,i)}^* \circ \pi_{(0,i)}^* (h^s) \in \text{CH}^{s-i}(G_i),$$

and  $w_{s-i}^i := W_{s-i}^i \pmod{2} \in \text{Ch}^{s-i}(G_i)$ . Since the variety  $X_K$  is cellular, the cycle  $[\mathcal{F}(i, 0)]$  decomposes as

$$(2) \quad [\mathcal{F}(i, 0)] = \sum_{s=0}^d z_{n-i-s}^i \times h^s + \sum_{s=i}^d w_{s-i}^i \times l_s \text{ in } \text{Ch}(G_{iK} \times X_K),$$

where  $l_d$  has to be replaced by the other class  $l'_d$  of maximal totally isotropic subspaces if  $n = 2d$  and  $l_d^2$  is not zero, i.e., if four divides  $n$  (see [2, Theorem 66.2]).

The two following lemmas, where we write  $p$  with underlined target for projections, can be proven the same way [3, Lemmas 3.2 and 3.10] have been proven but with  $Z_{n-i-j}^i$  (resp.  $z_{n-i-j}^i$ ) instead of  $Z_{n-i}^i$  (resp.  $z_{n-i}^i$ ).

LEMMA 2.2. — For any  $1 \leq i \leq d, 0 \leq j \leq d$  and  $x \in CH(X_K)$ , one has

$$((\theta_i)_*(Z_{n-i-j}^i))_*(x) = p_{G_i \times \underline{X}^i} \left( p_{G_i \times X^i}^* \left( \pi_{(0,i)} \circ \pi_{(0,i)}^*(x) \cdot Z_{n-i-j}^i \right) \cdot \eta_i \right),$$

where the cycle  $(\theta_i)_*(Z_{n-i-j}^i)$  is viewed as a correspondence  $X_K \rightsquigarrow X_K^i$ .

For any  $1 \leq i \leq d$ , we write  $\mathcal{F}(i-1, i)$  for the partial orthogonal flag variety of  $(i-1)$ -dimensional totally isotropic subspaces contained in  $i$ -dimensional totally isotropic subspaces and we consider the diagram

$$G_{i-1} \xleftarrow{\pi_{(i-1,i)}} \mathcal{F}(i-1, i) \xrightarrow{\pi_{(i-1,i)}} G_i,$$

given by the natural projections.

LEMMA 2.3. — For any  $2 \leq i \leq d, 0 \leq j \leq d$  and  $i \leq m \leq d$ , the cycle

$$p_{G_i \times \underline{X}^i} (w_{m-i}^i \cdot z_{n-i-j}^i \cdot \eta_i) \in Ch(X_K^i),$$

where we write  $\eta_i$  for  $\eta_i \pmod{2}$ , and this can be rewritten as

$$\sum_{s=0}^m \sum_{k=\max(i-s, 0)}^{\min(m-s, i)} p_{G_{i-1} \times \underline{X}^{i-1}} (w_{m-s-k}^{i-1} \cdot \sigma_{i-1}^k \cdot z_{n-i+1-j}^{i-1} \cdot \eta_{i-1}) \times h^s,$$

with  $\sigma_{i-1}^k = \pi_{(i-1,i)} \circ \pi_{(i-1,i)}^*(z_{n-2i+k}^i) \in Ch^j(G_{i-1K})$ .

### 3. Equivalence

In this section, we continue to use notation and material introduced in the previous sections and we prove Theorem 1.2.

For  $1 \leq i \leq d$  and  $0 \leq j \leq i-1$ , we set

$$\alpha_{i,j} := (\theta_i)_*(Z_{n-i-j}^i) + \rho_{i,j} \in CH(X_K^{i+1}),$$

and we view the cycle  $\alpha_{i,j}$  as a correspondence  $X_K \rightsquigarrow X_K^i$ .