

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

LOWER BOUNDS ON THE DIMENSION OF THE RAUZY GASKET

Rodolfo Gutiérrez-Romo & Carlos Matheus

Tome 148
Fascicule 2

2020

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

pages 321-327

Le Bulletin de la Société Mathématique de France est un périodique trimestriel
de la Société Mathématique de France.

Fascicule 2, tome 148, juin 2020

Comité de rédaction

Christine BACHOC
Yann BUGEAUD
Jean-François DAT
Clothilde FERMANIAN
Pascal HUBERT

Laurent MANIVEL
Julien MARCHÉ
Kieran O'GRADY
Emmanuel RUSS
Christophe SABOT

Marc HERZLICH (Dir.)

Diffusion

Maison de la SMF Case 916 - Luminy 13288 Marseille Cedex 9 France commandes@smf.emath.fr	AMS P.O. Box 6248 Providence RI 02940 USA www.ams.org
--	--

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Bulletin de la SMF

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 1 44 27 67 99 • Fax : (33) 1 40 46 90 96

bulletin@smf.emath.fr • smf.emath.fr

© Société Mathématique de France 2020

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Stéphane SEURET

LOWER BOUNDS ON THE DIMENSION OF THE RAUZY GASKET

BY RODOLFO GUTIÉRREZ-ROMO & CARLOS MATHEUS

ABSTRACT. — The Rauzy gasket R is the maximal invariant set of a certain renormalization procedure for special systems of isometries naturally appearing in the context of Novikov's problem in conductivity theory for monocrystals.

It was conjectured by Novikov and Maltsev in 2003 that the Hausdorff dimension $\dim_H(R)$ of the Rauzy gasket lies strictly between 1 and 2.

In 2016, Avila, Hubert and Skripchenko confirmed that $\dim_H(R) < 2$. In this note, we use some results by Cao–Pesin–Zhao in order to show that $\dim_H(R) > 1.19$.

RÉSUMÉ (*Bornes inférieures pour la dimension de la baderne de Rauzy*). — La baderne de Rauzy R est l'ensemble maximal invariant pour une certaine procédure de renormalisation sur les systèmes d'isométries spéciaux issus du problème de Novikov en théorie de conductivité des monocristaux.

Il fut conjecturé par Novikov et Maltsev en 2003 que la dimension de Hausdorff $\dim_H(R)$ de la baderne de Rauzy est strictement comprise entre 1 et 2.

En 2016, Avila, Hubert et Skripchenko ont confirmé que $\dim_H(R) < 2$. Dans cette note, on utilise des résultats par Cao–Pesin–Zhao afin de montrer que $\dim_H(R) > 1.19$.

Texte reçu le 22 février 2019, modifié le 5 juin 2019, accepté le 28 novembre 2019.

RODOLFO GUTIÉRREZ-ROMO, Institut de Mathématiques de Jussieu – Paris Rive Gauche, UMR 7586, Bâtiment Sophie Germain, 75205 Paris Cedex 13, France •
E-mail : rodolfo.gutierrez@imj-prg.fr • Url : <http://rodolfo>
CARLOS MATHEUS, Centre de Mathématiques Laurent Schwartz, CNRS (UMR 7640), École Polytechnique, 91128 Palaiseau, France • E-mail : carlos.matheus@math.cnrs.fr •
Url : <http://carlos.matheus.perso.math.cnrs.fr>

Mathematical subject classification (2010). — 37C45; 37D35, 28A78, 28A80.

Key words and phrases. — Rauzy gasket, Novikov's problem, Hausdorff dimension, Thermodynamic formalism, Topological pressure.

1. Introduction

The Rauzy gasket is a fractal subset of the standard 2-simplex. It was given life by Arnoux and Rauzy [1] in the context of representing low-complexity subshifts as interval exchange maps, although it was only given a name later. It is also related to frequencies of letters in ternary episturmian words [2], dynamics of special systems of isometries [7], and a particular case of Novikov's problem around the trajectories of electrons on Fermi surfaces in the presence of constant magnetic fields [6, 3]. It is depicted in Figure 1.1.




FIGURE 1.1. The Rauzy gasket.

Concretely, the Rauzy gasket is defined as follows. Consider the standard 2-simplex $\Delta = \{(x_1, x_2, x_3) \in \mathbb{R}_+^3 : x_1 + x_2 + x_3 = 1\}$. We decompose Δ into three simplices $\Delta_j = \{(x_1, x_2, x_3) \in \Delta : x_j \geq \sum_{k \neq j} x_k\}$ and a hole $\Delta \setminus \bigcup_{j=1}^3 \Delta_j$. The projectivizations of the matrices

$$M_1 = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad M_2 = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad M_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix}$$

induce weakly contracting maps $f_j: \Delta \rightarrow \Delta_j$, $j = 1, 2, 3$. In this context, recall from [2] that the *Rauzy gasket* is the unique non-empty compact subset of Δ such that

$$R = f_1(R) \cup f_2(R) \cup f_3(R).$$

The fact that the Rauzy gasket has zero Lebesgue measure has been proved by several authors, including Levitt [9]¹, Arnoux–Starosta [2] and De Leo–Dynnikov [6].

1. Using an argument attributed to Yoccoz.

A general conjecture by Novikov and Matlsev [10] from 2003 implies the following particular statement about the Rauzy gasket:

CONJECTURE 1.1 (Novikov–Matlsev). — $1 < \dim_H(R) < 2$.

Some numerical experiments by De Leo and Dynnikov [6] suggest that $1.7 < \dim_H(R) < 1.8$, and Avila–Hubert–Skripchenko [4] established that $\dim_H(R) < 2$.

The main result of this note is the following theorem:

THEOREM 1.2. — $\dim_H(R) > 1.19$.

The proof of this result occupies the remainder of this text.

2. Lower bounds on the Hausdorff dimension of the Rauzy gasket

In this section, we give a lower bound on $\dim_H(R)$ via the construction of appropriate uniformly expanding repellers inside R .

2.1. General framework. — We will use somewhat general methods to obtain bounds for the Hausdorff dimension of a uniformly expanding repeller in dimension 2. These methods rely on estimating the singular values of the derivatives of the maps defining the set. More precisely, given n uniformly contracting maps $T_1, \dots, T_n: X \rightarrow X$, where $X \subseteq \mathbb{R}^2$ is a compact set, and a repeller K defined as the unique non-empty compact set such that $K = \bigcup_{k=1}^n T_k(K)$, we need to estimate quantities of the form $\max_{x \in X} \|D_x T_k\|$ and $\min_{x \in X} \|(D_x T_k)^{-1}\|$, where $\|\cdot\|$ denotes the largest singular value. Since for any $a, b, c, d \in \mathbb{R}$ one has that

$$\left\| \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right\| = \sqrt{\frac{a^2 + b^2 + c^2 + d^2 + \sqrt{(a^2 + b^2 + c^2 + d^2)^2 - 4(ad - bc)^2}}{2}},$$

we obtain the simple estimates

$$\sqrt{\frac{a^2 + b^2 + c^2 + d^2}{2}} \leq \left\| \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right\| \leq \sqrt{a^2 + b^2 + c^2 + d^2},$$

which we will write as $\|\cdot\|^- \leq \|\cdot\| \leq \|\cdot\|^+$.

There are several methods in the literature to obtain lower bounds on the Hausdorff dimension of repellers. For our purposes, the thermodynamical method of Cao–Pesin–Zhao [5] is quite useful. In a nutshell, they consider a repeller Λ of a C^2 -expanding map g on a surface, a parameter $1 \leq s \leq 2$, and the potential $\psi^s(x, g) = \log \alpha_1(x, g) + (s-1) \log \alpha_2(x, g)$, where $\alpha_1(x, g) \geq \alpha_2(x, g)$ are the singular values of $D_x g$. Observe that $\|D_x g\|^- \leq \alpha_1(x, g) \leq \|D_x g\|^+$ and $\|(D_x g)^{-1}\|^- \leq \alpha_2(x, g)^{-1} \leq \|(D_x g)^{-1}\|^+$.

By Corollary 3.1 of [5], one has that

$$\dim(\Lambda) \geq s_1,$$

where s_1 is the unique root of the equation $P(g, -\psi^s(\cdot, g)) = 0$ and $P(g, \theta)$ stands for the topological pressure of the potential θ , i.e.,

$$P(g, \theta) := \sup \left\{ h_\mu(g) + \int \theta d\mu(x) : \mu \text{ is } g\text{-invariant} \right\}$$

(see (3.2) and (2.4) in [5]). The theory of (subadditive) thermodynamical formalism (as explained² in Section 3 of [8], for instance) states that

$$P(g, \theta) < 0 \iff \sum_{m \geq 1} \sum_{x \in \text{Fix}(g^m)} \exp(\theta_m(x)) < \infty,$$

where $\theta_m(x) := \sum_{j=0}^{m-1} \theta(g^j(x))$.

In general, $s \mapsto P(g, -\psi^s(\cdot, g))$ is a continuous and strictly decreasing function of s . Therefore, $s_1 \geq s_0$ for all s_0 with

$$\sum_{m \geq 1} \sum_{x \in \text{Fix}(g^m)} \exp \left(- \sum_{j=0}^{m-1} \psi^{s_0}(g^j(x), g) \right) = \infty.$$

2.2. The Rauzy gasket. — Observe that each composition $f_k \circ f_j$, with $k \neq j$, is a contraction on Δ (cf. Lemma 2 in [2]). Thus, for each integer $n \geq 2$, the unique non-empty compact subset K_n such that

$$K_n = \bigcup_{i \in S_n} f_{i_n} \circ \cdots \circ f_{i_1}(K_n),$$

where $S_n = \{1, 2, 3\}^n \setminus \{(1, \dots, 1), (2, \dots, 2), (3, \dots, 3)\}$, is a uniformly expanding repeller contained in R .

In the following, we consider the Riemannian metric on $T\Delta = \{(v_1, v_2, v_3) \in \mathbb{R}^3 : v_1 + v_2 + v_3 = 0\}$ induced by the usual Euclidean scalar product of \mathbb{R}^3 normalized so that the vectors $(\varepsilon_1, \varepsilon_2, \varepsilon_3)$, $\{\varepsilon_1, \varepsilon_2, \varepsilon_3\} = \{-1, 0, 1\}$ have norm 1. In particular, $\mathcal{B} = \{(1, -1, 0), (-1, -1, 2)/\sqrt{3}\}$ is an orthonormal basis of $T\Delta$.

REMARK 2.1. — A natural alternative is to consider the Fubini–Study metric $d(\mathbb{R}x, \mathbb{R}y) = \frac{\|x \wedge y\|}{\|x\| \|y\|}$ on the projective space $P\mathbb{R}^3$. However, we chose the *ad hoc* Riemannian metric above because the operation of taking exterior powers would lead to heavier calculations.

The repeller K_{13} , defined by g_{13} sending each $\Delta_i = f_{i_{13}} \circ \cdots \circ f_{i_1}(\Delta)$, with $i \in S_{13}$, onto Δ , is uniformly expanding with respect to this Riemannian metric.

2. Cf. Lemma 3.2 of [8] in particular.

Indeed, we can estimate the smallest expansion factor as

$$\frac{1}{\max_{i \in S_{13}} \max_{x \in \Delta_i} \|(D_x g_{13})^{-1}\|^+}$$

to obtain a value of at least $\sqrt{3}$.

Now, denote by

$$a = \log \left(\max_{i \in S_{13}} \max_{x \in \Delta_i} \|D_x g_{13}\|^+ \right), \quad b = \log \left(1 / \min_{i \in S_{13}} \min_{x \in \Delta_i} \|(D_x g_{13})^{-1}\|^+ \right)$$

and $\text{Fix}(g_{13}^m) = \exp(cm)$ for all m (i.e., $c = \log(|S_{13}|) = \log(3^{13} - 3)$). Observe that

$$\log \alpha_1(x, g) \leq a \quad \text{and} \quad \log \alpha_2(x, g) \leq b$$

for every $x \in \bigcup_{i \in S_{13}} \Delta_i$. Hence,

$$\sum_{j=0}^{m-1} \psi^{s_0}(g_{13}^j(x), g_{13}) \leq (a + b(s_0 - 1))m,$$

and we deduce that

$$\begin{aligned} & \sum_{m \geq 1} \sum_{x \in \text{Fix}(g_{13}^m)} \exp \left(- \sum_{j=0}^{m-1} \psi^{s_0}(g_{13}^j(x), g_{13}) \right) \\ & \geq \sum_{m \geq 1} \exp((c - a - b(s_0 - 1))m) = \infty, \end{aligned}$$

if $c - a - b(s_0 - 1) > 0$, i.e., $s_0 < 1 + (c - a)/b$.

This way, we obtain the bound

$$\dim_H(K_{13}) \geq s_1 \geq 1 + \frac{c - a}{b}.$$

With the help of a computer, we can find the exact values of a and b . We obtain:

$$a = \log \left(3208 \sqrt{\frac{86185}{3}} \right), \quad b = \log \left(4917248 \sqrt{\frac{2}{1595}} \right) \quad \text{and} \quad c = \log(1594320),$$

which yields $\dim_H(K_{13}) \geq 1 + \frac{c-a}{b} > 1.08$.

This lower bound can be improved by restricting it to a smaller fractal. Indeed, instead of using every sequence in S_{13} , we can take a subset of such sequences designed to optimize the previous bound by decreasing the values of a and b while trying to maintain a large value of c . The heuristic we use is as follows:

1. Sort the $i \in S_{13}$ according to $\max_{x \in \Delta_i} \|D_x g_{13}\|^+$ in ascending order, assigning a number $r^+(i)$ to each $i \in S_n$.