

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

CHARACTERISATION OF THE POLES OF THE ℓ -MODULAR ASAI L-FACTOR

Robert Kurinczuk & Nadir Matringe

Tome 148
Fascicule 3

2020

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

pages 481-514

Le Bulletin de la Société Mathématique de France est un périodique trimestriel

de la Société Mathématique de France.

Fascicule 3, tome 148, septembre 2020

Comité de rédaction

Christine BACHOC

Yann BUGEAUD

Jean-François DAT

Clothilde FERMANIAN

Pascal HUBERT

Laurent MANIVEL

Julien MARCHÉ

Kieran O'GRADY

Emmanuel RUSS

Christophe SABOT

Marc HERZLICH (Dir.)

Diffusion

Maison de la SMF

Case 916 - Luminy

13288 Marseille Cedex 9

France

commandes@smf.emath.fr

AMS

P.O. Box 6248

Providence RI 02940

USA

www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Bulletin de la SMF

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 1 44 27 67 99 • Fax : (33) 1 40 46 90 96

bulletin@smf.emath.fr • smf.emath.fr

© Société Mathématique de France 2020

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Fabien DURAND

CHARACTERISATION OF THE POLES OF THE ℓ -MODULAR ASAI L-FACTOR

BY ROBERT KURINCUK & NADIR MATRINGE

ABSTRACT. — Let F/F_\circ be a quadratic extension of non-archimedean local fields of odd residual characteristic, set $G = \mathrm{GL}_n(F)$, $G_\circ = \mathrm{GL}_n(F_\circ)$ and let ℓ be a prime number different from the residual characteristic of F . For a complex cuspidal representation π of G , the Asai L-factor $L_{\mathrm{As}}(X, \pi)$ has a pole at $X = 1$, if and only if π is G_\circ -distinguished. In this paper, we solve the problem of characterising the occurrence of a pole at $X = 1$ of $L_{\mathrm{As}}(X, \pi)$ when π is an ℓ -modular cuspidal representation of G ; we show that $L_{\mathrm{As}}(X, \pi)$ has a pole at $X = 1$, if and only if π is a *relatively banal* distinguished representation, namely π is G_\circ -distinguished but not $|\det(\)|_{F_\circ}$ -distinguished. This notion turns out to be an exact analogue for the symmetric space G/G_\circ of Mínguez and Sécherre's notion of banal cuspidal $\overline{\mathbb{F}_\ell}$ -representation of G_\circ . Along the way, we compute the Asai L-factor of all cuspidal ℓ -modular representations of G in terms of type theory and prove new results concerning lifting and reduction modulo ℓ of distinguished cuspidal representations. Finally, we determine when the natural G_\circ -period on the Whittaker model of a distinguished cuspidal representation of G is non-zero.

Texte reçu le 25 juillet 2018, accepté le 26 février 2020.

ROBERT KURINCUK, Department of Mathematics, Imperial College, London, SW7 2AZ,
United Kingdom • *E-mail :* robkurinczuk@gmail.com

NADIR MATRINGE, Université de Poitiers, Laboratoire de Mathématiques et Applications,
Téléport 2 - BP 30179, Boulevard Marie et Pierre Curie, 86962, Futuroscope Chasseneuil
Cedex. France • *E-mail :* nadir.matringe@math.univ-poitiers.fr

Mathematical subject classification (2010). — 22E50, 11F70, 11F66.

Key words and phrases. — Asai L-factor, Distinguished representations of p -adic groups,
Modular representations of p -adic groups.

This work has benefited from support from GDRI *Representation Theory* 2016–2020 and the
EPSRC grant EP/R009279/1, and the Heilbronn Institute for Mathematical Research.

RÉSUMÉ (*Caractérisation des pôles du facteur d’Asai ℓ -modulaire*). — Soit F/F_\circ une extension quadratique de corps locaux non archimédien de caractéristique résiduelle impaire. Posons $G = \mathrm{GL}_n(F)$, $G_\circ = \mathrm{GL}_n(F_\circ)$ et soit ℓ un nombre premier différent de la caractéristique résiduelle de F . Pour une représentation cuspidale complexe π de G , le facteur L d’Asai $L_{\mathrm{As}}(X, \pi)$ admet un pôle en $X = 1$ si et seulement si π est G_\circ -distinguée. Dans cet article nous résolvons le problème de l’occurrence d’un pôle en $X = 1$ de $L_{\mathrm{As}}(X, \pi)$ quand π est une représentation cuspidale ℓ -modulaire de G : dans ce cas $L_{\mathrm{As}}(X, \pi)$ admet un pôle en $X = 1$ si et seulement si π est *relativement banale* distinguée; autrement dit π est G_\circ -distinguée mais pas $|\det(\cdot)|_{F_\circ}$ -distinguée. Cette notion est l’analogue pour l’espace symétrique G/G_\circ de la notion de cuspidale banale introduite par Mínguez et Sécherre pour les $\overline{\mathbb{F}_\ell}$ -représentations de G_\circ . En cours de route, on calcule le facteur L d’Asai des représentations cuspidales ℓ -modulaires de G par la théorie des types, et on prouve de nouveaux résultats concernant le relèvement et la réduction modulo ℓ des représentations cuspidales distinguées. Finalement, on détermine quand la G_\circ -période sur le modèle de Whittaker d’une représentation cuspidale distinguée de G est non nulle.

1. Introduction

Let F/F_\circ be a quadratic extension of non-archimedean local fields of residual characteristic $p \neq 2$ and set $G = \mathrm{GL}_n(F)$ and $G_\circ = \mathrm{GL}_n(F_\circ)$. An irreducible representation of G is said to be *distinguished by G_\circ* , if it possesses a non-zero G_\circ -invariant linear form. In the case of complex representations, the equality of the Asai L-factor defined by the Rankin–Selberg method and its Galois avatar ([3], [21]) provides a bridge between functorial lifting from the quasi-split unitary group $U_n(F/F_\circ)$ and G_\circ -distinction of discrete series representations of G ; a discrete series of G is a (stable or unstable depending on the parity of n) lift of a (necessarily discrete series) representation of $U_n(F/F_\circ)$, if and only if the Asai L-factor of its Galois parameter has a pole at $X = 1$ ([25], [10]), whereas it is G_\circ -distinguished, if and only if its Asai L-factor obtained by the Rankin–Selberg method has a pole at $X = 1$ ([14], [1]).

Recently, motivated by the study of congruences between automorphic representations, there has been great interest in studying representations of G on vector spaces over fields of positive characteristic ℓ . There are two very different cases: when $\ell = p$ and when $\ell \neq p$. This article focuses on the latter $\ell \neq p$ case, where there is a theory of Haar measure that allows us to define Asai L-factors via the Rankin–Selberg method as in the complex case (Section 7).

The aim of this article is to show that in this case, a connection remains between the poles of Asai L-factor and distinction; however, this connection no longer characterises distinction, but a more subtle notion, which we call a *relatively banal* distinction. The easiest way to state that a cuspidal distinguished ℓ -modular representation is relatively banal is to say that it is not $|\det(\cdot)|_{F_\circ}$ -distinguished, where $|\det(\cdot)|_{F_\circ}$ is considered as an $\overline{\mathbb{F}_\ell}$ -valued character, but other compact definitions can also be given in terms of type theory, as well as in terms of its supercuspidal lifts:

PROPOSITION 1.1 (Definition 6.2, Theorem 6.11 and Corollary 6.3). — *Let π be an ℓ -modular cuspidal distinguished representation of G . Then, the following are equivalent, and when they are satisfied, we say that π is relatively banal:*

- (i) π is not $|\det(\cdot)|_{F_\circ}$ -distinguished.
- (ii) All supercuspidal lifts of π are distinguished by an unramified character of G_\circ .
- (iii) $q_\circ^{n/e_\circ(\pi)} \not\equiv 1[\ell]$, where $e_\circ(\pi)$ denotes the invariant associated to π in [4, Lemma 5.10] (see Section 5.2).

Relatively banal for G_\circ -distinguished cuspidal representations turns out to be the exact analogue of the definition of *banal cuspidal representations* of G_\circ (see [24, Remark 8.15] and [23]) after one identifies the cuspidal (irreducible) representations of G_\circ with the $\Delta(G_\circ)$ -distinguished cuspidal (irreducible) representations of $G_\circ \times G_\circ$, where $\Delta : G_\circ \rightarrow G_\circ \times G_\circ$ is the diagonal embedding, as we explain in Section 8.3.

The main theorem of this paper characterises the poles of the Asai L-factor:

THEOREM 1.2 (Theorem 8.1). — *Let π be a cuspidal ℓ -modular representation, then $L_{As}(X, \pi)$ has a pole at $X = 1$, if and only if π is relatively banal distinguished.*

Note that the proof of the above theorem is completely different from the proof of characterisation theorem in the complex case (see Remark 8.2 for more on the comparison of the proofs). Here, we show that the Asai L-factor of a cuspidal ℓ -modular representation is equal to 1 whenever π is not the unramified twist of a relatively banal representation using Theorem 6.11, which is the characterisation of relatively banal in terms of supercuspidal lifts. Then, when π is the unramified twist of relatively banal representation, following our paper [18], we get an explicit formula for $L_{As}(X, \pi)$ in Theorem 7.8 from the test vector computation of [4], which due to the relatively banal assumption (more precisely its type theory version) reduces modulo ℓ without vanishing. We then deduce Theorem 1.2 from this computation, together with the computation of the group of unramified characters μ of G_\circ , such that π is μ -distinguished (Corollary 5.17).

Finally, denoting by N the unipotent radical of the group of upper triangular matrices in G , by Z_\circ the centre of G_\circ and by N_\circ the group $N \cap G_\circ$, the most natural G_\circ -invariant linear form to consider on the Whittaker model of an ℓ -modular cuspidal representation π with respect to a distinguished non-degenerate character of N trivial on N_\circ is the local period

$$\mathcal{L}_\pi : W \mapsto \int_{Z_\circ N_\circ \backslash G_\circ} W(h) dh.$$

In fact, this period plays an essential role in the proof of Theorem 1.2 over the field of complex numbers (see Remark 8.2). One of the main differences in the