

*quatrième série - tome 48      fascicule 5      septembre-octobre 2015*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Syu KATO

*A homological study of Green polynomials*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> janvier 2015

N. ANANTHARAMAN B. KLEINER  
E. BREUILLARD E. KOWALSKI  
R. CERF P. LE CALVEZ  
A. CHAMBERT-LOIR M. MUSTAȚĂ  
I. GALLAGHER L. SALOFF-COSTE

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

### Édition / *Publication*

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

### Abonnements / *Subscriptions*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

### Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

---

© 2015 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Marc Peigné  
Périodicité : 6 n<sup>os</sup> / an

## A HOMOLOGICAL STUDY OF GREEN POLYNOMIALS\*

BY SYU KATO

---

**ABSTRACT.** – We interpret the orthogonality relation of Kostka polynomials arising from complex reflection groups ([51, 52] and [35]) in terms of homological algebra. This leads us to the notion of Kostka system, which can be seen as a categorical counterpart of Kostka polynomials. Then, we show that every generalized Springer correspondence ([34]) in a good characteristic gives rise to a Kostka system. This enables us to see the top-term generation property of the (twisted) homology of generalized Springer fibers, and the transition formula of Kostka polynomials between two generalized Springer correspondences of type BC. The latter provides an inductive algorithm to compute Kostka polynomials by upgrading [16] §3 to its graded version. In the appendices, we present purely algebraic proofs that Kostka systems exist for type A and asymptotic type BC cases, and therefore one can skip geometric sections §3–5 to see the key ideas and basic examples/techniques.

**RÉSUMÉ.** – La relation d’orthogonalité des polynômes de Kostka émanant des groupes de réflexions complexes ([51, 52] et [35]) est interprétée en termes d’algèbre homologique. Ceci nous conduit à la notion de système Kostka, qui peut être considérée comme une contrepartie catégorique des polynômes de Kostka. Puis, nous démontrons que chaque correspondance de Springer généralisée ([34]) dans une bonne caractéristique engendre un système de Kostka. Nous pouvons ainsi observer la propriété de génération du premier terme de l’homologie (tordue) des fibres de Springer généralisées, ainsi que la formule de transition de polynômes de Kostka entre deux correspondances de Springer généralisées de type BC. Cette dernière fournit un algorithme inductif de calcul des polynômes de Kostka par la mise à niveau de [16] §3 à sa version graduée. Dans les annexes, nous apportons les preuves algébriques que les systèmes de Kostka existent pour les cas de type A et de type BC asymptotique. Aussi, il est possible d’omettre de lire les sections géométriques 3 à 5 et pour entrevoir les idées-clés et parcourir des exemples/techniques de base.

---

\* The word “green” means ‘midori’ in Japanese.

### Introduction

Green polynomials attached to a connected reductive group over a finite field is a family of polynomials indexed by two conjugacy classes of their (little) Weyl groups<sup>(1)</sup>, depending on a variable  $t$  that corresponds to (some twist of) the cardinality of the base field. Introduced by Green [28] for  $GL(n, \mathbb{F}_q)$  and by Deligne-Lusztig [21] in general, they play a central role in the representation theory of finite groups of Lie types, affine Hecke algebras,  $p$ -adic groups, and so on. Equivalent to Green polynomials are Kostka polynomials ([28, 38]), which are  $t$ -analogues of Kostka numbers (Kostka-Foulkes polynomials) in the case of  $GL(n)$  (cf. [43] Chapter III). Hence, they appear almost everywhere in representation theory attached to root data.

Despite their natural appearance, not much is known about Kostka polynomials except for type A. One major reason seems to be the fact that the set of Kostka polynomials admits integral parameters, which actually yield different collections of polynomials even if they arise from character sheaves of Chevalley groups over finite fields ([34, 35, 38]). In such representation theoretic situation, Lusztig [34] introduced the notion of symbols, which govern the combinatorial data to determine Kostka polynomials by means of their *orthogonality relation* ([51, 35]). It is generalized by Malle [44] and Shoji [52, 53] to include the case of complex reflection groups, in which the orthogonality relation is employed as their definition.

In [5], Arthur initiated now so-called elliptic representation theory, that is the “cuspidal quotient” of (usual) representation theory. Green polynomials, in the guise of characters of discrete series representations, also appear in the context of elliptic representation theory ([50, 18]). In particular, the study of formal degrees of affine Hecke algebras/ $p$ -adic groups ([47, 48, 16]) revealed the transition pattern of Kostka polynomials evaluated at  $t = 1$ . This supplies connections among representation theories of infinitely many  $p$ -adic groups (of different types).

The goal of the present paper is two-fold: One is to afford an algebraic framework of the study of Green/Kostka polynomials of complex reflection groups. The other is to exhibit how the classical results on Kostka polynomials of Weyl groups and the above transition pattern unveil their finer versions in our framework. From these, we expect that our framework is suited to study global structures of families of (the sets of) Kostka polynomials, and to study their connections with elliptic/usual representation theory of reductive groups or “spetses” ([13]).

For more detailed explanation, we need notations: Let  $W$  be a complex reflection group, and let  $\text{Irr } W$  denote the set of isomorphism classes of irreducible  $W$ -modules. For each  $\chi \in \text{Irr } W$ , we denote by  $\chi^\vee$  its dual representation. Let  $\mathfrak{h}$  be a reflection representation of  $W$ . Form a graded algebra  $A_W := \mathbb{C}W \rtimes \mathbb{C}[\mathfrak{h}^*]$  with  $\deg w = 0$  ( $w \in W$ ) and  $\deg x = 2$  ( $x \in \mathfrak{h}$ ). Let  $A_W\text{-gmod}$  be the category of finitely generated  $\mathbb{Z}$ -graded  $A_W$ -modules. For  $E, F \in A_W\text{-gmod}$ , we define

$$\langle E, F \rangle_{\text{gEP}} := \sum_{i \geq 0} (-1)^i \text{gdim ext}_{A_W}^i(E, F) \in \mathbb{Z}((t^{1/2})),$$

<sup>(1)</sup> The subgroup of the Weyl group that preserves the cuspidal datum (cf. §3). In case the nilpotent orbit in the cuspidal datum is  $\{0\}$ , then it coincides with the whole Weyl group.

where  $\text{ext}$  means the graded extension (defined so that forgetting graded vector space structure yields the usual extension; see §1.1), and  $\text{gdim}$  means the graded dimension (which sends a  $\mathbb{Z}$ -graded vector space  $V = \bigoplus_{j \gg -\infty} V_j$  to  $\sum_j t^{j/2} \dim V_j$ ). For each  $\chi \in \text{lrr } W$ , we denote by  $L_\chi$  the irreducible graded  $A_W$ -module sitting at degree 0 that is isomorphic to  $\chi$  as a  $W$ -module.

DEFINITION A ( $\doteq$  Definition 2.13). – Let  $<$  be a total pre-order on  $\text{lrr } W$ . Then, a Kostka system  $\{K_\chi^\pm\}_\chi \subset A_W\text{-gmod}$  is a collection such that

1. Each  $K_\chi^\pm$  is an indecomposable  $A_W$ -module with simple head  $L_\chi$ ;
2. For each  $\chi, \eta \in \text{lrr } W$ , we have equalities

$$[K_\chi^+] = [L_\chi] + \sum_{\eta > \chi} K_{\chi, \eta}^+ [L_\eta] \quad \text{with } K_{\chi, \eta}^+ \in t\mathbb{N}[t] \text{ and}$$

$$[K_{\chi^\vee}^-] = [L_{\chi^\vee}] + \sum_{\eta > \chi} K_{\chi, \eta}^- [L_{\eta^\vee}] \quad \text{with } K_{\chi, \eta}^- \in t\mathbb{N}[t]$$

in the Grothendieck group of  $A_W\text{-gmod}$ ;

3. We have  $\langle K_\chi^+, (K_\eta^-)^* \rangle_{\text{gEP}} = 0$  for  $\chi \not\sim \eta^\vee$ , where  $(K_\eta^-)^*$  is the graded dual of  $K_\eta^-$ .

If  $W$  is a real reflection group, then we have  $K_\chi^+ = K_\chi^-$  by (the genuine) definition, and we denote them by  $K_\chi$ .

This definition is slightly weaker than the one presented in the main body of the paper (for simplicity). For Weyl groups, the classical preorders on  $\text{lrr } W$  reflect the geometry of nilpotent cones and the Springer correspondences.

THEOREM B (= Theorem 2.17). – For a Kostka system  $\{K_\chi^\pm\}_\chi$ , its graded character multiplicities  $K_{\chi, \eta}^\pm$  satisfy the orthogonality relation of Kostka polynomials in the sense of [51, 35, 52]. In particular, a Kostka system is an enhancement of Kostka polynomials.

There are a number of (conjectural) cases where Kostka polynomials of complex reflection groups satisfy the positivity of their coefficients ([44, 52, 53]). Theorem B supplies a possible framework in which such Kostka polynomials might obtain mathematical reality.

This possibility is supported by the following results that most of the Kostka polynomials in representation theory of reductive groups give rise to Kostka systems by giving graded categorifications of many of their properties:

THEOREM C (= part of Theorem 3.5 and Corollary 3.9). – Every set of Kostka polynomials arising from character sheaves of a connected reductive group over a finite field  $\mathbb{F}$  admits a realization as a Kostka system whenever  $\text{char } \mathbb{F}$  is good. In addition, such Kostka systems are semi-orthogonal in the sense

$$(0.1) \quad \text{ext}_{A_W}^\bullet(K_\chi, K_\eta) = \{0\} \quad \text{if } \chi < \eta.$$

REMARK D. – Note that for a Weyl group of type  $A_n$ , the set of Kostka polynomials is unique up to tensoring  $\text{sgn}$ , while for a Weyl group of type  $BC_n$ , we have at least  $4(n - 1)$  different sets of Kostka polynomials.