

quatrième série - tome 49 fascicule 3 mai-juin 2016

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Gavril FARKAS & Alessandro VERRA

The universal abelian variety over \mathcal{A}_5

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2016

N. ANANTHARAMAN I. GALLAGHER
P. BERNARD B. KLEINER
E. BREUILLARD E. KOWALSKI
R. CERF M. MUSTAȚĂ
A. CHAMBERT-LOIR L. SALOFF-COSTE

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition / *Publication*

Société Mathématique de France
Institut Henri Poincaré
11, rue Pierre et Marie Curie
75231 Paris Cedex 05
Tél. : (33) 01 44 27 67 99
Fax : (33) 01 40 46 90 96

Abonnements / *Subscriptions*

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 09
Fax : (33) 04 91 41 17 51
email : smf@smf.univ-mrs.fr

Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

© 2016 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret
Périodicité : 6 n^{os} / an

THE UNIVERSAL ABELIAN VARIETY OVER \mathcal{A}_5

BY GAVRIL FARKAS AND ALESSANDRO VERRA

ABSTRACT. – We establish a structure result for the universal abelian variety over \mathcal{A}_5 . This implies that the boundary divisor of $\overline{\mathcal{A}}_6$ is unirational and leads to a lower bound on the slope of the cone of effective divisors on $\overline{\mathcal{A}}_6$.

RÉSUMÉ. – On établit un théorème de structure pour la variété abélienne universelle sur \mathcal{A}_5 . Le résultat entraîne que le diviseur de la frontière de $\overline{\mathcal{A}}_6$ est unirationnel et ceci donne lieu à une borne inférieure pour la pente du cône des diviseurs effectifs en $\overline{\mathcal{A}}_6$.

The general principally polarized abelian variety $[A, \Theta] \in \mathcal{A}_g$ of dimension $g \leq 5$ can be realized as a Prym variety. Abelian varieties of small dimension can be studied in this way via the rich and concrete theory of curves, in particular, one can establish that \mathcal{A}_g is unirational in this range. In the case $g = 5$, the Prym map $P : \mathcal{R}_6 \rightarrow \mathcal{A}_5$ is finite of degree 27, see [7]; three different proofs [6, 17], [22] of the unirationality of \mathcal{R}_6 are known. The moduli space \mathcal{A}_g is of general type for $g \geq 7$, see [12, 18], [21]. Determining the Kodaira dimension of \mathcal{A}_6 is a notorious open problem.

The aim of this paper is to give a simple unirational parametrization of the universal abelian variety over \mathcal{A}_5 and hence of the boundary divisor of a compactification of \mathcal{A}_6 . We denote by $\phi : \mathcal{X}_{g-1} \rightarrow \mathcal{A}_{g-1}$ the universal abelian variety of dimension $g - 1$ (in the sense of stacks). The moduli space $\tilde{\mathcal{A}}_g$ of principally polarized abelian varieties of dimension g and their rank 1 degenerations is a partial compactification of \mathcal{A}_g obtained by blowing up \mathcal{A}_{g-1} in the Satake compactification, cf. [18]. Its boundary $\partial\tilde{\mathcal{A}}_g$ is isomorphic to the universal Kummer variety in dimension $g - 1$ and there exists a surjective double covering $j : \mathcal{X}_{g-1} \rightarrow \partial\tilde{\mathcal{A}}_g$. We establish a simple structure result for the boundary $\partial\tilde{\mathcal{A}}_6$:

THEOREM 0.1. – *The universal abelian variety \mathcal{X}_5 is unirational.*

This immediately implies that the boundary divisor $\partial\tilde{\mathcal{A}}_6$ is unirational as well. What we prove is actually stronger than Theorem 0.1. Over the moduli space \mathcal{R}_g of smooth Prym curves of genus g , we consider the universal Prym variety $\varphi : \mathcal{Y}_g \rightarrow \mathcal{R}_g$ obtained by pulling back $\mathcal{X}_{g-1} \rightarrow \mathcal{A}_{g-1}$ via the Prym map $P : \mathcal{R}_g \rightarrow \mathcal{A}_{g-1}$. Let $\overline{\mathcal{R}}_g$ be the moduli space of stable

Prym curves of genus g together with the universal Prym curve $\tilde{\pi} : \tilde{\mathcal{C}} \rightarrow \overline{\mathcal{R}}_g$ of genus $2g - 1$. If $\tilde{\mathcal{C}}^{g-1} := \tilde{\mathcal{C}} \times_{\overline{\mathcal{R}}_g} \cdots \times_{\overline{\mathcal{R}}_g} \tilde{\mathcal{C}}$ is the $(g - 1)$ -fold product, one has a universal *Abel-Prym* rational map $\mathbf{ap} : \tilde{\mathcal{C}}^{g-1} \dashrightarrow \mathcal{Y}_g$, whose restriction on each individual Prym variety is the usual Abel-Prym map. The rational map \mathbf{ap} is dominant and generically finite (see Section 4 for details). We prove the following result:

THEOREM 0.2. – *The five-fold product $\tilde{\mathcal{C}}^5$ of the universal Prym curve over $\overline{\mathcal{R}}_6$ is unirational.*

The key idea in the proof of Theorem 0.2 is to view smooth Prym curves of genus 6 as discriminants of conic bundles, via their representation as symmetric determinants of quadratic forms in three variables. We fix four general points $u_1, \dots, u_4 \in \mathbf{P}^2$ and set $w_i := (u_i, u_i) \in \mathbf{P}^2 \times \mathbf{P}^2$. Since the action of the automorphism group $\text{Aut}(\mathbf{P}^2 \times \mathbf{P}^2)$ on $\mathbf{P}^2 \times \mathbf{P}^2$ is 4-transitive, any set of four general points in $\mathbf{P}^2 \times \mathbf{P}^2$ can be brought to this form. We then consider the linear system

$$\mathbf{P}^{15} := \left| \mathcal{S}_{\{w_1, \dots, w_4\}}^2(2, 2) \right| \subset \left| \mathcal{O}_{\mathbf{P}^2 \times \mathbf{P}^2}(2, 2) \right|$$

of hypersurfaces $Q \subset \mathbf{P}^2 \times \mathbf{P}^2$ of bidegree $(2, 2)$ which are nodal at w_1, \dots, w_4 . For a general threefold $Q \in \mathbf{P}^{15}$, the first projection $p : Q \rightarrow \mathbf{P}^2$ induces a conic bundle structure with a sextic discriminant curve $\Gamma \subset \mathbf{P}^2$ such that $p(\text{Sing}(Q)) = \text{Sing}(\Gamma)$. The discriminant curve Γ is nodal precisely at the points u_1, \dots, u_4 . Furthermore, Γ is equipped with an unramified double cover $p_\Gamma : \tilde{\Gamma} \rightarrow \Gamma$, parametrizing the lines which are components of the singular fibres of $p : Q \rightarrow \mathbf{P}^2$. By normalizing, p_Γ lifts to an unramified double cover $f : \tilde{C} \rightarrow C$ between the normalization \tilde{C} of $\tilde{\Gamma}$ and the normalization C of Γ respectively. Note that there exists an exact sequence of generalized Prym varieties

$$0 \longrightarrow (\mathbf{C}^*)^4 \longrightarrow P(\tilde{\Gamma}/\Gamma) \longrightarrow P(\tilde{C}/C) \longrightarrow 0.$$

One can show without much effort that the assignment $\mathbf{P}^{15} \ni Q \mapsto [\tilde{C} \xrightarrow{f} C] \in \mathcal{R}_6$ is dominant. This offers an alternative, much simpler, proof of the unirationality of \mathcal{R}_6 . However, much more can be obtained with this construction.

Let $\mathbf{G} := \mathbf{P}^2 \times (\mathbf{P}^2)^\vee = \{(o, \ell) : o \in \mathbf{P}^2, \ell \in \{o\} \times (\mathbf{P}^2)^\vee\}$ be the Hilbert scheme of lines in the fibres of the first projection $p : \mathbf{P}^2 \times \mathbf{P}^2 \rightarrow \mathbf{P}^2$. Since containing a given line in a fibre of p imposes *three* linear conditions on the linear system \mathbf{P}^{15} of threefolds $Q \subset \mathbf{P}^2 \times \mathbf{P}^2$ as above, it follows that imposing the condition $\{o_i\} \times \ell_i \subset Q$ for *five* general lines, singles out a *unique* conic bundle $Q \in \mathbf{P}^{15}$. This induces an étale double cover $f : \tilde{C} \rightarrow C$, as above, over a smooth curve of genus 6. Moreover, f comes equipped with five marked points $\ell_1, \dots, \ell_5 \in \tilde{C}$. To summarize, we can define a rational map

$$\zeta : \mathbf{G}^5 \dashrightarrow \tilde{\mathcal{C}}^5, \quad \zeta\left((o_1, \ell_1), \dots, (o_5, \ell_5)\right) := \left(f : \tilde{C} \rightarrow C, \ell_1, \dots, \ell_5\right),$$

between two 20-dimensional varieties, where \mathbf{G}^5 denotes the 5-fold product of \mathbf{G} .

THEOREM 0.3. – *The morphism $\zeta : \mathbf{G}^5 \dashrightarrow \tilde{\mathcal{C}}^5$ is dominant, so that $\tilde{\mathcal{C}}^5$ is unirational.*

More precisely, we show that \mathbf{G}^5 is birationally isomorphic to the fibre product $\mathbf{P}^{15} \times_{\mathcal{R}_6} \tilde{\mathcal{C}}^5$. In order to set Theorem 0.3 on the right footing and in view of enumerative calculations, we introduce a \mathbf{P}^2 -bundle $\pi : \mathbf{P}(\mathcal{M}) \rightarrow S$ over the quintic del Pezzo surface S obtained by blowing up \mathbf{P}^2 at the points u_1, \dots, u_4 . The rank 3 vector bundle \mathcal{M} on S is obtained by making an elementary transformation along the four exceptional divisors E_1, \dots, E_4 over u_1, \dots, u_4 . The nodal threefolds $Q \subset \mathbf{P}^2 \times \mathbf{P}^2$ considered above can be thought of as sections of a tautological linear system on $\mathbf{P}(\mathcal{M})$ and, via the identification

$$\left| \mathcal{I}_{\{w_1, \dots, w_4\}}^2(2, 2) \right| = \left| \mathcal{O}_{\mathbf{P}(\mathcal{M})}(2) \right|,$$

we can view 4-nodal conic bundles in $\mathbf{P}^2 \times \mathbf{P}^2$ as *smooth* conic bundles over S . In this way the numerical characters of a pencil of such conic bundles can be computed (see Sections 2 and 3 for details).

Theorem 0.3 is then used to give a lower bound for the slope of the effective cone of $\overline{\mathcal{A}}_6$ (though we stop short of determining the Kodaira dimension of $\overline{\mathcal{A}}_6$). Recall that if E is an effective divisor on the perfect cone compactification $\overline{\mathcal{A}}_g$ of \mathcal{A}_g with no component supported on the boundary $D_g := \overline{\mathcal{A}}_g - \mathcal{A}_g$ and $[E] = a\lambda_1 - b[D_g]$, where $\lambda_1 \in CH^1(\tilde{\mathcal{A}}_g)$ is the Hodge class, then the slope of E is defined as $s(E) := \frac{a}{b} \geq 0$. The slope $s(\overline{\mathcal{A}}_g)$ of the effective cone of divisors of $\overline{\mathcal{A}}_g$ is the infimum of the slopes of all effective divisors on $\overline{\mathcal{A}}_g$. This important invariant governs to a large extent the birational geometry of $\overline{\mathcal{A}}_g$; for instance, since $K_{\overline{\mathcal{A}}_g} = (g + 1)\lambda_1 - [D_g]$, the variety $\overline{\mathcal{A}}_g$ is of general type if $s(\overline{\mathcal{A}}_g) < g + 1$, and uniruled when $s(\overline{\mathcal{A}}_g) > g + 1$. It is shown in the appendix of [14] that the slope of the moduli space $\overline{\mathcal{A}}_g$ is independent of the choice of a toroidal compactification.

It is known that $s(\overline{\mathcal{A}}_4) = 8$ and that the Jacobian locus $\mathcal{M}_4 \subset \overline{\mathcal{A}}_4$ achieves the minimal slope [19]; one of the results of [9] is the calculation $s(\overline{\mathcal{A}}_5) = \frac{54}{7}$. Furthermore, the only irreducible effective divisor on $\overline{\mathcal{A}}_5$ of minimal slope is the closure of the Andreotti-Mayer divisor N'_0 consisting of 5-dimensional ppav's $[A, \Theta]$ for which the theta divisor Θ is singular at a point which is not 2-torsion. Concerning $\overline{\mathcal{A}}_6$, we establish the following estimate:

THEOREM 0.4. – *The following lower bound holds: $s(\overline{\mathcal{A}}_6) \geq \frac{53}{10}$.*

Note that this is the first concrete lower bound on the slope of $\overline{\mathcal{A}}_6$. The idea of proof of Theorem 0.4 is very simple. Since $\tilde{\mathcal{C}}^5$ is unirational, we choose a suitable sweeping rational curve $i : \mathbf{P}^1 \rightarrow \tilde{\mathcal{C}}^5$, which we then push forward to $\overline{\mathcal{A}}_6$ as follows:

$$\mathbf{P}^1 \xrightarrow{i} \tilde{\mathcal{C}}^5 \xrightarrow{\text{ap}} \tilde{\mathcal{Y}}_6 \longrightarrow \tilde{\mathcal{X}}_5 \xrightarrow{j} D_6.$$

h

Here $\tilde{\mathcal{Y}}_6$ and $\tilde{\mathcal{X}}_5$ are partial compactifications of \mathcal{Y}_6 and \mathcal{X}_5 respectively which are described in Section 4, whereas D_6 is the boundary divisor of $\overline{\mathcal{A}}_6$. The curve $h(\mathbf{P}^1)$ sweeps the boundary divisor of $\overline{\mathcal{A}}_6$ and intersects non-negatively any effective divisor on $\overline{\mathcal{A}}_6$ not containing D_6 . In particular,

$$s(\overline{\mathcal{A}}_6) \geq \frac{h(\mathbf{P}^1) \cdot [D_6]}{h(\mathbf{P}^1) \cdot \lambda_1}.$$