

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

AEPPLI COHOMOLOGY CLASSES ASSOCIATED WITH GAUDUCHON METRICS ON COMPACT COMPLEX MANIFOLDS

Dan Popovici

Tome 143

Fascicule 4

2015

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 763-800

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 4, tome 143, décembre 2015

Comité de rédaction

Gérard BESSON	Daniel HUYBRECHTS
Emmanuel BREUILLARD	Julien MARCHÉ
Antoine CHAMBERT-LOIR	Christophe SABOT
Charles FAVRE	Laure SAINT-RAYMOND
Pascal HUBERT	Wilhelm SCHLAG
Marc HERZLICH	

Raphaël KRIKORIAN (dir.)

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 176 €, hors Europe : 193 € (\$ 290)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© *Société Mathématique de France* 2015

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

AEPPLI COHOMOLOGY CLASSES ASSOCIATED WITH GAUDUCHON METRICS ON COMPACT COMPLEX MANIFOLDS

BY DAN POPOVICI

ABSTRACT. — We propose the study of a Monge-Ampère-type equation in bidegree $(n-1, n-1)$ rather than $(1, 1)$ on a compact complex manifold X of dimension n for which we prove ellipticity and uniqueness of the solution subject to positivity and normalization restrictions. Existence will hopefully be dealt with in future work. The aim is to construct a special Gauduchon metric uniquely associated with any Aeppli cohomology class of bidegree $(n-1, n-1)$ lying in the Gauduchon cone of X that we hereby introduce as a subset of the real Aeppli cohomology group of type $(n-1, n-1)$ and whose first properties we study. Two directions for applications of this new equation are envisaged: to moduli spaces of Calabi-Yau $\partial\bar{\partial}$ -manifolds and to a further study of the deformation properties of the Gauduchon cone beyond those given in this paper.

RÉSUMÉ (*Classes de cohomologie d'Aeppli associées à des métriques de Gauduchon sur les variétés complexes compactes*)

Nous proposons l'étude d'une équation de type Monge-Ampère en bidegré $(n-1, n-1)$ plutôt que $(1, 1)$ sur une variété complexe compacte X de dimension n pour laquelle nous démontrons l'ellipticité et l'unicité des solutions soumises à des contraintes de positivité et de normalisation. Nous espérons que la question de l'existence pourra être traitée dans un travail ultérieur. Le but est de construire une métrique de Gauduchon spéciale associée de manière unique à une classe de cohomologie d'Aeppli quelconque de bidegré $(n-1, n-1)$ appartenant au cône de Gauduchon de X que nous introduisons comme un sous-ensemble du groupe de cohomologie d'Aeppli réel de type $(n-1, n-1)$ et dont nous étudions les premières

Texte reçu le 4 février 2014, révisé le 20 juillet 2014, accepté le 8 octobre 2014.

DAN POPOVICI, Institut de Mathématiques de Toulouse, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse, France • *E-mail* : popovici@math.univ-toulouse.fr
2010 Mathematics Subject Classification. — 53C55, 32Q25, 32G05, 14C30.

Key words and phrases. — Positivity in bidegree $(n-1, n-1)$, Gauduchon and sG cones, duality between the Bott-Chern and the Aeppli cohomologies, equation of the Monge-Ampère type in bidegree $(n-1, n-1)$.

propriétés. Des applications de cette nouvelle équation sont envisagées dans deux directions : aux espaces de modules de $\partial\bar{\partial}$ -variétés de Calabi-Yau et à une étude des propriétés de déformations du cône de Gauduchon au-delà de celles décrites dans ce travail.

1. Introduction

Let X be a compact complex manifold, $\dim_{\mathbb{C}} X = n$. The main theme of this paper is the interaction between various kinds of metrics (especially Gauduchon metrics) on X and certain cohomology theories (especially the Aeppli cohomology) often considered on X .

On the metric side, let $\omega > 0$ be a C^∞ positive definite $(1, 1)$ -form (i.e., a Hermitian metric) on X . The following diagram sums up the definitions of well-known kinds of Hermitian metrics and the implications among them.

$$\begin{array}{lll}
 d\omega = 0 & \implies \exists \alpha^{0,2} \in C_{0,2}^\infty(X, \mathbb{C}) \text{ s.t.} & \implies \partial\bar{\partial}\omega = 0 \\
 & d(\overline{\alpha^{0,2}} + \omega + \alpha^{0,2}) = 0 & \\
 (\omega \text{ is Kähler}) & (\omega \text{ is Hermitian-symplectic}) & (\omega \text{ is pluriclosed}) \\
 \Downarrow & & \\
 d\omega^{n-1} = 0 & \implies \exists \Omega^{n-2,n} \in C_{n-2,n}^\infty(X, \mathbb{C}) \text{ s.t.} & \implies \partial\bar{\partial}\omega^{n-1} = 0 \\
 & d(\overline{\Omega^{n-2,n}} + \omega^{n-1} + \Omega^{n-2,n}) = 0 & \\
 (\omega \text{ is balanced}) & (\omega \text{ is strongly Gauduchon (sG)}) & (\omega \text{ is Gauduchon}).
 \end{array}$$

Recall that of the above six types of metrics, only Gauduchon metrics always exist ([7]). Compact complex manifolds X carrying any of the other five types of metrics inherit the name of the metric (Kähler, balanced, etc).

The conditions on the top line in the above diagram bear on the metric in bidegree $(1, 1)$, while those on the bottom line are their analogues in bidegree $(n - 1, n - 1)$. It is a well-known fact in linear algebra (see e.g., [11]) that for every smooth $(n - 1, n - 1)$ -form $\Gamma > 0$ on X , there exists a unique smooth $(1, 1)$ -form $\gamma > 0$ on X such that $\gamma^{n-1} = \Gamma$. We denote it $\gamma = \Gamma^{\frac{1}{n-1}}$ and call γ the $(n - 1)$ st root of Γ . The power-root bijection between positive definite C^∞ forms of types $(1, 1)$ and $(n - 1, n - 1)$ suggests a possible duality between the metric properties in these two bidegrees. The following observation (noticed before in [8] and references therein as a consequence of more general results) gives a further hint. We give below a quick proof.

PROPOSITION 1.1. — *If ω is both pluriclosed and balanced, then ω is Kähler.*

Proof. — The pluriclosed assumption on ω translates to any of the following equivalent properties:

$$(1) \quad \partial\bar{\partial}\omega = 0 \iff \partial\omega \in \ker \bar{\partial} \iff \star(\partial\omega) \in \ker \partial^*,$$

the last equivalence following from the well known formula $\partial^* = -\star\bar{\partial}\star$, where $\star = \star_\omega : \Lambda^{p,q}T^*X \rightarrow \Lambda^{n-q,n-p}T^*X$ is the Hodge-star isomorphism defined by ω for arbitrary $p, q = 0, \dots, n$.

On the other hand, the balanced assumption on ω translates to any of the following equivalent properties:

$$(2) \quad d\omega^{n-1} = 0 \iff \partial\omega^{n-1} = 0 \iff \partial\omega \text{ is primitive,}$$

the last equivalence following from the formula $\partial\omega^{n-1} = (n-1)\omega^{n-2} \wedge \partial\omega$. Now, since $\partial\omega$ is primitive by (2), a well-known formula (valid for any Hermitian metric ω) given e.g., in [19, Proposition 6.29, p. 150] spells

$$\star(\partial\omega) = i \frac{\omega^{n-3}}{(n-3)!} \wedge \partial\omega = \frac{i}{(n-2)!} \partial\omega^{n-2}.$$

Since $\star(\partial\omega) \in \ker \partial^*$ by (1), we get by applying ∂^* in the above identities:

$$\partial^*\partial\omega^{n-2} = 0, \quad \text{so} \quad 0 = \langle\langle \partial^*\partial\omega^{n-2}, \omega^{n-2} \rangle\rangle = \|\partial\omega^{n-2}\|^2, \quad \text{so} \quad \partial\omega^{n-2} = 0.$$

Since $\partial\omega^{n-2} = (n-2)\omega^{n-3} \wedge \partial\omega$, the last identity above gives $\omega^{n-3} \wedge \partial\omega = 0$. Now, $\partial\omega$ is a form of degree 3 while the Lefschetz operator on 3-forms

$$L_\omega^{n-3} : \Lambda^3 T^*X \longrightarrow \Lambda^{2n-3} T^*X, \quad \alpha \mapsto \omega^{n-3} \wedge \alpha,$$

is an isomorphism (see e.g., [19, Lemma 6.20, p. 146]—no assumption on the Hermitian metric ω is needed). It follows that $\partial\omega = 0$, i.e., ω is Kähler. \square

On the cohomological side, recall that for all $p, q = 0, \dots, n$, the Bott-Chern cohomology group of type (p, q) is defined as

$$H_{BC}^{p,q}(X, \mathbb{C}) = \frac{\ker(d : C_{p,q}^\infty(X) \rightarrow C_{p+1,q}^\infty(X) + C_{p,q+1}^\infty(X))}{\text{Im}(\partial\bar{\partial} : C_{p-1,q-1}^\infty(X) \rightarrow C_{p,q}^\infty(X))},$$

while the Aeppli cohomology group of type (p, q) is defined as

$$H_A^{p,q}(X, \mathbb{C}) = \frac{\ker(\partial\bar{\partial} : C_{p,q}^\infty(X) \rightarrow C_{p+1,q+1}^\infty(X))}{\text{Im}(\partial : C_{p-1,q}^\infty(X) \rightarrow C_{p,q}^\infty(X)) + \text{Im}(\bar{\partial} : C_{p,q-1}^\infty(X) \rightarrow C_{p,q}^\infty(X))}.$$

There are always well-defined linear maps from $H_{BC}^{p,q}(X, \mathbb{C})$, from $H_{\bar{\partial}}^{p,q}(X, \mathbb{C})$ (the Dolbeault cohomology group of type (p, q)) and from $H^{p+q}(X, \mathbb{C})$ (the De Rham cohomology group of degree $p+q$) to $H_A^{p,q}(X, \mathbb{C})$ but, in general, they are neither injective, nor surjective.

We will be often considering the case when X is a $\partial\bar{\partial}$ -manifold. This means that the $\partial\bar{\partial}$ -lemma holds on X , i.e., for all p, q and for any smooth d -closed