

Mémoires

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Numéro 158
Nouvelle série

**POISSON ENSEMBLES
OF LOOPS
OF ONE-DIMENSIONAL
DIFFUSIONS**

2 0 1 8

Titus LUPU

SOCIÉTÉ MATHÉMATIQUE DE FRANCE
Publié avec le concours du Centre National de la Recherche Scientifique

Comité de rédaction

Christine BACHOC
Yann BUGEAUD
Jean-François DAT
Pascal HUBERT
Laurent MANIVEL

Marc HERZLICH (dir.)

Julien MARCHÉ
Kieran O'GRADY
Emmanuel RUSS
Christine SABOT

Diffusion

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 9
France
commandes@smf.emath.fr

AMS
P.O. Box 6248
Providence RI 02940
USA
www.ams.org

Tarifs

Vente au numéro : 32 € (\$ 48)
Abonnement électronique : 113 € (\$ 170)
Abonnement avec supplément papier : 167 €, hors Europe : 197 € (\$ 296)
Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat

Mémoires de la SMF
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris Cedex 05, France
Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96
memoires@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2018

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0249-633-X

ISBN 978-2-85629-891-6

[doi:10.24033/msmf.466](https://doi.org/10.24033/msmf.466)

Directeur de la publication : Stéphane SEURET

POISSON ENSEMBLES OF LOOPS
OF ONE-DIMENSIONAL DIFFUSIONS

Titus Lupu

T. Lupu

Laboratoire de Probabilités, Statistique et Modélisation, Sorbonne Université, 75005
Paris (France).

E-mail : titus.lupu@upmc.fr

2000 Mathematics Subject Classification. – 60-02, 60G15, 60G17, 60G55, 60G60,
60J55, 60J60, 60J65, 60J80.

Key words and phrases. – Poisson ensembles of Markov loops, loop-soup, one-dimensional diffusion processes, Vervaat's transformation, continuous state branching processes with immigration, Gaussian free field, Dynkin's isomorphism, Wilson's algorithm, uniform spanning tree, determinantal point processes.

POISSON ENSEMBLES OF LOOPS OF ONE-DIMENSIONAL DIFFUSIONS

Titus Lupu

Abstract. – There is a natural measure on loops (time-parametrized trajectories that in the end return to the origin), which one can associate to a wide class of Markov processes. The Poisson ensembles of Markov loops are Poisson point processes with intensity proportional to these measures. In wide generality, these Poisson ensembles of Markov loops are related, at intensity parameter $1/2$, to the Gaussian free field, and at intensity parameter 1 , to the loops done by a Markovian sample path. Here, we study the specific case when the Markov process is a one-dimensional diffusion. After a detailed description of the measure, we study the Poisson point processes of loops, their occupation fields, and explain how to sample these Poisson ensembles of loops out of diffusion sample path perturbed at their successive minima. Finally, we introduce a couple of interwoven determinantal point processes on the line, which is a dual through Wilson’s algorithm of Poisson ensembles of loops, and study the properties of these determinantal point processes.

Résumé (Ensembles poissoniens de boucles des diffusions unidimensionnelles)

Il y a une mesure naturelle sur les boucles (trajectoires paramétrées par le temps, qui à la fin retournent à leur origine) qu'on peut associer à une large classe de processus de Markov. Les ensembles poissoniens de boucles markoviennes sont des processus ponctuels de Poisson d'intensité proportionnelle à ces mesures. Dans une grande généralité, ces ensembles poissoniens de boucles markoviennes sont reliés, au paramètre d'intensité $1/2$, au champ libre gaussien, et au paramètre d'intensité 1 , aux boucles créées par une trajectoire markovienne. Ici nous étudions le cas spécifique où le processus de Markov est une diffusion unidimensionnelle. Après une description détaillée de la mesure, nous étudions les processus ponctuels de Poisson des boucles, leurs champs d'occupation et expliquons comment séquencer ces ensembles poissoniens de boucles à partir de trajectoires de diffusions perturbées à leur minima successifs. Enfin, nous introduisons un couple de processus ponctuels déterminantaux sur la droite, entrelacés, qui est un dual, à travers l'algorithme de Wilson, de l'ensemble poissonien de boucles, et étudions les propriétés de ces processus ponctuels déterminantaux.

CONTENTS

1. Introduction	1
1.1. Measure on loops: history	1
1.2. Measure on loops: present work and recent developments	2
1.3. Results and layout	4
1.4. A list of commonly used notations	6
Chapter 2	6
Chapter 3	7
Chapter 4	8
Chapter 5	9
Chapter 6	9
Chapter 7	10
2. Preliminaries on generators and semi-groups	11
2.1. A second order ODE	11
2.2. One-dimensional diffusions	17
2.3. “Generators” with creation of mass	21
3. Measure on loops and its basic properties	29
3.1. Spaces of loops	29
3.2. Measures $\mu^{x,y}$ on finite life-time paths	30
3.3. The measure μ^* on unrooted loops	37
3.4. Multiple local times	41
3.5. Measure on loops rooted at the minimum	44
3.6. A generalization of the Vervaat’s transformation	48
3.7. Restricting loops to a discrete subset	53
3.8. Measure on loops in case of creation of mass	55
4. Occupation fields of the Poisson ensembles of Markov loops	59
4.1. Inhomogeneous continuous state branching processes with immigration ..	59
4.2. Occupation field	62
4.3. Isomorphism with the Gaussian free field	72

5. Decomposing paths into Poisson ensembles of loops	77
5.1. Gluing together excursions ordered by their minima	77
5.2. Loops represented as excursions and glued together	80
5.3. The case $\alpha = 1$	87
6. Wilson's algorithm in dimension one	93
6.1. Description of the algorithm	93
6.2. The erased paths	95
6.3. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: Brownian case	97
6.4. Determinantal point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$: general case	118
7. Monotone couplings for the point processes $(\mathcal{Y}_\infty, \mathcal{Z}_\infty)$	119
7.1. Conditioning	119
7.2. Couplings	134
Bibliography	153

CHAPTER 1

INTRODUCTION

1.1. Measure on loops: history

There is a measure on loops one can naturally associate to a wide class of Markov processes. In general, it is expressed as

$$(1.1.1) \quad \mu(d\gamma) := \int_{t>0} \int_x \mathbb{P}_{x,x}^t(d\gamma) p_t(x,x) m(dx) \frac{dt}{t},$$

where $p_t(x,y)$ are the transition densities with respect to a σ -finite measure $m(dy)$, and $\mathbb{P}_{x,y}^t$ are the bridge probability measures. Here, a loop γ is a path parametrized by continuous time, which at the end returns to the origin ($\gamma(0) = \gamma(T(\gamma))$).

Such a measure first appeared, to my knowledge, in the work of Symanzik [61, 62, 63]. There, studying the ϕ^4 model in dimension d , he expressed the moments, and some other functionals, of such fields in terms of multiple integrals over measures on massive Brownian loops and massive Brownian paths, corresponding to a d -dimensional Brownian motion killed at independent exponential time. He called this loop expansion, and saw it as Euclidean analog of Feynman diagrams in Minkowski space QFT. Later, in the work of Brydges, Frölich and Spencer [9], an analogous measure appeared, but for discrete time random walk loops.

Then, an analogously defined measure in the setting of the two-dimensional Brownian motion appeared in the work of Lawler and Werner [34]. They also considered the Poisson point processes of intensity proportional to this measure on loops, which they called “loop-soups”. The main motivation for studying this object was that the obtained loops, up to rerooting and time-reparametrization, were invariant in law by conformal transformations. This properties were used by Sheffield and Werner in [57] to construct the Conformal Loop Ensemble (CLE) as outer boundaries of clusters in a Brownian loop-soup.

Lawler and Trujillo-Ferreras initiated in [33] the study of discrete time random walk loop-soups. See also a recent survey by Lawler [31]. The measure they used was actually the same that appeared in Brydges, Frölich and Spencer [9]. Le Jan

considered loops parametrized by continuous time, associated to Markov jump process on an electrical network, with intensity following the pattern (1.1.1), rather than the discrete time random walk loops [36]. He also called the object Poisson ensemble of Markov loops rather than loop-soup. We will adopt Le Jan's terminology. If one takes the discrete skeleton of Le Jan's loops, one gets the random walk loops studied in [33].

Taking loops parametrized by continuous time allowed Le Jan to consider the occupation field of Poisson ensembles of Markov loops, that is to say the total time spent by loops on each vertex. Le Jan identified two properties universally satisfied by these Poisson ensembles of Markov loops. The first one, is that at intensity parameter $1/2$, the occupation field has same law as half the square of a Gaussian free field. This is an extension of Dynkin's isomorphism [16, 18, 17], and is closely related to Symanzik's identities [63]. The second universal property is that, at intensity parameter 1 , the loops erased during the loop-erasure algorithm applied to a Markovian sample path, are part of a Poisson ensembles of Markov loops. This property has been previously observed in the two-dimensional Brownian setting. See [34], Conjecture 1, and [32], Theorem 7.3. See also [56] for recent developments in dimension three. On an electrical network, Wilson's algorithm [69] samples a uniform spanning tree by running successive loop erased random walks. Le Jan observed that the loops erased during this algorithm give a Poisson ensembles of Markov loops of intensity parameter 1 .

Since then, the definition of the measure of loops was extended to a wide class of Markov processes, including some not having transition densities [38, 25, 23]. Actually, one rather considers measures on unrooted loops, seen as parametrized by a circle, where the cut between the start and the endpoint is not identified. What makes such measures on unrooted loops natural is actually the covariance by time change by an inverse of a Continuous Additive Functional (CAF). The time change by an inverse of a CAF transforms a Markov process into another one, possibly on a smaller state space. The measure on unrooted loops of the time changed process is the pushforward by time-change of the measure on loops of the initial Markov process. This covariance is no longer true at the level of rooted loops. In dimension two for Brownian loops, the covariance by time-change implies the invariance by conformal transformations of the range [34].

1.2. Measure on loops: present work and recent developments

These notes are devoted to the study of loop measures, Poisson ensembles of loops and of an analog of Wilson's algorithm in the setting of one-dimensional diffusions. This is a work done during my Ph.D. under the direction of Le Jan at Université Paris-Sud, Orsay [39]. The aim of these notes is to give a complete and self-contained presentation of the topic in the continuous one-dimensional setting, displaying both