Astérisque

# BURKHARD KÜLSHAMMER Morita equivalent blocks in Clifford theory of finite groups

Astérisque, tome 181-182 (1990), p. 209-215

<http://www.numdam.org/item?id=AST\_1990\_\_181-182\_\_209\_0>

© Société mathématique de France, 1990, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (http://smf4.emath.fr/ Publications/Asterisque/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

# $\mathcal{N}$ umdam

Article numérisé dans le cadre du programme Numérisation de documents anciens mathématiques http://www.numdam.org/

## Morita Equivalent Blocks in Clifford Theory of Finite Groups

BURKHARD KÜLSHAMMER

Let F be an algebraically closed field of prime characteristic p, and let

 $1 \longrightarrow K \longrightarrow H \longrightarrow G \longrightarrow 1$ 

be an extension of finite groups. Let B be a block of FK (considered as a subalgebra of FK), and let A be a block of FH covering B (i. e.  $t_A t_B \neq 0$ ). Following a suggestion by J. L. Alperin [1] we consider the following

**QUESTION.** When are A and B Morita equivalent?

Our main results concerning this question are given by theorems 1, 7, 8 and proposition 10 below. Special cases of this question are dealt with in [2] and [7].

**THEOREM 1.** With notation as above, the map  $B \longrightarrow 1_A B \subset A$ ,  $b \longmapsto 1_A b$ , is an isomorphism of F-algebras.

Before proving theorem 1 we introduce some notation and state some preliminary results. Obviously K is contained in  $H(B) := \{h \in H: hBh^{-1} = B\}$ , the stabilizer of B in H, and we set G(B) := H(B)/K. The following facts are well-known (see [8; theorem 1], for example).

**PROPOSITION 2.** (i)  $FH1_BFH$  is the sum of all blocks of FH covering B. (ii) If  $h_p$ , ...,  $h_t$  denote a transversal for H(B) in H then the map

$$\operatorname{Mat}(t, I_{B}FH(B)) \longrightarrow FHI_{B}FH, \ [a_{ij}]_{i,j=1}^{t} \longrightarrow \sum_{i,j=1}^{t} h_{i}a_{ij}h_{j}^{-1},$$

S.M.F. Astérisque 181-182 (1990)

### B. KÜLSHAMMER

is an isomorphism of F-algebras.

(iii) The maps

$$Z(1_BFH(B)) \longrightarrow Z(FH1_BFH), z \longmapsto \sum_{i=1}^{t} h_i z h_i^{-1}$$

and

$$Z(FH1_{B}FH) \longrightarrow Z(1_{B}FH(B)), z \longmapsto 1_{B}z,$$

are isomorphisms of F-algebras and inverse to each other.

For  $h \in H(B)$ , the map  $B \longrightarrow B$ ,  $b \longmapsto hbh^{-1}$ , is an *F*-algebra automorphism of *B*. It is easy to see that the elements  $h \in H$  for which the map  $B \longrightarrow B$ ,  $b \longmapsto hbh^{-1}$ , is an inner automorphism of *B* form a normal subgroup H(B) of H(B) containing K (cf. [3; proposition 2.7]). Define G(B) := H(B)/K.

Setting  $C := 1_B C_{FH}(K)$  and  $C_g := C \cap hFK$  for  $g = hK \in G$  we obtain  $C = \bigoplus_{g \in G} C_g$ and  $C_g C_{g'} \in C_{gg'}$  for  $g,g' \in G$ , i. e. C is a G-graded F-algebra in the sense of [4]. It is easy to see that  $C_g = 0$  for  $g \in G \setminus G(B)$ . Thus  $C = \bigoplus_{g \in G(B)} C_g$  can also be viewed as a G(B)-graded F-algebra.

**PROPOSITION 3.** (13; lemma 3.31)  $I := \bigoplus_{g \in G[B]} (JZB)C_g \oplus \bigoplus_{g \in G(B) \setminus G[B]} C_g$  is an ideal of C contained in the radical JC of C.

Setting  $C[B] := \bigoplus_{g \in G[B]} C_g$  we thus have C = C[B] + JC. By lifting theorems for idempotents one obtains the following result.

**COROLLARY 4.** ([3; theorem 3.5]) All idempotents of ZC are contained in C[B].

It is easy to see that C[B] is a crossed product of G[B] with ZB, in the sense of [4]; in particular, C[B] is free as a ZB-module, and  $\overline{C[B]} := C[B]/(JZB)C[B]$  is a crossed product of G[B] with  $ZB/(JZB) \cong F$ , i. e. a twisted group algebra of G[B] over F. Our next result is [8; theorem C].

#### MORITA EQUIVALENT BLOCKS

**PROPOSITION 5.** If G = G[B] then the map  $B \otimes_{ZB} C \longrightarrow 1_B FH$ ,  $b \otimes c \longmapsto bc$ , is an isomorphism of F-algebras.

We are now in a position to prove theorem 1.

**Proof of theorem 1.** Obviously the map  $B \longrightarrow 1_A B$ ,  $b \longmapsto 1_A b$ , is an epimorphism of *F*-algebras. Hence it suffices to prove injectivity. By proposition 2,  $1_A 1_B$  is the block idempotent of a block of *FH(B)* covering *B*. Hence we may replace *H* by *H(B)* and assume H = H(B). By corollary 4,  $1_A$  is contained in *FH(B)*. Replacing *A* by a block of  $1_A FH(B)$  we may assume that H = H(B). In this case the map  $B \otimes_{ZB} C \longrightarrow 1_B FH$ ,  $b \otimes c \longmapsto bc$ , is an isomorphism of *F*-algebras by proposition 5. Moreover, *C* is free over *ZB*. This isomorphism maps  $B \otimes_{ZB} 1_A C$  onto *A*. Since  $C = 1_A C \oplus (1_B - 1_A)C$ ,  $1_A C$ is projective over *ZB*. Since *ZB* is local,  $1_A C$  is even free over *ZB*. Thus *A* is free over *B*, and the result follows.  $\boxtimes$ 

In order to prove our next theorem we need a result on the behaviour of defect groups.

**PROPOSITION 6.** (13; theorem 7.7.1)  $1_A + (JZB)C[B]$  is a primitive idempotent in  $C_{\overline{C(B)}}(G(B))$ , and A has a defect group P such that  $P \cap K$  is a defect group of B and PK/K is a defect group of  $1_A + (JZB)C[B]$  in G(B).

Part of proposition 6 has also been proved in [6; 4.2]. We will say that A and B are "naturally" Morita equivalent of degree n if there exists a simple F-subalgebra S of A of dimension  $n^2$  such that the map  $l_A B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of F-algebras. In this case A and B are Morita equivalent since  $l_A B$  is isomorphic to B by theorem 1 and S is a complete matrix algebra of degree n over F.

**THEOREM 7.** A and B are "naturally" Morita equivalent if and only if G = G[B] and A and B have the same defect.

**Proof.** Suppose first that G = G[B] and that A and B have the same defect. By proposition 6, the block  $I_AC + (JZB)C/(JZB)C$  of the twisted group algebra C/(JZB)C of G[B] = G over F has defect 0 in G(B) = G. It is well-known that this implies that

### B. KÜLSHAMMER

the block  $I_AC + (JZB)C/(JZB)C$  of C/(JZB)C is a simple *F*-algebra; in particular,  $I_AJC = (JZB)I_AC$ . By the Wedderburn-Malcev theorem there is a simple *F*-subalgebra *S* of  $I_AC$  such that  $I_AC = S \oplus I_AJC = S \oplus (JZB)I_AC$ . Then  $I_AC = (ZB)S + (JZB)I_AC$ , and Nakayama's lemma implies that  $I_AC = (ZB)S$ . In the proof of theorem 1 we had shown that  $I_AC$  is free over *ZB*. Thus  $I_AC/I_AJC$  is free of the same rank over *ZB/JZB*  $\cong$  *F*. Therefore the rank of  $I_AC$  over *ZB* equals the dimension of *S* over *F*. Comparing dimensions we see that the map  $ZB \otimes_F S \longrightarrow I_AC$ ,  $z \otimes s \longmapsto zs$ , is an isomorphism of *F*-algebras. By proposition 5, the map  $B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism as well.

Suppose now conversely that A and B are "naturally" Morita equivalent, and let S be a simple F-subalgebra of A such that the map  $1_A B \otimes_F S \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of F-algebras. Then  $1_A = 1_S = 1_A 1_B$ . On the other hand, it follows from proposition 2 that  $1_A = \sum_{i=1}^{t} 1_A (h_i 1_B h_i^{-1})$  with pairwise orthogonal idempotents  $1_A (h_i 1_B h_i^{-1})$  where t = |H:H(B)|. Thus H(B) = H and G(B) = G.

We know from proposition 3 that C = C[B] + JC; in particular,  $I_A C = I_A C[B] + I_A JC$ . On the other hand, since A and B are "naturally" Morita equivalent the map

$$1_A ZB \otimes_F S \longrightarrow 1_A ZB \cdot S = C_A(B) = 1_A C, \ z \otimes s \longmapsto zs,$$

is an isomorphism of *F*-algebras. By the Wedderburn-Malcev theorem we may find a unit *u* in  $1_AC$  such that  $S^u$  is contained in  $1_AC[B]$ . Then the map  $1_AB \otimes_F S^u \longrightarrow A$ ,  $b \otimes s \longmapsto bs$ , is an isomorphism of *F*-algebras as well. Hence we may assume that *S* is contained in *FH[B]*. Since also  $1_A \in FH[B]$  by corollary 4 we obtain  $A \subset FH[B]$  which clearly implies that H[B] = H.

Since  $I_AC$  is isomorphic to  $ZB \otimes_F S$ .  $I_AC + (JZB)C/(JZB)C$  is a simple F-algebra. It is well-known that this implies that the block  $I_AC + (JZB)C/(JZB)C$  of  $\overline{C(B)}$  has defect 0 in G[B] = G. By proposition 6, A and B have the same defect.  $\square$ 

In the following we assume that G(B) = G; in view of proposition 2, this is not an important restriction. In this case we can reduce the question of whether A and B are "naturally" Morita equivalent to their Brauer correspondents. Let Q be a defect group of B, and let B' be the Brauer correspondent of B in  $N_K(Q)$ . Since G(B) = G the Frattini argument shows that  $H = N_H(Q)K$ , and we obtain a finite group extension