Astérisque

JOHAN PAS Some applications of uniform *p*-adic cell decomposition

Astérisque, tome 198-199-200 (1991), p. 265-271

<http://www.numdam.org/item?id=AST_1991__198-199-200__265_0>

© Société mathématique de France, 1991, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (http://smf4.emath.fr/ Publications/Asterisque/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

\mathcal{N} umdam

Article numérisé dans le cadre du programme Numérisation de documents anciens mathématiques http://www.numdam.org/

SOME APPLICATIONS OF UNIFORM *p*-ADIC CELL DECOMPOSITION

JOHAN PAS¹

K.U. Leuven

In this paper we summarize some applications of uniform p-adic cell decomposition. The technique of p-adic cell decomposition was developed by Denef [3,4], using ideas of Cohen [2]. Denef gave two applications of his theorem. A first one was an elementary proof of Macintyre's quantifier elimination for p-adic fields; a second one was a proof of the rationality of the Igusa local zeta function without using Hironaka's resolution of singularities.

Denef's cell decomposition theorem holds for every *p*-adic field but the decomposition procedure depends on the particular field. In [13] and [14] we obtain cell decomposition theorems which are uniform in a certain class of *p*-adic fields. In [13] we prove a cell decomposition for a class of henselian valued fields of equicharacteristic zero. By the use of an ultraproduct construction, this provides a decomposition which is uniform, for almost all *p*, in the class $\{\mathbf{Q}_p\}_{p \text{ prime}}$ of all fields of *p*-adic numbers. The cell decomposition proved in [14] is uniform in the class of all finite unramified extensions of the field \mathbf{Q}_p (with *p* a fixed prime number). Macintyre [10] obtained independently a different cell decomposition which is also uniform in these classes of fields.

The basic idea of cell decomposition is that, given a *p*-adic field Kand a polynomial f(x) with $x = (x_1, \ldots, x_m)$, one can partition K^m into

¹Research Assistant of the National Fund of Scientific Research (Belgium)

so-called cells, such that on each of these cells the function f(x) takes, in a certain sense, a simpler form. The precise meaning of 'simpler' here is determined by the formalism (defined by a first order language for *p*-adic fields) in which one proves the cell decomposition theorem. The use of uniform cell decomposition allows one to simplify, uniformly, formulas in the first order language and to obtain a quantifier elimination in this language which is uniform in the class of fields. This is formulated more precisely in section 1. A second application of cell decomposition is the computation of certain *p*-adic integrals on definable sets. Cell decomposition enables one to partition definable sets in simpler parts for which the integral can be computed. The results here are stated in section 2.

Section 1 Quantifier elimination

We define the first order language \mathcal{L} , in which we prove the cell decomposition for valued fields of equicharacteristic zero, and in which we obtain quantifier elimination. The language \mathcal{L} is a language with three sorts of variables, namely variables for the elements of the valued field, variables for elements of the residue field and variables for elements of the value group. The language contains symbols for the standard field operations in the valued field and in the residue field, and symbols for the usual operations in the value group. One also has a function symbol for the valuation map from the field to the value group, and another function symbol for an angular component map modulo P from the field to the residue field. Such an angular component map modulo P is a multiplicative morphism from the group of units of the valued field to the group of units of the residue field, such that the restriction of this morphism to the set of elements of valuation zero is the canonical projection onto the residue field.

Cell Decomposition Theorem 3.2 of [13] implies

THEOREM. ([13] Theorem 4.1) Let K be a henselian valued field of equicharacteristic zero which has an angular component map modulo P. Then K has elimination of field quantifiers in the language \mathcal{L} .

Applying this to ultraproducts $\prod \mathbf{Q}_p / \mathcal{D}$ where \mathcal{D} is a non-principal ultrafilter on the index set of prime numbers we obtain a uniform quantifier elimination for the fields of *p*-adic numbers.

COROLLARY. ([13] Corollary 4.3) Let φ be a formula in \mathcal{L} . Then there exists an \mathcal{L} -formula ψ (which is independent of p) without field quantifiers, such that, for almost all primes p, we have

 $\varphi \longleftrightarrow \psi$ on \mathbf{Q}_p .

Section 2 Poincaré series and Igusa local zeta functions

Let h(x) be a polynomial in m variables $x = (x_1, \ldots, x_m)$ over \mathbb{Z} , and let K be a finite field extension of \mathbb{Q}_p for some prime number p, such that K is unramified over \mathbb{Q}_p , that is, p is a generator for the maximal ideal of the valuation ring R of K. Suppose that the residue field of Khas cardinality q. The valuation on K is denoted by |.| and normalized by |p| = 1/q.

For every $n \in \mathbb{N}$ we consider the number of solutions of the congruence $h(x) \equiv 0 \mod p^n$ in R,

$$N(n,K) = \operatorname{Card} \{ x \mod p^n \mid x \in R^m, h(x) \equiv 0 \mod p^n \}.$$

The generating function of this sequence

$$P(T,K) = \sum_{n=0}^{\infty} N(n,K) T^n$$

is called a *Poincaré series* associated to the polynomial h. Borevich and Shafarevich [1] conjectured that this series is a rational function of T. This conjecture was first proved by Igusa [8,7] using Hironaka's resolution of singularities.

Another type of Poincaré series for h is the series associated to the *p*-adic points on the variety h = 0. Put for $n \in \mathbb{N}$,

$$\widetilde{N}(n,K) = \operatorname{Card} \{ x \mod p^n \mid x \in R^m, h(x) = 0 \},$$

 \mathbf{and}

$$\widetilde{P}(T,K) = \sum_{n=0}^{\infty} \widetilde{N}(n,K) T^{n}.$$

This series was studied by Oesterlé [12] and Serre [16]. Denef [3] proved that $\tilde{P}(T, K)$ is a rational function of T.

A first step in the rationality proofs for these Poincaré series, is to relate the series to a certain *p*-adic integral. For the first series P(T, K)this integral is the so-called *Igusa local zeta function* of *h*, which is defined as

(1)
$$Z(s,K) = \int_{R^m} |h(x)|^s |dx| \quad \text{for } s \in \mathbf{R}, \, s > 0,$$

where |dx| is the Haar measure on K^m normalized on R^m . The relation of this integral with the Poincaré series P(T, K) is given by

(2)
$$Z(s,K) = \frac{1 + (T-1)P(q^{-m}T)}{T}$$
 for $T = q^{-s}$.

For $\widetilde{P}(T, K)$ the corresponding integral is

$$I(s,K) = \int_{D} |w|^{s} |dx| |dw| \quad \text{for } s \in \mathbf{R}, \, s > 0,$$

where

$$D = \{ (x, w) \in \mathbb{R}^m \times \mathbb{R} \mid \exists y \in \mathbb{R}^m : x \equiv y \bmod w \text{ and } h(y) = 0 \}.$$