

*quatrième série - tome 49    fascicule 3    mai-juin 2016*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Vincent SÉCHERRE & Shaun STEVENS

*Block decomposition of the category of  $\ell$ -modular smooth representations  
of  $GL_n(\mathbb{F})$  and its inner forms*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> janvier 2016

N. ANANTHARAMAN I. GALLAGHER  
P. BERNARD B. KLEINER  
E. BREUILLARD E. KOWALSKI  
R. CERF M. MUSTAȚĂ  
A. CHAMBERT-LOIR L. SALOFF-COSTE

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

### Édition / *Publication*

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

### Abonnements / *Subscriptions*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

### Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

---

© 2016 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret  
Périodicité : 6 n<sup>os</sup> / an

# BLOCK DECOMPOSITION OF THE CATEGORY OF $\ell$ -MODULAR SMOOTH REPRESENTATIONS OF $GL_n(\mathbb{F})$ AND ITS INNER FORMS

BY VINCENT SÉCHERRE AND SHAUN STEVENS

---

**ABSTRACT.** – Let  $F$  be a nonarchimedean locally compact field of residue characteristic  $p$ , let  $D$  be a finite dimensional central division  $F$ -algebra and let  $R$  be an algebraically closed field of characteristic different from  $p$ . To any irreducible smooth representation of  $G = GL_m(D)$ ,  $m \geq 1$  with coefficients in  $R$ , we can attach a uniquely determined inertial class of supercuspidal pairs of  $G$ . This provides us with a partition of the set of all isomorphism classes of irreducible representations of  $G$ . We write  $\mathcal{R}(G)$  for the category of all smooth representations of  $G$  with coefficients in  $R$ . To any inertial class  $\Omega$  of supercuspidal pairs of  $G$ , we can attach the subcategory  $\mathcal{R}(\Omega)$  made of smooth representations all of whose irreducible subquotients are in the subset determined by this inertial class. We prove that the category  $\mathcal{R}(G)$  decomposes into the product of the  $\mathcal{R}(\Omega)$ 's, where  $\Omega$  ranges over all possible inertial class of supercuspidal pairs of  $G$ , and that each summand  $\mathcal{R}(\Omega)$  is indecomposable.

**RÉSUMÉ.** – Soit  $F$  un corps commutatif localement compact non archimédien de caractéristique résiduelle  $p$ , soit  $D$  une  $F$ -algèbre à division centrale de dimension finie et soit  $R$  un corps algébriquement clos de caractéristique différente de  $p$ . A toute représentation lisse irréductible du groupe  $G = GL_m(D)$ ,  $m \geq 1$  à coefficients dans  $R$  correspond une classe d'inertie de paires supercuspidales de  $G$ . Ceci définit une partition de l'ensemble des classes d'isomorphisme de représentations irréductibles de  $G$ . Notons  $\mathcal{R}(G)$  la catégorie des représentations lisses de  $G$  à coefficients dans  $R$  et, pour toute classe d'inertie  $\Omega$  de paires supercuspidales de  $G$ , notons  $\mathcal{R}(\Omega)$  la sous-catégorie formée des représentations lisses dont tous les sous-quotients irréductibles appartiennent au sous-ensemble déterminé par cette classe d'inertie. Nous prouvons que  $\mathcal{R}(G)$  est le produit des  $\mathcal{R}(\Omega)$ , où  $\Omega$  décrit les classes d'inertie de paires supercuspidales de  $G$ , et que chaque facteur  $\mathcal{R}(\Omega)$  est indécomposable.

## Introduction

When considering a category of representations of some group or algebra, a natural step is to attempt to decompose the category into *blocks*; that is, into subcategories which are indecomposable summands. Thus any representation can be decomposed uniquely as a direct sum of pieces, one in each block; any morphism comes as a product of morphisms, one in

---

The second-named author is supported by EPSRC grant EP/H00534X/1.

each block; and this decomposition of the category is the finest decomposition for which these properties are satisfied. Then a full understanding of the category is equivalent to a full understanding of all of its blocks.

In the case of representations of a finite group  $G$ , over an algebraically closed field  $R$ , there is always a block decomposition. In the simplest case, when the characteristic of  $R$  is prime to the order of  $G$ , this is particularly straightforward: all representations are semisimple so each block consists of representations isomorphic to a direct sum of copies of a fixed irreducible representation. In the general case, there is a well-developed theory, beginning with the work of Brauer and Nesbitt, and understanding the block structure is a major endeavor.

Now suppose  $G$  is the group of rational points of a connected reductive algebraic group over a nonarchimedean locally compact field  $F$ , of residue characteristic  $p$ . When  $R$  has characteristic zero, a block decomposition of the category  $\mathcal{R}_R(G)$  of smooth  $R$ -representations of  $G$  was given by Bernstein [1], in terms of the classification of representations of  $G$  by their cuspidal support. Any irreducible representation  $\pi$  of  $G$  is a quotient of some (normalized) parabolically induced representation  $i_M^G \varrho$ , with  $\varrho$  a cuspidal irreducible representation of a Levi subgroup  $M$  of  $G$ ; the pair  $(M, \varrho)$  is determined up to  $G$ -conjugacy by  $\pi$  and is called its *cuspidal support*. Then each such pair  $(M, \varrho)$  determines a block, whose objects are those representations of  $G$  all of whose subquotients have cuspidal support  $(M, \varrho\chi)$ , for some unramified character  $\chi$  of  $M$ .

One important tool in proving this block decomposition is the equivalence of the following two properties of an irreducible  $R$ -representation  $\pi$  of  $G$ :

- $\pi$  is not a quotient of any properly parabolically induced representation; equivalently, all proper Jacquet modules of  $\pi$  are zero ( $\pi$  is *cuspidal*);
- $\pi$  is not a *sub*quotient of any properly parabolically induced representation  $i_M^G \varrho$  with  $\varrho$  an irreducible representation ( $\pi$  is *supercuspidal*).

When  $R$  is an algebraically closed field of positive characteristic different from  $p$  (the *modular* case), these properties are no longer equivalent and the methods used in the characteristic zero case cannot be applied. Instead, one can attempt to define the *supercuspidal support* of a smooth irreducible  $R$ -representation  $\pi$  of  $G$ : it is a pair  $(M, \varrho)$  consisting of an irreducible supercuspidal representation  $\varrho$  of a Levi subgroup  $M$  of  $G$  such that  $\pi$  is a *sub*quotient of  $i_M^G \varrho$ . However, for a general group  $G$ , it is not known whether the supercuspidal support of a representation is well-defined up to conjugacy; indeed, the analogous question for finite reductive groups of Lie type is also open.

In any case, one can define the notion of an *inertial supercuspidal class*  $\Omega = [M, \varrho]_G$ : it is the set of pairs  $(M', \varrho')$ , consisting of a Levi subgroup  $M'$  of  $G$  and a supercuspidal representation  $\varrho'$  of  $M'$ , which are  $G$ -conjugate to  $(M, \varrho\chi)$ , for some unramified character  $\chi$  of  $M$ . Given such a class  $\Omega$ , we denote by  $\mathcal{R}_R(\Omega)$  the full subcategory of  $\mathcal{R}_R(G)$  whose objects are those representations all of whose subquotients are isomorphic to a subquotient of  $i_{M'}^G \varrho'$ , for some  $(M', \varrho') \in \Omega$ .

The main purpose of this paper is then to prove the following result:

**THEOREM.** – *Let  $G$  be an inner form of  $\mathrm{GL}_n(\mathbb{F})$  and let  $R$  be an algebraically closed field of characteristic different from  $p$ . Then there is a block decomposition*

$$\mathcal{R}_R(G) = \prod_{\Omega} \mathcal{R}_R(\Omega),$$

where the product is taken over all inertial supercuspidal classes.

This theorem generalizes the Bernstein decomposition in the case that  $R$  has characteristic zero, and also a similar statement, for general  $R$ , stated by Vignéras [24] in the split case  $G = \mathrm{GL}_n(\mathbb{F})$ ; however, the authors were unable to follow all the steps in [24] so our proof is independent, even if some of the ideas come from there.

Our proof builds on work of Mínguez and the first author [16, 15], in which they give a classification of the irreducible  $R$ -representations of  $G$ , in terms of supercuspidal representations, and of the supercuspidal representations in terms of the theory of types. In particular, they prove that supercuspidal support is well-defined up to conjugacy, so that the irreducible objects in  $\mathcal{R}_R(\Omega)$  are precisely those with supercuspidal support in  $\Omega$ .

One question we do not address here is the structure of the blocks  $\mathcal{R}_R(\Omega)$ . Given the explicit results on supertypes here, it is not hard to construct a progenerator  $\Pi$  for  $\mathcal{R}_R(\Omega)$  as a compactly-induced representation; for  $G = \mathrm{GL}_n(\mathbb{F})$  this was done (independently) by Guiraud [11] (for level zero blocks) and Helm [12]. Then  $\mathcal{R}_R(\Omega)$  is equivalent to the category of  $\mathrm{End}_G(\Pi)$ -modules. In the case that  $R$  has characteristic zero, the algebra  $\mathrm{End}_G(\Pi)$  was described as a tensor product of affine Hecke algebras of type A in [22] (or [7] in the split case); indeed, we use this description in our proof here. For  $R$  an algebraic closure  $\overline{\mathbb{F}}_{\ell}$  of a finite field of characteristic  $\ell \neq p$ , and a block  $\mathcal{R}_R(\Omega)$  with  $\Omega = [\mathrm{GL}_n(\mathbb{F}), \varrho]_{\mathrm{GL}_n(\mathbb{F})}$ , Dat [9] has described this algebra; it is an algebra of Laurent polynomials in one variable over the  $R$ -group algebra of a cyclic  $\ell$ -group. It would be interesting to obtain a description in the general case.

We now describe the proof of the theorem, which relies substantially on the theory of semisimple types developed in [22] (see [7] for the split case). Given an inner form  $G$  of  $\mathrm{GL}_n(\mathbb{F})$ , in [22] the authors constructed a family of pairs  $(\mathbf{J}, \boldsymbol{\lambda})$ , consisting of a compact open subgroup  $\mathbf{J}$  of  $G$  and an irreducible complex representation  $\boldsymbol{\lambda}$  of  $\mathbf{J}$ . This family of pairs  $(\mathbf{J}, \boldsymbol{\lambda})$ , called semisimple types, satisfies the following condition: for every inertial cuspidal class  $\Omega$ , there is a semisimple type  $(\mathbf{J}, \boldsymbol{\lambda})$  such that the irreducible complex representations of  $G$  with cuspidal support in  $\Omega$  are exactly those whose restriction to  $\mathbf{J}$  contains  $\boldsymbol{\lambda}$ .

In [16], Mínguez and the first author extended this construction to the modular case: they constructed a family of pairs  $(\mathbf{J}, \boldsymbol{\lambda})$ , consisting of a compact open subgroup  $\mathbf{J}$  of  $G$  and an irreducible modular representation  $\boldsymbol{\lambda}$  of  $\mathbf{J}$ , called semisimple supertypes. However, they did not give the relation between these semisimple supertypes and inertial supercuspidal classes of  $G$ . In this paper, we prove:

- for each inertial supercuspidal class  $\Omega$ , there is a semisimple supertype  $(\mathbf{J}, \boldsymbol{\lambda})$  such that the irreducible  $R$ -representations of  $G$  with supercuspidal support in  $\Omega$  are precisely those which appear as subquotients of the compactly induced representation  $\mathrm{ind}_{\mathbf{J}}^G(\boldsymbol{\lambda})$ ;