Astérisque

GUNTHER VOGEL ULRICH GÖRTZ TORSTEN WEDHORN VOLKER MEUSERS EVA VIEHMANN KONSTANTIN ZIEGLER STEFAN WEWERS INKEN VOLLAARD IRENE I. BOUW MICHAEL RAPOPORT **Argos seminar on intersections of modular correspondences**

Astérisque, tome 312 (2007)

<http://www.numdam.org/item?id=AST_2007__312__R1_0>

© Société mathématique de France, 2007, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (http://smf4.emath.fr/ Publications/Asterisque/) implique l'accord avec les conditions générales d'utilisation (http://www.numdam.org/conditions). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

\mathcal{N} umdam

Article numérisé dans le cadre du programme Numérisation de documents anciens mathématiques http://www.numdam.org/

ARGOS SEMINAR ON INTERSECTIONS OF MODULAR CORRESPONDENCES

Société Mathématique de France 2007 Publié avec le concours du Centre National de la Recherche Scientifique **2000** *Mathematics Subject Classification.* — 11G18, 11E08, 11F03, 11F30, 11F32, 11G15, 11G18, 14B12, 14G35, 14K07, 14K22, 14L05.

Key words and phrases. — Modular correspondences, endomorphisms of formal groups, quadratic forms over the ring of *p*-adic integers, Siegel-Eisenstein series.

ARGOS SEMINAR ON INTERSECTIONS OF MODULAR CORRESPONDENCES

Abstract. — This volume contains the written account of the Bonn seminar on arithmetic geometry 2003/2004. It gives a coherent exposition of the theory of intersections of modular correspondences. The focus of the seminar is the formula for the intersection number of arithmetic modular correspondences due to Gross and Keating. Other topics treated are Hurwitz's theorem on the intersection of modular correspondences over the field of complex numbers, and the relation of the arithmetic intersection numbers to Fourier coefficients of Siegel-Eisenstein series.

Also included is background material on one-dimensional formal groups and their endomorphisms, and on quadratic forms over the ring of *p*-adic integers.

Résumé (Séminaire ARGOS sur les intersections de correspondances modulaires)

Ce volume consiste des exposés faits dans le cadre du séminaire de géométrie arithmétique de Bonn en 2003/2004. Il donne une exposition systématique de la théorie des intersections de correspondances modulaires. Le but principal est la formule de Gross-Keating du nombre d'intersection de correspondances modulaires arithmétiques. Autres sujets traités sont le théorème de Hurwitz sur l'intersection de correspondances modulaires sur le corps des nombres complexes, et la relation des nombres d'intersection arithmétiques aux coefficients de Fourier des séries de Siegel-Eisenstein.

On a aussi inclus des rappels sur les groupes formels à un paramètre et leurs endomorphismes, et sur les formes quadratiques sur l'anneau des entiers p-adiques.

CONTENTS

1. Foreword	vii
Notation	xv
2. Modular Polynomials by Gunther Vogel	1
3. A sum of representation numbers by Ulrich Görtz	9
4. Arithmetic Intersection Numbers by Ulrich Görtz	15
5. The genus of the endomorphisms of a supersingular elliptic curve by Torsten Wedhorn	25
6. Lubin-Tate Formal Groups by Volker Meusers	49
7. Formal moduli of formal \mathcal{O}_K -modules by Eva Viehmann & Konstantin Ziegler	57
8. Canonical and quasi-canonical liftings by Stefan Wewers	67
9. Canonical and quasi-canonical liftings in the split case by Volker Meusers	87

CONTENTS

10. Lifting endomorphisms of formal \mathcal{O}_K -modules
11. Endomorphisms of quasi-canonical lifts
12. Invariants of ternary quadratic forms
13. Deformations of isogenies of formal groups
14. An alternative approach using ideal bases
15. Calculation of representation densities
16. The connection to Eisenstein series
Index

1. FOREWORD

1. Motivation and main results

This book is based on the notes for the $AP\GammaO\Sigma^{(1)}$ seminar of the winter semester 2003/2004 at the University of Bonn. Its aim was to go through the paper On the intersection of modular correspondences by Gross and Keating [**GK**], and understand it thoroughly. This subject was chosen for three reasons. First of all, it was felt that the mathematics contained in this paper (and the papers on which Gross and Keating base their article) is extremely interesting, and has become even more important recently, due to the use that S. Kudla and others have made of these results. Secondly, thanks to the elementary methods employed in the proofs of the main theorems, the seminar provided a rapid access, even to a novice in the field, to a deep and sophisticated topic in arithmetic algebraic geometry. Thirdly, it was felt from the start that the literature on the subject was not easy to penetrate and that therefore the effort made by all speakers to master this material should not be lost, and that a written account of the seminar should be made available.

The origin of the topics treated in the seminar goes back to the 19th century. Let $j = j(\tau)$ be the elliptic modular function on the upper half plane. For $m \ge 1$ let $\varphi_m(j,j') \in \mathbb{Z}[j,j']$ be the classical modular polynomial, defined by

(1.1)
$$\varphi_m(j(\tau), j(\tau')) = \prod_{\substack{A \in M_2(\mathbb{Z}) \\ \det(A) = m \\ \mod SL_2(\mathbb{Z})}} (j(\tau) - j(A\tau'))$$

Kronecker and Hurwitz established a number of important properties of these polynomials, as for instance their factorization into irreducible factors. They also proved degree formulas like

(1.2)
$$\deg f_m = \sum_{dd'=m} \max(d, d') ,$$

where $f_m(j) = \varphi_m(j, j)$, for m not a square.

⁽¹⁾Acronym for **Ar**ithmetische **G**eometrie **O**berseminar.

1. FOREWORD

From the point of view of the seminar, the interest in these results lies in the fact that they can be interpreted as giving intersection numbers on the complex surface $S_{\mathbb{C}} = \operatorname{Spec} \mathbb{C}[j, j']$. Let $T_{m,\mathbb{C}} \subseteq S_{\mathbb{C}}$ be the divisor defined by $\varphi_m = 0$. Then (1.2) can be interpreted as the intersection formula

(1.3)
$$(T_{m,\mathbb{C}} \cdot T_{1,\mathbb{C}}) = \sum_{dd'=m} \max(d,d') \quad ,$$

if m is not a square. Here $(T_{m_1,\mathbb{C}} \cdot T_{m_2,\mathbb{C}})$ is defined by

(1.4)
$$(T_{m_1,\mathbb{C}} \cdot T_{m_2,\mathbb{C}}) = \dim_{\mathbb{C}} \mathbb{C}[j,j']/(\varphi_{m_1},\varphi_{m_2}) \quad .$$

More generally, Hurwitz showed that the divisors $T_{m_1,\mathbb{C}}$ and $T_{m_2,\mathbb{C}}$ intersect properly on $S_{\mathbb{C}}$ if and only if m_1m_2 is not a perfect square and gave an explicit expression for the intersection number $(T_{m_1,\mathbb{C}} \cdot T_{m_2,\mathbb{C}})$. This in turn leads to the famous class number relations of Kronecker and Hurwitz.

Gross and Keating took up this classical subject by adding an arithmetic dimension to it. Instead of usual intersection numbers they consider arithmetic intersection numbers. Let $S = \text{Spec } \mathbb{Z}[j, j']$, which we consider as an arithmetic threefold. Let T_m be the arithmetic divisor defined by $\varphi_m = 0$. The arithmetic intersection number is defined for any triple of positive integers m_1, m_2, m_3 by

(1.5)
$$(T_{m_1} \cdot T_{m_2} \cdot T_{m_3}) = \log \# \mathbb{Z}[j, j'] / (\varphi_{m_1}, \varphi_{m_2}, \varphi_{m_3}) .$$

Gross and Keating derive a criterion for when this number is finite and give in this case an explicit expression for it (see below). This result is the main focus of the present book. Let us state it from the point of view adopted in these notes. Let \mathcal{M} be the moduli space of elliptic curves over Spec \mathbb{Z} (since we impose no level structure, \mathcal{M} is not a scheme, but a Deligne-Mumford stack). Put $\mathcal{S} = \mathcal{M} \times_{\text{Spec }\mathbb{Z}} \mathcal{M}$. For a positive integer m, let \mathcal{T}_m be the moduli space of isogenies of elliptic curves $E \to E'$ of degree m. Then \mathcal{T}_m maps by a finite unramified morphism to $\mathcal{M} \times \mathcal{M}$. From this point of view, the intersection number above should be interpreted as

$$\sum_{p} \log(p) \cdot \sum_{x \in \mathcal{X}(\overline{\mathbb{F}}_p)} \frac{1}{\# \operatorname{Aut}(x)} \lg \widehat{\mathcal{O}}_{\mathcal{X},x},$$

where we denote by \mathcal{X} the triple fiber product of \mathcal{T}_{m_1} , \mathcal{T}_{m_2} , and \mathcal{T}_{m_3} over $\mathcal{M} \times \mathcal{M}$. Here the weighting factor $\frac{1}{\#\operatorname{Aut}(x)}$ is due to the fact that \mathcal{X} is a stack.

We now state the main results contained in this volume.

We denote by $S_{\mathbb{C}}$ resp. by $T_{m,\mathbb{C}}$ the base change of S resp. T_m to Spec C. The first result is Hurwitz's theorem.

Theorem 1.1. — The cycles $T_{m_1,\mathbb{C}}$ and $T_{m_2,\mathbb{C}}$ intersect properly on $S_{\mathbb{C}}$ if and only if the integer $m = m_1m_2$ is not a perfect square. In this case, the intersection $T_{m_1,\mathbb{C}} \times_{S_{\mathbb{C}}} T_{m_2,\mathbb{C}}$ lies over the locus in $S_{\mathbb{C}}$ corresponding to pairs (E, E') of elliptic curves with complex multiplication by orders in the imaginary-quadratic field $\mathbb{Q}(\sqrt{-m})$ of discriminant $\geq -4m$. The intersection number is equal to

$$(T_{m_1,\mathbb{C}} \cdot T_{m_2,\mathbb{C}}) = \sum_{\substack{t \in \mathbb{Z} \\ t^2 < 4m}} \sum_{d \mid gcd(m_1,m_2,t)} d \cdot H\left(\frac{4m - t^2}{d^2}\right) \quad . \quad \Box$$

viii

ASTÉRISQUE 312

1. FOREWORD

Here H(n) denotes the Hurwitz class number (the number of $SL_2(\mathbb{Z})$ -equivalence classes of positive definite binary quadratic forms over \mathbb{Z} with determinant n).

The second result is the theorem of Gross and Keating.

Theorem 1.2. — The cycles T_{m_1}, T_{m_2} and T_{m_2} intersect properly on S if and only if there is no positive definite binary quadratic form over \mathbb{Z} which represents the three integers m_1, m_2, m_3 . In this case the intersection $T_{m_1} \times_S T_{m_2} \times_S T_{m_3}$ lies over the locus in S corresponding to pairs (E, E') of elliptic curves which are supersingular in some characteristic p with $p < 4m_1m_2m_3$. The arithmetic intersection number is equal to

$$(T_{m_1} \cdot T_{m_2} \cdot T_{m_3}) = \sum_{p < 4m_1m_2m_3} n(p)\log p$$

where

$$n(p) = \frac{1}{2} \cdot \sum_{Q} \left(\prod_{\substack{\ell \mid \Delta, \\ \ell \neq p}} \beta_{\ell}(Q) \right) \cdot \alpha_{p}(Q) \quad . \quad \Box$$

Here the sum is the taken over all positive definite integral ternary quadratic forms Q with diagonal (m_1, m_2, m_3) which are isotropic over \mathbb{Q}_{ℓ} for all $\ell \neq p$. Furthermore $\Delta = \frac{1}{2} \det Q$ and $\beta_{\ell}(Q)$ is a normalized representation density of Q by the \mathbb{Z}_{ℓ} -lattice $M_2(\mathbb{Z}_{\ell})$ with its norm form. Finally, $\alpha_p(Q)$ is the length of a certain local deformation space. Namely, one considers the universal deformation space of a triple of isogenies of formal groups of dimension 1 and height 2 over $\overline{\mathbb{F}}_p$. Here the passage from a global problem involving elliptic curves to a local problem on formal groups is provided by the Serre-Tate theorem. In [**GK**], Gross and Keating give completely explicit expressions for the factors $\beta_{\ell}(Q)$ and $\alpha_p(Q)$, comp. Chapters 5 and 13. They express these quantities in terms of new invariants of ternary quadratic forms over \mathbb{Z}_p which are defined by them for this purpose (the Gross-Keating invariant in $(\mathbb{Z}_{\geq 0})^3$ and the Gross-Keating epsilon factor in $\{\pm 1\}$). This is especially striking in the cases when $\ell = 2$ resp. p = 2, in the other cases these invariants can be expressed in terms of classical quantities.

The invariant $\alpha_p(Q)$ is probably the most interesting ingredient in the formula above, and we now give a precise definition.

Let G be a formal group of dimension 1 and height 2 over \mathbb{F}_p . Let $W = W(\mathbb{F}_p)$ be the ring of Witt vectors. The universal deformation of the pairs (G, G) is then (Γ, Γ') over the formal scheme $\hat{\mathcal{S}} = \operatorname{Spf} W[[t, t']]$. If now $f_1, f_2, f_3 : G \to G$ are three endomorphisms $\neq 0$, we let $I_i \subset W[[t, t']]$ for i = 1, 2, 3 be the minimum ideal such that f_i lifts to a homomorphism $\tilde{f}_i : \Gamma \to \Gamma' \pmod{I_i}$. Then I_i defines a divisor $\hat{\mathcal{T}}_i$ on $\hat{\mathcal{S}}$. Consider

(1.6)
$$(\hat{T}_1 \cdot \hat{T}_2 \cdot \hat{T}_3) = \text{length}_W W[[t, t']]/(I_1 + I_2 + I_3)$$

On End(G) we have the usual quadratic form Nm with values in \mathbb{Z}_p (the norm form, after identifying End(G) with the maximal order in the quaternion division algebra over \mathbb{Q}_p). It turns out that (1.6) only depends on the quadratic form $Q(f_1, f_2, f_3)$: