

*quatrième série - tome 50    fascicule 4    juillet-août 2017*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Laura DEMARCO & Kevin PILGRIM

*The classification of polynomial basins of infinity*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

**Responsable du comité de rédaction / *Editor-in-chief***

Emmanuel KOWALSKI

**Publication fondée en 1864 par Louis Pasteur**

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

**Comité de rédaction au 1<sup>er</sup> janvier 2017**

P. BERNARD      A. NEVES  
S. BOUCKSOM    J. SZEFTEL  
E. BREUILLARD   S. VŨ NGỌC  
R. CERF          A. WIENHARD  
G. CHENEVIER    G. WILLIAMSON  
E. KOWALSKI

**Rédaction / *Editor***

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

**Édition / *Publication***

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

**Abonnements / *Subscriptions***

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

**Tarifs**

Europe : 519 €. Hors Europe : 548 €. Vente au numéro : 77 €.

---

© 2017 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret  
Périodicité : 6 n<sup>os</sup> / an

# THE CLASSIFICATION OF POLYNOMIAL BASINS OF INFINITY

BY LAURA DEMARCO AND KEVIN PILGRIM

---

**ABSTRACT.** – We consider the problem of classifying the dynamics of complex polynomials  $f : \mathbb{C} \rightarrow \mathbb{C}$  restricted to the basins of infinity  $X(f)$ . We synthesize existing combinatorial tools—tableaux, trees, and laminations—into a new invariant of basin dynamics we call the pictograph. For polynomials with all critical points escaping to infinity, we obtain a complete description of the set of topological conjugacy classes with given pictograph. For arbitrary polynomials, we compute the total number of topological conjugacy classes of basins  $(f, X(f))$  with a given pictograph. We also define abstract pictographs and prove that every abstract pictograph is realized by a polynomial. Extra details are given in degree 3, and we provide examples that show the pictograph is a finer invariant than both the tableau of [5] and the tree of [10].

**RÉSUMÉ.** – Nous étudions la question de la classification de la dynamique des polynômes complexes  $f : \mathbb{C} \rightarrow \mathbb{C}$  restreints à leur bassin de l’infini. Nous faisons la synthèse d’outils de combinatoire — tableaux, arbres, laminations — en un nouvel invariant du bassin dynamique que nous appelons *pictogramme*. Pour les polynômes dont tous les points critiques s’échappent vers l’infini, nous obtenons une description complète de l’ensemble des classes de conjugaison topologiques ayant un pictogramme donné. Plus généralement, pour tout polynôme, nous calculons le nombre de classes de conjugaison topologiques du bassin  $(f, X(f))$  à pictogramme donné. Nous définissons les pictogrammes de façon abstraite et prouvons que chacun d’eux est réalisable par un polynôme. Nous donnons plus de détails en degré 3 et donnons des exemples montrant que le pictogramme est un invariant plus fin que les tableaux de [5] et que les arbres de [10].

## 1. Introduction

This article continues a study of the moduli space of complex polynomials  $f : \mathbb{C} \rightarrow \mathbb{C}$ , in each degree  $d \geq 2$ , in terms of the dynamics of polynomials on their basins of infinity [4, 5, 10, 8, 7]. Our main goal is to classify the topological conjugacy classes of a polynomial  $f$  restricted to its basin

$$X(f) = \{z \in \mathbb{C} : f^n(z) \rightarrow \infty\}.$$

The basin  $X(f)$  is an open, connected subset of  $\mathbb{C}$ . In degree  $d = 2$ , there are only two topological conjugacy classes of basins  $(f, X(f))$ , distinguished by the Julia set being connected or disconnected; see, for example, [19, Theorem 10.1]. In every degree  $d > 2$ , there are infinitely many topological conjugacy classes of basins, even among the structurally stable polynomials in the shift locus. The main objective of this article is the development of combinatorial methods that allow us to distinguish and enumerate these conjugacy classes in all degrees.

By definition, a polynomial  $f$  of degree  $d$  is in the shift locus if all of its  $d - 1$  critical points are in  $X(f)$ . In this case, the basin  $X(f)$  is a rigid Riemann surface, admitting up to Möbius transformations a unique embedding into the Riemann sphere ([18, §2.8], [1, §IV.4]). In this case, the restriction  $f : X(f) \rightarrow X(f)$  uniquely determines the conformal conjugacy class of  $f : \mathbb{C} \rightarrow \mathbb{C}$ . Thus, our results on basin dynamical systems  $(f, X(f))$ —given as Theorems 1.1, 1.2, 1.3, and 1.4 below—also provide a combinatorial classification of topological conjugacy classes of polynomials in the shift locus.

In the theory of dynamical systems, the study of a system like  $(f, X(f))$  is somewhat nonstandard. On the one hand, since all points tend to  $\infty$  under iteration, the system is transient. On the other hand, the structure of  $(f, X(f))$ , with an induced dynamical system on its Cantor set of ends as a topological space, carries enough information to recover the full entropy of the polynomial  $(f, \mathbb{C})$ ; see [10, Theorem 1.1]. Our methods and perspective are inspired by the two foundational articles of Branner and Hubbard on polynomial dynamics which lay the groundwork and treat the case of cubic polynomials in detail [4, 5].

### 1.1. The pictograph, informally described

We begin with a rough description of the pictograph  $\mathcal{D}(f)$  associated to a polynomial basin dynamics  $(f, X(f))$ . A formal presentation is given in §2.2 (for cubic polynomials) and Section 10 (in arbitrary degree).

*Global setup.* – The basin dynamics  $f : X(f) \rightarrow X(f)$  fits naturally into a sequence of dynamical systems related by semiconjugacies. These are organized in the diagram below, and are explained in the following paragraphs.

$$(1.1) \quad \begin{array}{ccccccc} \begin{array}{c} \mathcal{F} \\ \curvearrowright \\ \mathcal{X}(f) \end{array} & \xrightarrow{g_f} & \begin{array}{c} f \\ \curvearrowright \\ X(f) \end{array} & \xrightarrow{\pi_f} & \begin{array}{c} F \\ \curvearrowright \\ T(f) \end{array} & \xrightarrow{h_f} & \begin{array}{c} \cdot d \\ \curvearrowright \\ (0, \infty) \end{array} \\ & & & \searrow & \swarrow & & \\ & & & & G_f & & \end{array}$$

The map  $G_f$  is the harmonic *Green's function*; its values we call *heights* or sometimes *escape rates*. The grand orbits (under multiplication by  $d$ ) of heights of critical points are called *nongeneric heights*. We endow  $(0, \infty)$  with a simplicial structure in which the nongeneric heights are vertices. The map  $\pi_f$  collapses connected components of level sets of  $G_f$  to points; its image is the *DeMarco-McMullen tree*  $T(f)$ , and  $f$  induces a self-map  $F : T(f) \rightarrow T(f)$ . By construction, the factor  $h_f : T(f) \rightarrow (0, \infty)$  is simplicial.




FIGURE 1.1. The local features of the quotient maps in (1.1) at a vertex  $v$  of  $T(f)$  and the associated lamination  $L_v$ .

*Local features.* – See Figure 1.1. Every vertex  $v$  of  $T(f)$  has a nongeneric height  $h_f(v)$ ; and for each vertex, there is a maximal interval  $(a, b) \subset (0, \infty)$  containing  $h_f(v)$  for which all heights  $t \in (a, b) - \{h_f(v)\}$  are generic. The connected component  $X_v(f)$  of  $G_f^{-1}(a, b)$  containing  $\pi_f^{-1}(v)$  is a planar Riemann surface that we call a *local model surface*. The intersection  $L_{X_v} := X_v(f) \cap \pi_f^{-1}(v)$  is called the *central leaf* of  $X_v(f)$ . The central leaf  $L_{X_v}$  is a connected component of a fiber of  $G_f$ , is homeomorphic to the underlying space of a finite planar graph, and is the boundary of the unbounded component of its complement. This implies it is naturally the quotient of the unit circle by a certain kind of equivalence relation, a *finite lamination*, denoted  $L_v$ . The lamination  $L_v$  is encoded by a simple picture in a disk, a *lamination diagram*. The lamination diagram is not endowed with coordinates—rotating the picture does not change it—but the circle is equipped with a metric induced by the 1-form  $|\partial G_f|$ . For example: if two critical points  $c_1, c_2$  belong to the same central leaf  $L_{X_v}$ , their relative angles in  $L_v$  are determined by the metric.

*The pictograph.* – The pictograph  $\mathcal{D}(f)$  is a diagram consisting of a collection of laminations  $L_v$ , associated to vertices in the convex hull of the critical points of  $(F, T(f))$ , together with labels that mark the orbits of the critical points. For illustration, Figure 1.2 shows a pictograph associated to a polynomial of degree 4 in the shift locus. The markings on each lamination indicate which iterate of a critical point lands on a central leaf or is seen through the “pant leg” of the local model surface. We emphasize two things.

1. The pictograph contains both combinatorial and metric information. For example, if the iterates of two critical points, say  $f^i(c_1)$  and  $f^j(c_2)$ , both lie on a central leaf  $L_{X_v}$ , then the lamination  $L_v$  is labeled by symbols  $i_1$  and  $j_2$ , placed on the unit circle at a distance recording the metric information of how these points are deployed in  $L_{X_v}$ .
2. The pictograph is a static object. It does not, by definition, include a self-map of an object. However, it allows for reconstruction of dynamics, as described in the main results presented below.

The next paragraph describes what the pictograph captures.

*The tree of local models.* – The *tree of local models* is the disjoint union  $\mathcal{X}(f) := \bigsqcup_v X_v(f)$ , indexed by the vertices  $v$  of the tree  $T(f)$ . It is equipped with a holomorphic self-map  $\mathcal{F}$  induced by  $f$ . The collection of inclusions  $\{X_v(f) \hookrightarrow X(f)\}_{v \in T(f)}$  induces a (generically two-to-one) *gluing quotient map*  $g_f : \mathcal{X}(f) \rightarrow X(f)$ , which is not part of the data of the tree of local models.