

338

ASTÉRISQUE

2011

OPERADS AND CHAIN RULES
FOR THE CALCULUS OF FUNCTORS

Greg Arone & Michael Ching

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Astérisque est un périodique de la Société mathématique de France.

Numéro 338, 2011

Comité de rédaction

Ahmed ABBES	Bob OLIVER
Viviane BALADI	Fabrice PLANCHON
Gérard BESSON	Pierre SCHAPIRA
Laurent BERGER	Éric VASSEROT
Hélène ESNAULT	Wendelin WERNER
Damien GABORIAU	
Yves ANDRÉ (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 40 € (\$ 60)

Abonnement Europe : 463 €, hors Europe : 512 € (\$ 768)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2011

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0303-1179

ISBN 978-2-85629-308-9

Directeur de la publication : Aline BONAMI

338

ASTÉRISQUE

2011

OPERADS AND CHAIN RULES
FOR THE CALCULUS OF FUNCTORS

Greg Arone & Michael Ching

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Greg Arone
Department of Mathematics
University of Virginia
Charlottesville, VA 22904

Michael Ching
Department of Mathematics
University of Georgia
Athens, GA 30602

OPERADS AND CHAIN RULES FOR THE CALCULUS OF FUNCTORS

Greg ARONE and Michael CHING

Abstract. — We study the structure possessed by the Goodwillie derivatives of a pointed homotopy functor of based topological spaces. These derivatives naturally form a bimodule over the operad consisting of the derivatives of the identity functor. We then use these bimodule structures to give a chain rule for higher derivatives in the calculus of functors, extending that of Klein and Rognes. This chain rule expresses the derivatives of FG as a derived composition product of the derivatives of F and G over the derivatives of the identity.

There are two main ingredients in our proofs. Firstly, we construct new models for the Goodwillie derivatives of functors of spectra. These models allow for natural composition maps that yield operad and module structures. Then, we use a cosimplicial cobar construction to transfer this structure to functors of topological spaces. A form of Koszul duality for operads of spectra plays a key role in this.

Résumé (Opérades et règles de la chaîne pour le calcul fonctoriel). — Nous étudions la structure des dérivées de Goodwillie d'un foncteur d'homotopie pointé d'espaces topologiques possédant une base. Ces dérivées forment, de manière naturelle, un bimodule au-dessus de l'opérade, celui des dérivées du foncteur identité. Nous utilisons ces structures de bimodule pour donner une règle de la chaîne pour les dérivées supérieures en calcul fonctoriel, étendant celle de Klein et Rognes. La règle de la chaîne exprime les dérivées de FG en tant que produits de composition des dérivées de F et de G au-dessus des dérivées de l'identité.

Il y a deux ingrédients principaux dans nos preuves. Premièrement, nous construisons des nouveaux modèles pour les dérivées de Goodwillie des foncteurs de spectres. Ces modèles fournissent des applications de composition naturelles avec des structures de module et d'opérade. Ensuite, nous utilisons une construction de cobarre cosimplicielle pour porter cette structure aux foncteurs d'espaces topologiques. Une forme de dualité de Koszul pour les opérades de spectres joue un rôle-clé dans cette preuve.

CONTENTS

Introduction	1
1. Basics	17
1.1. Categories of spaces and spectra	17
1.2. Taylor tower and derivatives of functors of simplicial sets and spectra	26
1.3. Basic results on Taylor towers of composite functors	29
1.4. The category of functors	31
1.5. Subcomplexes of presented cell functors	37
1.6. Pro-spectra	40
2. Operads and modules	47
2.1. Composition products, operads and bar constructions	47
2.2. Homotopy invariance of the bar construction	55
2.3. Cofibrant replacements and model structures for operads and modules	59
2.4. Derived mapping objects for modules	68
2.5. Pro-symmetric sequences and Spanier-Whitehead Duality	72
3. Functors of spectra	83
3.1. Models for Goodwillie derivatives of functors of spectra	83
3.2. Composition maps for Goodwillie derivatives	92
3.3. Chain rule for functors from spectra to spectra	97
3.4. Operad structures for comonads	100
4. Functors of spaces	109
4.1. The cobar construction	109
4.2. Functors from spaces to spectra	112
4.3. Functors from spectra to spaces	128
4.4. Functors from spaces to spaces	134
4.5. A Koszul duality result for operads of spectra	137
4.6. Chain rules for functors of spaces and spectra	141
Appendix	143
A. Categories of operads, modules and bimodules	145
Bibliography	155

INTRODUCTION

In a landmark series of papers, [16], [17] and [18], Goodwillie outlines his ‘calculus of homotopy functors’. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ (where \mathcal{C} and \mathcal{D} are each either \mathbf{Top}_* , the category of pointed topological spaces, or \mathbf{Spec} , the category of spectra) be a pointed homotopy functor. One of the things that Goodwillie does is associate with F a sequence of spectra, which are called the *derivatives* of F . We denote these spectra by $\partial_1 F, \partial_2 F, \dots, \partial_n F, \dots$, or, collectively, by $\partial_* F$. Importantly, for each n the spectrum $\partial_n F$ has a natural action of the symmetric group Σ_n . Thus, $\partial_* F$ is a *symmetric sequence* of spectra.

The importance of the derivatives of F is that they contain substantial information about the homotopy type of F . Goodwillie’s main construction in [18] defines a sequence of ‘approximations’ to F together with natural transformations forming a so-called ‘Taylor tower’. This tower takes the form

$$F \rightarrow \dots \rightarrow P_n F \rightarrow P_{n-1} F \rightarrow \dots \rightarrow P_0 F$$

with $P_n F$ being the universal ‘ n -excisive’ approximation to F . (A functor is *n -excisive* if it takes any $n + 1$ -dimensional cube with homotopy pushout squares for faces to a homotopy cartesian cube.) For ‘analytic’ F , this tower *converges* for sufficiently highly connected X , that is

$$F(X) \simeq \operatorname{holim}_n P_n F(X).$$

In order to understand the functors $P_n F$ better, Goodwillie analyzes the fibre $D_n F$ of the map $P_n F \rightarrow P_{n-1} F$. This fibre is an ‘ n -homogeneous’ functor in an appropriate sense, and Goodwillie shows in [18] that $D_n F$ is determined by $\partial_n F$, via the following formula. If F takes values in \mathbf{Spec} then

$$D_n F(X) \simeq (\partial_n F \wedge X^{\wedge n})_{h\Sigma_n}.$$

If F takes values in \mathbf{Top}_* then one needs to prefix the right hand side with Ω^∞ .

This paper investigates the question of what additional structure the collection $\partial_* F$ naturally possesses, beyond the symmetric group actions. The first example of such structure was given by the second author in [9]. There, he constructed an operad structure on the sequence $\partial_* I_{\mathbf{Top}_*}$, where $I_{\mathbf{Top}_*}$ is the identity functor on \mathbf{Top}_* . Our first main result says that if F is a functor from \mathbf{Top}_* to \mathbf{Top}_* , then $\partial_* F$ has the structure of a bimodule over the operad $\partial_* I_{\mathbf{Top}_*}$. (For functors either only from or to \mathbf{Top}_* , we get left or right module structures respectively.)

It turns out that these bimodule structures are exactly what is needed to write a ‘chain rule’ for the calculus of functors. By a chain rule we mean a formula for describing the derivatives of a composite functor FG in terms of ∂_*F and ∂_*G . Such a chain rule was first studied by Klein and Rognes, in [27], who provided a complete answer to this question for first derivatives. In this paper we extend some of their work to higher derivatives, although with some restrictions. In particular, we only consider reduced functors (those with $F(*) \simeq *$) and only derivatives based at the trivial object $*$. (Klein and Rognes consider derivatives at a general base object.) Our result expresses $\partial_*(FG)$ as a derived ‘composition product’ of the $\partial_*I_{\mathbf{Top}_*}$ -bimodule structures on ∂_*F and ∂_*G .

The proofs of our main theorems are rather roundabout, but give us additional interesting results along the way. We first treat the case of functors from \mathbf{Spec} to \mathbf{Spec} , and construct new models for the derivatives of such functors. Then, to pass from \mathbf{Spec} to \mathbf{Top}_* , we rely heavily on the close connection between topological spaces and coalgebras over the cooperad $\Sigma^\infty\Omega^\infty$, and also on a form of ‘Koszul duality’ for operads in \mathbf{Spec} . Koszul duality for operads was first introduced by Ginzburg and Kapranov, in [14], in the context of operads of chain complexes. Some of their ideas were extended to operads of spectra by the second author in [9]. In particular, it was shown there that the operad $\partial_*I_{\mathbf{Top}_*}$ plays the role of the Koszul dual of the commutative cooperad, and hence is a spectrum-level version of the Lie operad. In this paper, we give a deeper topological reason behind this observation. The commutative cooperad appears because it is equivalent to the derivatives of the comonad $\Sigma^\infty\Omega^\infty$. One of our main results is that the derivatives of $I_{\mathbf{Top}_*}$ and $\Sigma^\infty\Omega^\infty$ are related by this form of Koszul duality.

The module and bimodule structures on the derivatives of a general functor F also arise via an extension of Koszul duality ideas to spectra. For example, we show that for $F : \mathbf{Top}_* \rightarrow \mathbf{Top}_*$, the derivatives of F and $\Sigma^\infty F$ are related by a corresponding duality between left $\partial_*I_{\mathbf{Top}_*}$ -modules and left $\partial_*(\Sigma^\infty\Omega^\infty)$ -comodules.

It seems to us that this paper gives a satisfactory answer to one of the open-ended questions proposed in the introduction to [9]: is there a deeper connection between calculus of functors and the theory of operads? Yes, there is a deeper connection. It stems from two basic sources. The first is the fact that composition of functors is related to the composition product of symmetric sequences. The second is the relationship between $I_{\mathbf{Top}_*}$ and $\Sigma^\infty\Omega^\infty$, which translates, on the level of derivatives, to Koszul duality of operads.

We now proceed with a more precise statement of our main results.

Our results. — As we mentioned already, our results are stated in the language of operads and modules over them. The collection of derivatives of a functor F (of either based spaces or spectra) forms a symmetric sequence of spectra, that is, a sequence

$$\partial_*F = (\partial_1F, \partial_2F, \dots)$$