351

ASTÉRISQUE

2013

DIFFRACTION OF SINGULARITIES FOR THE WAVE EQUATION ON MANIFOLDS WITH CORNERS

Richard MELROSE & András VASY & Jared WUNSCH

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Astérisque est un périodique de la Société mathématique de France.
Numéro 351, 2013

Comité de rédaction

Ahmed Abbes
Viviane Baladi
Gérard Besson
Laurent Berger
Philippe Biane
Hélène Esnault
Damien Gaboriau
Michael Harris
Fabrice Planchon
Pierre Schapira
Bertrand Toen

Yves André (dir.)

Diffusion

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 9
France
smf@smf.univ-mrs.fr

Hindustan Book Agency
O-131, The Shopping Mall
Arjun Marg, DLF Phase 1
Gurgaon 122002, Haryana
Inde

AMS
P.O. Box 6248
Providence RI 02940
USA
www.ams.org

Tarifs

Vente au numéro : 35 € ($ 52)
Abonnement Europe : 484 €, hors Europe : 523 € ($ 784)
Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris Cedex 05, France
Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96
revues@smf.ens.fr • http://smf.emath.fr/

© Société Mathématique de France 2013
Tous droits réservés (article L 122–4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335–2 et suivants du CPI.

ISSN 0303-1179
ISBN 978-2-85629-367-6

Directrice de la publication : Aline Bonami

351

ASTÉRISQUE

2013

DIFFRACTION OF SINGULARITIES FOR THE WAVE EQUATION ON MANIFOLDS WITH CORNERS

Richard MELROSE & András VASY & Jared WUNSCH

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Richard MELROSE
Department of Mathematics
MIT
Cambridge MA, USA
rbm@math.mit.edu

András VASY
Department of Mathematics
Stanford University
Stanford CA, USA
andras@math.stanford.edu

Jared WUNSCH
Department of Mathematics
Northwestern University
Evanston IL, USA
jwunsch@math.northwestern.edu

DIFFRACTION OF SINGULARITIES FOR THE WAVE EQUATION ON MANIFOLDS WITH CORNERS

Richard MELROSE, András VASY and Jared WUNSCH

Abstract. — We consider the fundamental solution to the wave equation on a manifold with corners of arbitrary codimension. If the initial pole of the solution is appropriately situated, we show that the singularities which are diffracted by the corners (i.e., loosely speaking, are not propagated along limits of transversely reflected rays) are smoother than the main singularities of the solution. More generally, we show that subject to a hypothesis of nonfocusing, diffracted wavefronts of any solution to the wave equation are smoother than the incident singularities. These results extend our previous work on edge manifolds to a situation where the fibers of the boundary fibration, obtained here by blowup of the corner in question, are themselves manifolds with corners.

***Résumé* (Diffraction des singularités de l'équation d'onde sur les variétés à coins).** — Nous considérons la solution fondamentale à l'équation d'onde sur une variété à coins de codimension arbitraire. Si le pôle initial de la solution est situé arbitrairement, nous montrons que les singularités diffractées par les coins (autrement dit, intuitivement, ne sont pas propagées le long des limites de rayons réfléchis de manière transverse) sont plus lisses que les singularités principales de la solution. Plus généralement, nous montrons que sous une condition de non-focalisation, les fronts d'onde diffractés de toute solution de l'équation d'onde sont plus lisses que les singularités incidentes. Ces résultats étendent nos travaux précédents sur les variétés à bord, à une situation où les fibres de la fibration de bord, obtenue ici par un blow-up du coin en question, sont elles-mêmes des variétés à coins.

© Astérisque 351, SMF 2013

CONTENTS

1. Introduction 1
1.1. The problem and its history 1
1.2. The hypothesis 5
1.3. Structure of the proof 6
1.4. Sketch of the propagation results 7
1.5. Organization of the paper 10
1.6. Acknowledgements 11

2. Geometry: metric and Laplacian 13

3. Bundles and bicharacteristics 17
3.1. b-Cotangent bundle 17
3.2. Edge-smooth cotangent bundle 22
3.3. Edge-b cotangent bundle 24
3.4. Bicharacteristics 27
3.5. A summary 38

4. Edge-b calculus 41

5. Differential-pseudodifferential operators 45
5.1. The calculus 45
5.2. Dual spaces and adjoints 51
5.3. Domains 56
5.4. The wave operator as an element of $x^{-2}\operatorname{Diff}^2_{\mathrm{es},\sharp}(M)$ 58

6. Coisotropic regularity and non-focusing 67

7. Edge propagation 75
7.1. Radial points in the interior of the edge 75
7.2. Propagation into radial points over the boundary of the edge 76
7.3. Propagation at hyperbolic points within the edge 89
7.4. Propagation at glancing points within the edge 96

8. Propagation of fiber-global coisotropic regularity 103

9. Geometric theorem 107

A. Some functional analysis 117

B. The edge-b calculus 123

Index of Notation 131

Bibliography 133

CHAPTER 1

INTRODUCTION

1.1. The problem and its history

Let X_0 be a manifold with corners, of dimension n, i.e., a manifold locally modeled on $(\mathbb{R}^+)^{f+1} \times \mathbb{R}^{n-f-1}$, endowed with an incomplete metric, smooth and nondegenerate up to the boundary. We consider the wave equation

$$\Box u \equiv D_t^2 u - \Delta u = 0 \text{ on } M_0 = \mathbb{R} \times X_0, \tag{1.1.1}$$

where $D_t = \imath^{-1}(\partial/\partial t)$ and Δ is the nonnegative Laplace-Beltrami operator; we will impose either Dirichlet or Neumann conditions at ∂X_0. As is well known by the classic result of Duistermaat-Hörmander[(1)] (see **[4]**), the wavefront set of a solution u propagates along null-bicharacteristics in the interior. However, the behavior of singularities striking the boundary and corners of M_0 is considerably subtler.

Indeed the propagation of singularities for the wave equation on manifolds with boundary is already a rather subtle problem owing the the difficulties posed by "glancing" bicharacteristics, those which are tangent to the boundary. Chazarain **[1]** showed that singularities striking the boundary transversely simply reflect according to the usual law of geometric optics (conservation of energy and tangential momentum, hence "angle of incidence equals angle of reflection") for the reflection of bicharacteristics. This result was extended in **[21]** and **[22]** by showing that, at glancing points, singularities may only propagate along certain generalized bicharacteristics. The continuation of these curves may fail to be unique at (non-analytic) points of infinite-order tangency as shown by Taylor **[27]**. Whether all of these branches of bicharacteristics can carry singularities is still not known.

As was shown initially in several special examples (namely those amenable to separation of variables) the interaction of wavefront set with a corner gives rise to new, *diffractive* phenomena, in which a single bicharacteristic carrying a singularity into a corner produces singularities departing from the corner along a whole family of bicharacteristics. For instance, a ray carrying a singularity transversely into a codimension-two corner will in general produce singularities on the entire cone of rays reflected in

(1) This result, viewed in the context of hyperbolic equations, built on a considerable body of work prior to the introduction of the wavefront set; see especially **[9, 12]**.

such a way as to conserve both energy and momentum tangent to the corner (see Figure 1) The first diffraction problem to be rigorously treated was that of the exterior



FIGURE 1. A ray carrying a singularity may strike a corner of codimension two and give rise to a whole family of diffracted singularities, conserving both energy and momentum along the corner.

of a wedge,[(2)] which was analyzed by Sommerfeld **[26]**; subsequently, many related examples were analyzed by Friedlander **[5]**, and more generally the case of exact cones was worked out explicitly by Cheeger-Taylor **[2]**, **[3]** in terms of the functional calculus for the Laplace operator on the cross section of the cone. All of these explicit examples reveal that generically a diffracted wave arises from the interaction of wavefront set of the solution with singular strata of the boundary of the manifold; this has long been understood at a heuristic level, with the geometric theory of diffraction of Keller **[8]** describing the classes of trajectories that ought to contribute to the asymptotics of the solution in various regimes.

Subsequent work has been focused primarily on characterizing the bicharacteristics on which singularities can propagate, and on describing the strength and form of the singularities that arise. The propagation of singularities on manifolds with boundary was first understood in the analytic case by Sjöstrand **[23, 24, 25]**, and subsequently generalized to a very wide class of manifolds, including manifolds with corners, by Lebeau **[10, 11]**. In the $\mathcal{C}^\infty$ setting employed here, the special case of manifolds with conic singularities was studied by Melrose-Wunsch **[16]** and *edge manifolds* (i.e., cone bundles) were considered by Melrose-Vasy-Wunsch **[15]**. Vasy **[30]** obtained results analogous to Lebeau's in the case of manifolds with corners, and it is the results of this work that directly bear on the situation studied here.

While the foregoing results characterize which bicharacteristics may carry singularities for solutions to the wave equation, they ignore the question of the regularity

(2) This is not in fact a manifold with corners, but is quite closely related.