

389

ASTÉRISQUE

2017

THE CUBIC SZEGŐ EQUATION AND
HANKEL OPERATORS

Patrick Gérard
Sandrine Grellier

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Astérisque est un périodique de la Société mathématique de France

Numéro 389

Comité de rédaction

Ahmed ABBES
Viviane BALADI
Laurent BERGER
Philippe BIANE
Hélène ESNAULT

Philippe EYSSIDIEUX
Damien GABORIAU
Michael HARRIS
Fabrice PLANCHON
Pierre SCHAPIRA

Éric VASSEROT (dir.)

Diffusion

Maison de la SMF
B.P. 67
13274 Marseille CEDEX 9
France
christian.smf@cirm-math.fr

AMS
P.O. Box 6248
Providence RI 02940
USA
www.ams.org

Tarifs 2017

Vente au numéro : 35 € (\$ 52)

Abonnement électronique : 500 € (\$ 750)

Abonnement avec supplément papier : 657 €, hors Europe : 699 € (\$ 1049)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris CEDEX 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

astsmf@ihp.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2017

Tous droits réservés (article L122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L335-2 et suivants du CPI.

ISSN : print 0303-1179, electronic 2492-5926

ISBN 978-2-85629-854-1

Directeur de la publication : Stéphane SEURET

ASTÉRISQUE 389

THE CUBIC SZEGŐ EQUATION AND
HANKEL OPERATORS

Patrick Gérard
Sandrine Grellier

P. Gérard

Laboratoire de Mathématiques d'Orsay, Université Paris-Sud, CNRS,
Université Paris–Saclay, 91405 Orsay, France.

E-mail : `Patrick.Gerard@math.u-psud.fr`

S. Grellier

Fédération Denis Poisson, MAPMO-UMR 6628, Département de Mathématiques,
Université d'Orléans, 45067 Orléans Cedex 2, France.

E-mail : `Sandrine.Grellier@univ-orleans.fr`

2010 Mathematics Subject Classification. — 35B15, 47B35, 37K15.

Key words and phrases. — Cubic Szegő equation, integrable system, nonlinear Schrödinger equation, Hankel operator, spectral analysis.

The authors are grateful to Z. Hani, T. Kappeler, S. Kuksin, V. Peller, A. Pushnitski and N. Tzvetkov for valuable discussions and comments about earlier versions of this work, to L. Baratchart for pointing them references [1] and [3], and to A. Nazarov for references concerning Bateman–type formulae. They also thank the anonymous referees for making helpful suggestions and comments. The first author is supported by ANR ANAE 13-BS01-0010-03.

THE CUBIC SZEGŐ EQUATION AND HANKEL OPERATORS

Patrick Gérard, Sandrine Grellier

Abstract. — This monograph is devoted to the dynamics on Sobolev spaces of the cubic Szegő equation on the circle \mathbb{S}^1 ,

$$i\partial_t u = \Pi(|u|^2 u).$$

Here Π denotes the orthogonal projector from $L^2(\mathbb{S}^1)$ onto the subspace $L^2_+(\mathbb{S}^1)$ of functions with nonnegative Fourier modes. We construct a nonlinear Fourier transformation on $H^{1/2}(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$ allowing to describe explicitly the solutions of this equation with data in $H^{1/2}(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$. This explicit description implies almost-periodicity of every solution in this space. Furthermore, it allows to display the following turbulence phenomenon. For a dense G_δ subset of initial data in $C^\infty(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$, the solutions tend to infinity in H^s for every $s > \frac{1}{2}$ with super-polynomial growth on some sequence of times, while they go back to their initial data on another sequence of times tending to infinity. This transformation is defined by solving a general inverse spectral problem involving singular values of a Hilbert–Schmidt Hankel operator and of its shifted Hankel operator.

Résumé (Équation de Szegő cubique et opérateurs de Hankel)

Cette monographie est consacrée à l'étude de la dynamique, dans les espaces de Sobolev, de l'équation de Szegő cubique sur le cercle \mathbb{S}^1 ,

$$i\partial_t u = \Pi(|u|^2 u),$$

où Π désigne le projecteur orthogonal de $L^2(\mathbb{S}^1)$ sur le sous-espace $L^2_+(\mathbb{S}^1)$ des fonctions à modes de Fourier positifs ou nuls. On construit une transformée de Fourier non linéaire sur $H^{1/2}(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$ permettant de résoudre explicitement cette équation avec données initiales dans $H^{1/2}(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$. Ces formules explicites entraînent la presque périodicité des solutions dans cet espace. Par ailleurs, elles permettent de mettre en évidence le phénomène de turbulence suivant. Pour un G_δ dense de données initiales de $C^\infty(\mathbb{S}^1) \cap L^2_+(\mathbb{S}^1)$, les solutions tendent vers l'infini à vitesse sur-polynomiale en norme $H^s(\mathbb{S}^1)$ pour tout $s > \frac{1}{2}$ sur une suite de temps, alors qu'elles retournent vers leur donnée initiale sur une autre suite de temps tendant vers l'infini. Cette transformation est définie *via* la résolution d'un problème spectral inverse lié aux valeurs singulières d'un opérateur de Hankel Hilbert-Schmidt et de son opérateur décalé.

CONTENTS

1. Introduction	1
2. Hankel operators and the Lax pair structure	9
2.1. Hankel operators H_u and K_u	9
2.2. The Lax pair structure	11
2.3. Application: an exponential bound for Sobolev norms	13
3. Spectral analysis	15
3.1. Spectral decomposition of the operators H_u and K_u	15
3.2. Some Bateman-type formulae	19
3.3. Finite Blaschke products	24
3.4. Two results by Adamyan–Arov–Krein	25
3.5. Multiplicity and Blaschke products	29
4. The inverse spectral theorem	35
4.1. The inverse spectral theorem in the finite rank case	36
4.2. Extension to the Hilbert-Schmidt class	53
4.3. Extension to compact Hankel operators	57
5. The Szegő dynamics	59
5.1. Evolution under the cubic Szegő flow	59
5.2. Application: traveling waves revisited	62
5.3. Application to almost periodicity	64
6. Long Time instability and unbounded Sobolev orbits	65
6.1. Instability and genericity of unbounded orbits	65
6.2. A family of quasiperiodic solutions	66
6.3. Construction of the smooth family of data	68
6.4. The singular behavior	75
7. Geometry of the Fourier transform	89

7.1. Evolution under the Szegő hierarchy	89
7.2. The symplectic form on $\mathcal{V}_{(d_1, \dots, d_n)}$	94
7.3. Invariant tori of the Szegő hierarchy and unitary equivalence of pairs of Hankel operators	100
Bibliography	109
Index	113

CHAPTER 1

INTRODUCTION

The large time behavior of solutions to Hamiltonian partial differential equations is an important problem in mathematical physics. In the case of finite dimensional Hamiltonian systems, many features of the large time behavior of trajectories are described using the topology of the phase space. For a given infinite dimensional system, several natural phase spaces, with different topologies, can be chosen, and the large time properties may strongly depend on the choice of such topologies. For instance, it is known that the cubic defocusing nonlinear Schrödinger equation

$$i\partial_t u + \Delta u = |u|^2 u$$

posed on a Riemannian manifold M of dimension $d = 1, 2, 3$ with sufficiently uniform properties at infinity, defines a global flow on the Sobolev spaces $H^s(M)$ for every $s \geq 1$. In this case, a typical large time behavior of interest is the boundedness of trajectories. On the energy space $H^1(M)$, the conservation of energy trivially implies that all the trajectories are bounded. On the other hand, the existence of unbounded trajectories in $H^s(M)$ for $s > 1$ as well as bounds of the growth in time of the $H^s(M)$ norms is a long standing problem [7]. As a way to detect and to measure the transfer of energy to small scales, this problem is naturally connected to wave turbulence. In [6] and [38], it has been established that big H^s norms grow at most polynomially in time. Existence of unbounded trajectories only recently [22] received a positive answer in some very special cases, while several instability results were already obtained [9], [20], [21], [24], [19]. Natural model problems for studying these phenomena seem to be those for which the calculation of solutions is the most explicit, namely integrable systems. Continuing with the example of nonlinear Schrödinger equations, a typical example is the one dimensional cubic nonlinear Schrödinger defocusing equation ([43]). However, in this case, the set of conservation laws is known to control the whole regularity of the solution, so that all the trajectories of $H^s(M)$ are bounded in $H^s(M)$ for every nonnegative integer s . In fact, when the equation is posed on the

circle, the recent results of [18] show that, for every such s , the trajectories in $H^s(M)$ are almost periodic in $H^s(M)$.

The goal of this monograph is to study an integrable infinite dimensional system, connected to a nonlinear wave equation, with a dramatically different large time behavior of its trajectories according to the regularity of the phase space.

Following [30] and [42], it is natural to change the dispersion relation by considering the family of equations, $\alpha < 2$,

$$i\partial_t u - |D|^\alpha u = |u|^2 u$$

posed on the circle \mathbb{S}^1 , where the operator $|D|^\alpha$ is defined by

$$\widehat{|D|^\alpha u}(k) = |k|^\alpha \hat{u}(k)$$

for every distribution $u \in \mathcal{D}'(\mathbb{S}^1)$. Numerical simulations in [30] and [42] suggest weak turbulence. For $1 < \alpha < 2$, it is possible to recover at most polynomial growth of the Sobolev norms ([39]). In the case $\alpha = 1$, the so-called half-wave equation

$$(1.0.1) \quad i\partial_t u - |D|u = |u|^2 u$$

defines a global flow on $H^s(\mathbb{S}^1)$ for every $s \geq \frac{1}{2}$ ([11] — see also [36]). In that case, the only available bound of the H^s -norms is e^{ct^2} ([39]), due to the lack of dispersion. Therefore, the special case $\alpha = 1$ seems to be a more favorable framework for displaying wave turbulence effects.

Notice that this equation can be reformulated as a system, using the Szegő projector Π defined on $\mathcal{D}'(\mathbb{S}^1)$ as

$$\widehat{\Pi u}(k) = \mathbf{1}_{k \geq 0} \hat{u}(k).$$

Indeed, setting $u_+ := \Pi u$, $u_- := (I - \Pi)u$, equation (1.0.1) is equivalent to the system

$$\begin{cases} i(\partial_t + \partial_x)u_+ &= \Pi(|u|^2 u) \\ i(\partial_t - \partial_x)u_- &= (I - \Pi)(|u|^2 u) \end{cases}$$

Furthermore, if the initial datum u_0 satisfies $u_0 = \Pi u_0$ and belongs to $H^s(\mathbb{S}^1)$, $s > 1$, with a small norm ε , then the corresponding solution u is approximated by the solution v of

$$(1.0.2) \quad i(\partial_t + \partial_x)v = \Pi(|v|^2 v),$$

for $|t| \leq \frac{1}{\varepsilon^2} \log \frac{1}{\varepsilon}$ [11] [36]. In other words, the first nonlinear effects in the above system arise through a decoupling of the two equations. Notice that an elementary change of variable in equation (1.0.2) reduces it to

$$(1.0.3) \quad i\partial_t u = \Pi(|u|^2 u).$$

Equation (1.0.3) therefore appears as a model evolution for describing the dynamics of the half-wave equation (1.0.1). We introduced it in [10] under the name *cubic*