

397

ASTÉRISQUE

2018

TILTING MODULES AND THE p -CANONICAL BASIS

Simon Riche & Geordie Williamson

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Astérisque est un périodique de la Société Mathématique de France.

Numéro 397, 2018

Comité de rédaction

Ahmed ABBES Hélène ESNAULT
Viviane BALADI Philippe EYSSIDIEUX
Laurent BERGER Michael HARRIS
Philippe BIANE Alexandru OANCEA
Nicolas BURQ Fabrice PLANCHON
Damien CALAQUE
Éric VASSEROT (dir.)

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
Christian.munusami@smf.emath.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 45 € (\$ 67)

Abonnement Europe : 665 €, hors Europe : 718 € (\$ 1 077)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Astérisque

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

nathalie.christiaen@smf.emath.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2018

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN : 0303-1179 (print) 2492-5926 (electronic)

ISBN 978-2-85629-880-0

Directeur de la publication : Stéphane Seuret

397

ASTÉRISQUE

2018

TILTING MODULES AND THE p -CANONICAL BASIS

Simon Riche & Geordie Williamson

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE

Simon Riche

Université Clermont Auvergne, CNRS, LMBP, F-63000 Clermont-Ferrand, France.
`simon.riche@uca.fr`

Geordie Williamson

School of Mathematics and Statistics F07, University of Sydney NSW 2006, Australia.
`g.williamson@sydney.edu.au`

Classification mathématique par sujet (2010). — 17B10, 20G15, 14F05.

Mots-clefs. — Modules basculants, groupes algébriques, faisceaux à parité, catégorie de Hecke.

S.R. was partially supported by ANR Grant No. ANR-13-BS01-0001-01. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 677147).

TILTING MODULES AND THE p -CANONICAL BASIS

by Simon RICHE & Geordie WILLIAMSON

Abstract. — In this book we propose a new approach to tilting modules for reductive algebraic groups in positive characteristic. We conjecture that translation functors give an action of the (diagrammatic) Hecke category of the affine Weyl group on the principal block. Our conjecture implies character formulas for the simple and tilting modules in terms of the p -canonical basis, as well as a description of the principal block as the antispherical quotient of the Hecke category. We prove our conjecture for $GL_n(\mathbb{k})$ using the theory of 2-Kac-Moody actions. Finally, we prove that the diagrammatic Hecke category of a general crystallographic Coxeter group may be described in terms of parity complexes on the flag variety of the corresponding Kac-Moody group.

Résumé — Dans cet ouvrage nous proposons une nouvelle approche à l'étude des modules basculants pour les groupes algébriques réductifs sur des corps de caractéristique positive. Nous conjecturons que les foncteurs de translation induisent une action de la catégorie de Hecke (diagrammatique) du groupe de Weyl affine sur le bloc principal. Cette conjecture implique des formules de caractères pour les modules simples et les modules basculants en termes de la base p -canonique, ainsi qu'une description du bloc principal comme le quotient anti-sphérique de la catégorie de Hecke. Nous démontrons notre conjecture pour le groupe $GL_n(\mathbb{k})$ en utilisant la théorie des représentations des algèbres 2-Kac-Moody. Enfin, nous prouvons que la catégorie de Hecke diagrammatique d'un groupe de Coxeter cristallographique général peut être décrite en termes de faisceaux à parité sur la variété de drapeaux du groupe de Kac-Moody correspondant.

CONTENTS

1. Introduction	1
1.1. Overview	1
1.2. The “categorical” conjecture	2
1.3. Tilting modules and the antispherical module	4
1.4. Tilting characters	6
1.5. The case of the group $GL_n(\mathbb{k})$	8
1.6. The diagrammatic Hecke category and parity sheaves	9
1.7. Variants	10
1.8. Simple characters	11
1.9. Comparison with Lusztig’s conjecture	15
1.10. Acknowledgements	16
1.11. Organization of the book	17
Part I. General conjecture	19
2. Tilting objects and sections of the ∇-flag	21
2.1. Highest weight categories	21
2.2. Canonical ∇ -flags	22
2.3. Tilting objects and sections of the ∇ -flag	23
3. Regular and subregular blocks of reductive groups	27
3.1. Definitions	27
3.2. Translation functors	29
3.3. Sections of the ∇ -flag and translation to a wall	30
3.4. Sections of the ∇ -flag and translation from a wall	33
3.5. Morphisms between “Bott-Samelson type” tilting modules	36
4. Diagrammatic Hecke category and the antispherical module	39
4.1. The affine Hecke algebra and the antispherical module	39
4.2. Diagrammatic Soergel bimodules	41
4.3. Two lemmas on \mathcal{D}_{BS}	45
4.4. Categorified antispherical module	46
4.5. Morphisms in the categorified antispherical module	47
5. Main conjecture and consequences	53

5.1. Statement of the conjecture	53
5.2. Tilting modules and antispherical Soergel bimodules	55
5.3. Surjectivity	56
5.4. Dimensions of morphism spaces	59
5.5. Proof of Theorem 5.2.1	60
5.6. Graded form of $\text{Rep}_0(G)$	61
5.7. Integral form of $\text{Tilt}(\text{Rep}_0(G))$	64
5.8. Integral form of $\text{Rep}_0(G)$	66
Part II. The case of $\text{GL}_n(\mathbb{k})$	73
6. Representations of GL_n in characteristic p as a 2-representation of $\widehat{\mathfrak{gl}}_p$	77
6.1. The affine Lie algebra $\widehat{\mathfrak{gl}}_N$	77
6.2. The natural representation of $\widehat{\mathfrak{gl}}_N$	78
6.3. Realization of $\bigwedge^n \text{nat}_p$ as a Grothendieck group	79
6.4. $\text{Rep}(G)$ as a 2-representation	84
7. Restriction of the representation to $\widehat{\mathfrak{gl}}_n$	93
7.1. Combinatorics	93
7.2. Categorifying the combinatorics	94
7.3. First relations	95
7.4. Restriction of the 2-representation to $\mathcal{U}(\widehat{\mathfrak{gl}}_n)$	100
8. From categorical $\widehat{\mathfrak{gl}}_n$-actions to \mathcal{D}_{BS}-modules	103
8.1. Strategy	103
8.2. Preliminary lemmas	105
8.3. Polynomials	107
8.4. One color relations	108
8.5. Cyclicity	109
8.6. Jones-Wenzl relations	110
8.7. Two color associativity	112
8.8. Zamolodchikov (three color) relations	117
Part III. Relation to parity sheaves	125
9. Parity complexes on flag varieties	127
9.1. Reminder on Kac-Moody groups and their flag varieties	127
9.2. Partial flag varieties	130
9.3. Derived categories of sheaves on \mathcal{X} and \mathcal{X}^s	132
9.4. Parity complexes on flag varieties	133
9.5. Sections of the $!$ -flag	136
9.6. Sections of the $!$ -flag and pushforward to \mathcal{X}^s	137
9.7. Sections of the $!$ -flag and pullback from \mathcal{X}^s	140
9.8. Morphisms between “Bott-Samelson type” parity complexes	142

10. Parity complexes and the Hecke category	145
10.1. Diagrammatic category associated with \mathcal{G}	145
10.2. More on Bott-Samelson parity complexes	146
10.3. Statement of the equivalences	147
10.4. Construction of the functor Δ_{BS}	148
10.5. Verification of the relations	152
10.6. Fully-faithfulness of Δ_{BS}	154
10.7. The case of the affine flag variety	158
11. Whittaker sheaves and antispherical diagrammatic categories	161
11.1. Definition of Whittaker sheaves	161
11.2. Whittaker parity complexes	163
11.3. Sections of the $!$ -flag for Whittaker parity complexes	165
11.4. Surjectivity	167
11.5. Description of the antispherical diagrammatic category in terms of Whittaker sheaves	168
11.6. Application to the light leaves basis in the antispherical category ..	169
11.7. Iwahori-Whittaker sheaves on the affine flag variety	170
List of notation	173
Bibliography	179

CHAPTER 1

INTRODUCTION

1.1. Overview

In this book we give new conjectural character formulas for simple and indecomposable tilting modules for a connected reductive algebraic group in characteristic p ,⁽¹⁾ and we prove our conjectures in the case of the group $\mathrm{GL}_n(\mathbb{k})$ when $n \geq 3$. These conjectures are formulated in terms of the p -canonical basis of the corresponding affine Hecke algebra. They should be regarded as evidence for the philosophy that Kazhdan-Lusztig polynomials should be replaced by p -Kazhdan-Lusztig polynomials in modular representation theory. From this point of view several conjectures (Lusztig's conjecture, James' conjecture, Andersen's conjecture) become the question of agreement between canonical (or Kazhdan-Lusztig) and p -canonical bases.

In the general setting we prove that the new character formulas follow from a very natural conjecture of a more categorical nature, which has remarkable structural consequences for the representation theory of reductive algebraic groups. It is a classical observation that wall-crossing functors provide an action of the affine Weyl group W on the Grothendieck group of the principal block; in this way, the principal block gives a categorification of the antispherical module for W . We conjecture that this action can be categorified: namely, that the action of wall-crossing functors on the principal block gives rise to an action of the diagrammatic Bott-Samelson Hecke category attached to W as in [31]. From this conjecture we deduce the following properties.

1. The principal block is equivalent (as a module category over the diagrammatic Bott-Samelson Hecke category) to a categorification of the antispherical module defined by diagrammatic generators and relations.
2. The principal block admits a grading. Moreover, this graded category arises via extension of scalars from a category defined over the integers. Thus the principal block of any reductive algebraic group admits a “graded integral form”.

1. This book is written under the assumption that p is (strictly) larger than the Coxeter number, and so p cannot be too small. However there is a variant of our conjectures for any p involving singular variants of the Hecke category. In particular, it seems likely that p -Kazhdan-Lusztig polynomials give correct character formulas for tilting modules in *any* characteristic (see Conjecture 1.4.3). We hope to return to this subject in a future work.

3. The (graded) characters of the simple and tilting modules are determined by the p -canonical basis in the antispherical module for the Hecke algebra of W .

From (1) one may describe the principal block in terms of parity sheaves on the affine flag variety, which raises the possibility of calculating simple and tilting characters topologically. Point (2) gives a strong form of “independence of p ”. Finally, point (3) implies Lusztig’s character formula for large p .

We prove this “categorical” conjecture (hence in particular the character formulas) for the groups $\mathrm{GL}_n(\mathbb{k})$ using the Khovanov-Lauda-Rouquier theory of 2-Kac-Moody algebra actions. We view this, together with the agreement with Lusztig’s conjecture for large p and character formulas of Soergel and Lusztig in the context of quantum groups (see §1.7 for details) as strong evidence for our conjecture.

1.2. The “categorical” conjecture

Let G be a connected reductive algebraic group over an algebraically closed field \mathbb{k} of characteristic p with simply connected derived subgroup. We assume that $p > h$, where h is the Coxeter number of G . Let $T \subset B \subset G$ be a maximal torus and a Borel subgroup in G . Denote by $\mathbf{X} := X^*(T)$ the lattice of characters of T and by $\mathbf{X}^+ \subset \mathbf{X}$ the subset of dominant weights. Consider a regular block $\mathrm{Rep}_0(G)$ of the category of finite-dimensional algebraic G -modules, corresponding to a weight $\lambda_0 \in \mathbf{X}$ in the fundamental alcove, with its natural highest weight structure. If Φ is the root system of (G, T) , $W_f = N_G(T)/T$ is the corresponding Weyl group, and $W := W_f \ltimes \mathbb{Z}\Phi$ is the affine Weyl group, then the simple, standard, costandard and indecomposable tilting objects in the highest weight category $\mathrm{Rep}_0(G)$ are all parametrized by $W \bullet \lambda_0 \cap \mathbf{X}^+$. (Here “ \bullet ” denotes “ p -dilated dot action of W of \mathbf{X} ,” see §3.1.) If ${}^fW \subset W$ is the subset of elements w which are minimal in their coset $W_f w$, then there is a natural bijection

$${}^fW \xrightarrow{\sim} W \bullet \lambda_0 \cap \mathbf{X}^+$$

sending w to $w \bullet \lambda_0$. In this way we can parametrize the simple, standard, costandard and indecomposable tilting objects in $\mathrm{Rep}_0(G)$ by fW , and denote them by $\mathbb{L}(x \bullet \lambda_0)$, $\Delta(x \bullet \lambda_0)$, $\nabla(x \bullet \lambda_0)$ and $\mathbb{T}(x \bullet \lambda_0)$ respectively. In particular, on the level of Grothendieck groups we have

$$(1.2.1) \quad [\mathrm{Rep}_0(G)] = \bigoplus_{x \in {}^fW} \mathbb{Z}[\nabla(x \bullet \lambda_0)].$$

Let $S \subset W$ denote the simple reflections. To any $s \in S$ one can associate a “wall-crossing” functor Ξ_s by translating to and from an s -wall of the fundamental alcove. Consider the “antispherical” right $\mathbb{Z}[W]$ -module $\mathbb{Z}_\varepsilon \otimes_{\mathbb{Z}[W_f]} \mathbb{Z}[W]$, where \mathbb{Z}_ε denotes the sign module for the finite Weyl group W_f (viewed as a right $\mathbb{Z}[W_f]$ -module). This module has a basis (as a \mathbb{Z} -module) consisting of the elements $1 \otimes w$ with $w \in {}^fW$. Then we can reformulate (1.2.1) as an isomorphism

$$(1.2.2) \quad \phi: \mathbb{Z}_\varepsilon \otimes_{\mathbb{Z}[W_f]} \mathbb{Z}[W] \xrightarrow{\sim} [\mathrm{Rep}_0(G)]$$