

quatrième série - tome 54 fascicule 1 janvier-février 2021

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Alessio FIGALLI & David JERISON

*A sharp Freiman type estimate for semisums in two and three
dimensional Euclidean spaces*

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

YVES DE CORNULIER

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2021

P. BERNARD D. HARARI
S. BOUCKSOM C. IMBERT
G. CARON S. MOREL
G. CHENEVIER P. SHAN
A. DUCROS J. SZEFTEL
B. FAYAD S. VŪ NGỌC
G. GIACOMIN G. WILLIAMSON
D. HÄFNER

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France
Case 916 - Luminy
13288 Marseille Cedex 09
Tél. : (33) 04 91 26 74 64
Fax : (33) 04 91 41 17 51
email : abonnements@smf.emath.fr

Tarifs

Abonnement électronique : 428 euros.
Abonnement avec supplément papier :
Europe : 576 €. Hors Europe : 657 € (\$ 985). Vente au numéro : 77 €.

© 2021 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

A SHARP FREIMAN TYPE ESTIMATE FOR SEMISUMS IN TWO AND THREE DIMENSIONAL EUCLIDEAN SPACES

BY ALESSIO FIGALLI AND DAVID JERISON

ABSTRACT. – Freiman’s theorem is a classical result in additive combinatorics concerning the approximate structure of sets of integers that contain a high proportion of their internal sums. As a consequence, one can deduce an estimate for sets of real numbers: “If $A \subset \mathbb{R}$ and $\left| \frac{1}{2}(A + A) \right| - |A| \ll |A|$, then A is close to its convex hull.” In this paper we prove a sharp form of the analogous result in dimensions 2 and 3.

RÉSUMÉ. – Le théorème de Freiman est un résultat classique de la combinatoire additive concernant la structure approximative des ensembles d’entiers qui contiennent une forte proportion de leurs sommes internes. En conséquence, on déduit l’estimée suivante : “Si $A \subset \mathbb{R}$ et $\left| \frac{1}{2}(A + A) \right| - |A| \ll |A|$, alors A est proche de son enveloppe convexe.” Dans cet article, nous prouvons une forme optimale du résultat correspondant en dimensions 2 et 3.

1. Introduction

Given a set $A \subset \mathbb{R}^n$, define the semisum by

$$\frac{1}{2}(A + A) := \left\{ \frac{x+y}{2} : x \in A, y \in A \right\}.$$

Evidently, $\frac{1}{2}(A + A) \supset A$, and for convex sets K , $\frac{1}{2}(K + K) = K$. Also, $\left| \frac{1}{2}(A + A) \right| = |A| > 0$ implies that A is equal to its convex hull $\text{co}(A)$ minus a set of measure zero (see [3, Théorème 6]).

The stability of this statement is a natural question that has already been extensively investigated in the one dimensional case. Indeed, by approximating sets in \mathbb{R} with finite unions of intervals, one can translate the problem to \mathbb{Z} and in the discrete setting the question becomes a well studied problem in additive combinatorics. More precisely, set

$$\delta(A) := \left| \frac{1}{2}(A + A) \right| - |A|,$$

where $|\cdot|$ denotes the outer Lebesgue measure. The following theorem can be obtained as a corollary of a result of G. Freiman [9] about the structure of additive subsets of \mathbb{Z} (see [5] for more details, and also [11] and the references therein for more recent developments on this one dimensional problem):

THEOREM 1.1. – *Let $A \subset \mathbb{R}$ be a measurable set of positive Lebesgue measure, and assume that $\delta(A) < |A|/2$. Then*

$$|\text{co}(A) \setminus A| \leq 2\delta(A).$$

Note that the assumption $\delta(A) < |A|/2$ is necessary, as can be seen by considering the set $A = [0, 1] \cup [R, R + 1]$ with $R \gg 1$.

In [5, Theorem 1.2] we extended Theorem 1.1 to every dimension, but with a dimensional dependence in the exponent (see also [6] for a stability result when one considers the semisum of two different sets). Our result was as follows.

THEOREM 1.2. – *Let $n \geq 2$. There exist computable dimensional constants $\delta_n, C_n > 0$ such that if $A \subset \mathbb{R}^n$ is a measurable set of positive Lebesgue measure with $\delta(A) \leq \delta_n|A|$, then*

$$\frac{|\text{co}(A) \setminus A|}{|A|} \leq C_n \left(\frac{\delta(A)}{|A|} \right)^{\alpha_n}, \quad \text{where } \alpha_n := \frac{1}{8 \cdot 16^{n-2} n! (n-1)!}.$$

Note that the dimensional smallness assumption on $\delta(A)$ is necessary. Indeed, consider $t = 1/2$ and the set

$$A := B_1(0) \cup \{Re_1\}, \quad R \gg 1.$$

Then $|\text{co}(A) \setminus A| \approx R$ is arbitrarily large, while $\delta(A) = |B_{1/2}(\frac{R}{2}e_1)| = 2^{-n}|A|$, hence $\delta_n \leq 2^{-n}$.

The proof in [5] is based on induction on dimension and Fubini-type arguments, and it leads to a bad estimate for the exponent α_n . In fact, we believe that $\alpha_n = 1$, which we formulate more precisely in the following conjecture.

CONJECTURE 1.3. – *Suppose that A is a measurable subset of \mathbb{R}^n , of positive Lebesgue measure. There exist computable constants C_n and $d_n > 0$, depending only on n , such that the following holds: if $\delta(A) \leq d_n|A|$, then*

$$|\text{co}(A) \setminus A| \leq C_n \delta(A).$$

In this paper we introduce a completely new strategy that allows us to prove this *sharp* stability estimate in dimensions 2 and 3.

THEOREM 1.4. – *Conjecture 1.3 is valid for $n \leq 3$.*

The exponent $\alpha_n = 1$ may look surprising at first sight, as most sharp stability results for minimizers of geometric inequalities in dimension $n \geq 2$ hold with the exponent $1/2$. In particular, the best possible stability exponent for the Brunn-Minkowski inequality on convex sets is $1/2$, see [8, 7]. In contrast, our stability inequality with exponent 1 is affine invariant and additive under partitions of the set by convex tilings, and these properties are crucial to the proof. Even though we have stopped at $n = 3$, the proof is by induction

on n and is organized with the hope that parts of it will ultimately apply to the case of general n . There is at least one other stability inequality in which the exponent 1 is optimal in all dimensions, namely the one proved in [4]. (Observe that the exponent 1 becomes natural when looking at critical points instead of minimizers, see for instance [2, Theorem 1.2], but this is a consequence of the different definition of the “deficit” δ .)

Acknowledgments. – AF was partially supported by NSF Grants DMS-1262411 and DMS-1361122, and by ERC Grant “Regularity and Stability in Partial Differential Equations (RSPDE)”. DJ was partially supported by NSF grant DMS 1500771 and a 2018 Simons Fellowship. The authors thank Emanuel Milman for pointing out to them reference [3]. The authors are very grateful to an anonymous referee, who found a major error in the first version of this paper. In that version, we asserted that the conjecture was true in all dimensions, based on the false claim that inequality (2.4) was valid with the exponent 1 rather than n . In this revision we had to make substantial changes that allow us to prove the conjecture in dimensions 2 and 3. In Remark 2.5 we briefly comment on the case $n \geq 4$.

2. Proof of Theorem 1.4

As the reader will see, many of the arguments for the proof of Theorem 1.4 are valid in any dimension. For this reason we shall work with a generic n for most of the proof, and we shall use some geometric considerations specific to $n = 2$ and $n = 3$ only towards the end.

Basic considerations

Since Theorem 1.4 is known for $n = 1$ (see Theorem 1.1), we can assume that $n \geq 2$ and, by induction on dimension, we can also assume that Theorem 1.4 holds in dimension $n - 1$.

Denote the convex hull of A by $K := \text{co}(A)$. Since the theorem is affine invariant, after dilation we can assume, with no loss of generality, that $|A| = 1$. Assuming that $\delta(A) \ll 1$, it follows by [1] and/or [5, Theorem 1.2] that ⁽¹⁾

$$(2.1) \quad \mu := |K \setminus A| \ll 1.$$

In particular, $1 \leq |K| \leq 2$. Therefore, using the lemma of F. John [10], up to an affine transformation with Jacobian bounded from above and below by a dimensional constant, we can assume that K satisfies

$$(2.2) \quad B_{1/\sqrt{n}} \subset K \subset B_{\sqrt{n}}$$

for balls of radius $1/\sqrt{n}$ and \sqrt{n} centered at the origin.

By approximation, ⁽²⁾ we can assume the set A is compact and that ∂K consists of finitely many polygonal faces. In particular, $\frac{1}{2}(A + A)$ is compact, hence measurable. Furthermore,

⁽¹⁾ Although this estimate can be deduced as a consequence of [1], that result does not provide computable constants, as the proof is based on a contradiction argument relying on compactness.

⁽²⁾ One way to define a suitable approximation is to consider a sequence of finite sets $V_k \subset V_{k+1} \subset A$ such that the polyhedra $P_k = \text{co}(V_k)$ satisfy $|P_k| \rightarrow |\text{co}(A)|$ as $k \rightarrow \infty$, and a sequence of compact subsets $A'_k \subset A$ such that $|A'_k| \rightarrow |A|$ as $k \rightarrow \infty$. Then let $A_k := V_k \cup [A'_k \cap (1 - 1/k)P_k]$. Since $|A_k| \rightarrow |A|$, it suffices to prove the estimate of Theorem 1.4 for A_k and then let $k \rightarrow \infty$.