

*quatrième série - tome 49      fascicule 6      novembre-décembre 2016*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Jayadev S. ATHREYA & Alex ESKIN & Anton ZORICH

*Right-angled billiards and volumes of moduli spaces of quadratic  
differentials on  $\mathbb{C}P^1$*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> janvier 2016

N. ANANTHARAMAN I. GALLAGHER  
P. BERNARD B. KLEINER  
E. BREUILLARD E. KOWALSKI  
R. CERF M. MUSTAȚĂ  
A. CHAMBERT-LOIR L. SALOFF-COSTE

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

### Édition / *Publication*

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

### Abonnements / *Subscriptions*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

### Tarifs

Europe : 519 €. Hors Europe : 548 €. Vente au numéro : 77 €.

---

© 2016 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret  
Périodicité : 6 n<sup>os</sup> / an

# RIGHT-ANGLED BILLIARDS AND VOLUMES OF MODULI SPACES OF QUADRATIC DIFFERENTIALS ON $\mathbb{CP}^1$

BY JAYADEV S. ATHREYA, ALEX ESKIN AND ANTON ZORICH  
WITH AN APPENDIX BY JON CHAIKA

---

**ABSTRACT.** – We use the relation between the volumes of the strata of meromorphic quadratic differentials with at most simple poles on  $\mathbb{CP}^1$  and counting functions of the number of (bands of) simple closed geodesics in associated flat metrics with singularities to prove a very explicit formula for the volume of each such stratum conjectured by M. Kontsevich a decade ago.

Applying ergodic techniques to the Teichmüller geodesic flow we obtain quadratic asymptotics for the number of (bands of) closed trajectories and for the number of generalized diagonals in almost all right-angled billiards.

**RÉSUMÉ.** – Nous utilisons le lien entre les volumes des strates de différentielles méromorphes quadratiques avec des pôles simples sur  $\mathbb{CP}^1$  et les fonctions de comptage du nombre de (cylindres de) géodésiques fermées simples pour la métrique plate associée afin de démontrer une formule très explicite pour le volume des strates, conjecturée par M. Kontsevich il y a une décennie.

En appliquant des techniques ergodiques au flot géodésique de Teichmüller nous obtenons une asymptotique quadratique pour le nombre de (bandes de) trajectoires fermées et le nombre de diagonales généralisées dans presque tout billard à angles « droits ».

## 1. Introduction

Motivated by the study of computing asymptotics for the number of generalized diagonals and for the number of closed billiard trajectories in right-angled polygons, we were naturally led to questions on Masur-Veech volumes of strata of moduli spaces of quadratic differentials on  $\mathbb{CP}^1$ . Our main result, explicitly computing these volumes, resolves a conjecture of M. Kontsevich.

---

J.S.A. is partially supported by NSF grants DMS 1069153, NSF grants DMS 1107452, 1107263, 1107367 RNMS: GEometric structures And Representation varieties (the GEAR Network), and NSF CAREER grant DMS 1351853. A.E. is partially supported by NSF grants DMS 0244542 and DMS 0604251. A.Z. is partially supported by IUF and by ANR “GeoDyM”.

### 1.1. Volumes of moduli spaces of quadratic differentials

**THEOREM 1.1 (Kontsevich Conjecture).** – *The volume of any stratum  $\mathcal{Q}_1(d_1, \dots, d_k)$  of meromorphic quadratic differentials with at most simple poles on  $\mathbb{CP}^1$  (i.e.,  $d_i \in \{-1; 0\} \cup \mathbb{N}$  for  $i = 1, \dots, k$ , and  $\sum_{i=1}^k d_i = -4$ ) is equal to*

$$(1.1) \quad \text{Vol } \mathcal{Q}_1(d_1, \dots, d_k) = 2\pi^2 \cdot \prod_{i=1}^k v(d_i)$$

(where all the zeroes and poles are “named”).

Here, the function  $v$  is defined on integers  $n$  greater than or equal to  $-1$  by

$$(1.2) \quad v(n) := \frac{n!!}{(n+1)!!} \cdot \pi^n \cdot \begin{cases} \pi & \text{when } n \text{ is odd} \\ 2 & \text{when } n \text{ is even} \end{cases}$$

for  $n = -1, 0, 1, 2, 3, \dots$ , and the double factorial  $n!! = n \cdot (n-2) \cdots$  is the product of all even (respectively odd) positive integers smaller than or equal to  $n$ . By convention we set

$$(-1)!! = 0!! = 1,$$

which implies that

$$v(-1) = 1 \quad \text{and} \quad v(0) = 2.$$

This formula for the volume (up to some normalization factor) was conjectured by M. Kontsevich about ten years ago. It is much simpler than the formula for the volumes of the strata of Abelian differentials found by A. Eskin and A. Okounkov [25].

When this paper was written, there was not a single stratum of quadratic differentials for which the explicit volume was known, though an algorithm of computation was presented in [26]. In addition to this work, there is some very recent progress in evaluation of volumes of low-dimensional strata in genera different from 0. Rigorous formal methods used in [29] (in particular, implementation of the algorithm [26]) are confirmed by independent numerical experiments [12]. However, any known approach involves significant computer-assisted computations, and is limited to volumes of strata of sufficiently small dimension, while Theorem 1.1 provides a simple formula for *all* strata in genus 0.

Returning to our original motivation, we obtain as an important application of Theorem 1.1 asymptotics for the number of closed trajectories and for the number of generalized diagonals in right-angled polygons (see §1.3 below). This choice is particularly natural in the context of this paper since we have to solve an analogous problem for quadratic differentials and to compute the corresponding Siegel-Veech constants  $c_{\mathcal{Q}}$  for the strata of quadratic differentials in genus 0 anyway: it makes part of the proof of Theorem 1.1. This theorem also immediately provides asymptotics for certain Hurwitz numbers, see §1.2. Another example of applications is discussed in [14] where the values of volumes and the related Siegel-Veech constants are used to compute Lyapunov exponents of the Hodge bundle over hyperelliptic loci in the strata of quadratic differentials and to compute the diffusion rate for interesting families of generalized wind-tree billiards [13].

**Strategy of the proof**

We start by solving the counting problems for quadratic differentials. The Siegel-Veech constant  $c_{\text{area}}$  responsible for the exact quadratic asymptotics of the weighted number of bands of regular closed geodesics on almost any flat sphere in a given stratum  $\mathcal{Q}(d_1, \dots, d_n)$  of meromorphic quadratic differentials with at most simple poles on  $\mathbb{CP}^1$  was recently computed in [16],

$$(1.3) \quad c_{\text{area}}(\mathcal{Q}(d_1, \dots, d_n)) = -\frac{1}{8\pi^2} \sum_{j=1}^n \frac{d_j(d_j + 4)}{d_j + 2}.$$

Developing techniques elaborated in [21] for the strata of Abelian differentials and using the further results from [8] and [39] on the *principal boundary* of the strata of quadratic differentials we express the Siegel-Veech constant  $c_{\text{area}}$  in genus 0 in terms of the ratio of the volumes of appropriate strata,

$$(1.4) \quad c_{\text{area}}(\mathcal{Q}(d_1, \dots, d_n)) = \frac{\text{explicit polynomial in volumes of simpler strata}}{\text{Vol}(\mathcal{Q}(d_1, \dots, d_n))}.$$

In this way we obtain a series of identities on the volumes of the strata of meromorphic quadratic differentials with at most simple poles in genus zero. The resulting identities recursively determine the volumes of all strata. The proof of Theorem 1.1, given in §5, consists in verifying that the expression (1.1) for the volume satisfies the combinatorial identities implied by (1.3) and (1.4). Part of this verification is performed in Appendix A.

REMARK 1.2 (Normalization conventions). – Note that the convention that all zeroes and poles are “named” affects the normalization: we compute the volumes of the corresponding covers over strata with “anonymous” singularities. For example, the stratum  $\mathcal{Q}(1, -1^5)$  of quadratic differentials with “anonymous” zeroes and poles is isomorphic to the stratum  $\mathcal{H}(2)$  of holomorphic Abelian differentials; by convention the volume elements are chosen to be invariant under this isomorphism. However, by (1.1) we have

$$\text{Vol } \mathcal{Q}_1(1, -1^5) = 2\pi^2 \cdot v(1) \cdot (v(-1))^5 = 2\pi^2 \cdot \frac{\pi^2}{2} \cdot 1^5 = 5! \cdot \frac{\pi^4}{120} = 5! \cdot \text{Vol } \mathcal{H}_1(2),$$

which corresponds to 5! ways to *give names* to five simple poles.

Similarly,

$$\text{Vol } \mathcal{Q}_1(2, -1^6) = 2\pi^2 \cdot v(2) \cdot (v(-1))^6 = 2\pi^2 \cdot \frac{4\pi^2}{3} \cdot 1^6 = \frac{6!}{2!} \cdot \frac{\pi^4}{135} = \frac{6!}{2!} \cdot \text{Vol } \mathcal{H}_1(1, 1).$$

This time there is an extra factor  $\frac{1}{2!}$  responsible for *forgetting the names* of the two zeroes of  $\mathcal{H}(1, 1)$ .

**1.2. Counting pillowcase covers**

One of the ways to compute the volumes of the strata of Abelian or quadratic differentials (actually, the only one before the current paper) is to count *square-tiled surfaces* or *pillowcase covers*, see [25], [26], [27], [51]. In the current paper we proceed in the other direction: we first compute volumes of the strata by an alternative method, and then, as a corollary of volume computation, we get an explicit expression for the leading term of the function counting associated pillowcase covers, when the degree of the cover tends to infinity.