Astérisque **330**, 2010, p. 1–11

## BANACH $\ell$ -ADIC REPRESENTATIONS OF *p*-ADIC GROUPS

by

Marie-France Vigneras

Abstract. — Let  $p \neq \ell$  be two distinct prime numbers, let F be a p-adic field and let E be an  $\ell$ -adic field. We prove that the smooth part and the completion are inverse equivalences of categories between the category of admissible Banach unitary E-representations of GL(n, F) and the category of admissible smooth E-representations of GL(n, F) equipped with a commensurability class of lattices. We formulate the  $\ell$ -adic local Langlands correspondence as a canonical bijection between the n-dimensional  $\ell$ -adic representations of the absolute Galois group  $\operatorname{Gal}_F$ and the topologically irreducible admissible Banach unitary  $\ell$ -adic representations of GL(n, F).

*Résumé* (Représentations  $\ell$ -adiques de groupes p-adiques). — Soient  $p \neq \ell$  deux nombres premiers distincts, soit F un corps p-adique et soit E un corps  $\ell$ -adique. Nous démontrons que la partie lisse et la complétion définissent des équivalences de catégories inverses l'une de l'autre entre la catégorie des représentations admissibles de Banach unitaires de GL(n, F) sur E et la catégorie des représentations lisses admissibles de GL(n, F) sur E munies d'une classe de commensurabilité de réseaux. Nous formulons la correspondance de Langlands locale  $\ell$ -adique comme une bijection canonique entre les représentations  $\ell$ -adiques de dimension n du groupe de Galois absolu  $\operatorname{Gal}_F$  et les représentations topologiquement irréductibles admissibles de Banach unitaires  $\ell$ -adique de GL(n, F).

## 1. Introduction

Let p be a prime number, let F be a finite extension of  $\mathbf{Q}_p$  or a field of Laurent series k((T)) over a finite field k of characteristic p, let  $\overline{F}$  be an algebraic closure of F and let n be an integer  $\geq 1$ .

For any topological field C, the continuous representations of GL(n, F) on topological vector spaces over C are interesting for their applications in arithmetic, geometry or physics, via the theory of L-functions associated to automorphic representations. When C varies, the theories of C-representations of GL(n, F) present simultaneously

<sup>2010</sup> Mathematics Subject Classification. — 11F70, 11F80, 11F85.

Key words and phrases. — Banach  $\ell$ -adic representation, reductive *p*-adic group, admissible representation, smooth representation, lattice.

strong similarities and strong different features but the Langlands insight, when C is the complex field, to use the smooth complex representations of  $\operatorname{Gal}_F = \operatorname{Gal}(\overline{F}/F)$ as a classifying scheme, seems to extend to other fields.

Why moving the coefficient field C? There are many reasons.

1) The representations of  $\operatorname{Gal}_F$  appearing naturally are not smooth complex. In the étale cohomology of proper smooth algebraic varieties, they are continuous  $\ell$ -adic representations on finite dimensional vector spaces V over finite extensions  $E/\mathbf{Q}_{\ell}$ , for a prime number  $\ell$ . By a reduction of a stable  $O_E$ -lattice of V, they give smooth mod  $\ell$ -representations over the residual field of E.

2) The local Langlands correspondence for GL(n, F), over any algebraically closed field R of characteristic different from p, is a bijection

$$\pi \leftrightarrow (\rho, N)$$

between the equivalence classes of the smooth irreducible *R*-representations  $\pi$  of GL(n, F) and of the pairs  $(\rho, N)$  where  $\rho$  is a *n*-dimensional smooth semi-simple *R*-representation of the Weil group  $W_F$  and N a nilpotent endomorphism of the space of  $\rho$  such that  $\rho(w)N = N|w|\rho(w)$  where |?| is the unramified *R*-character of  $W_F$  sending a geometric Frobenius to q, the order of the residual field of F.

Our purpose is to obtain a local Langlands correspondence for continuous  $\ell$ -adic representations.

**Theorem 1.** — Let  $\ell$  be a prime number different from p. The  $\ell$ -adic local Langlands correspondence for GL(n, F) is a canonical bijection between the equivalence classes of

a) n-dimensional continuous  $\ell$ -adic representations of  $\operatorname{Gal}_F$  with a semi-simple action of the Frobenius,

b) topologically irreducible admissible Banach unitary  $\ell$ -adic representations of GL(n, F).

This theorem<sup>(1)</sup> is motivated by the fascinating work and conjectures of Christophe Breuil on the *p*-adic local Langlands correspondence, where topologically irreducible admissible Banach unitary *p*-adic representations of  $GL(2, \mathbf{Q}_p)$  appear naturally.

With the existing literature, one translates the local Langlands complex correspondence for GL(n, F) into a canonical bijection between the isomorphism classes of a) and of

c) Irreducible smooth  $\overline{\mathbf{Q}}_{\ell}$ -representations of GL(n, F) with a stable lattice.

Indeed, as is well known,

(i) The smooth complex local Langlands correspondence  $LL(\rho, N)$  twisted by a suitable unramified character,

 $(\rho, N) \leftrightarrow LL(\rho, N) \otimes |\det?|^{-(n-1)/2},$ 

<sup>&</sup>lt;sup>(1)</sup> Proved in a letter to Breuil in september 2003, and announced in the Emmy Noether lectures 2005 of Goettingen.

called the smooth complex local Hecke correspondence, is Aut C-equivariant [H prop.6].

(ii) Transporting the correspondence (i) with an algebraic isomorphism  $j : \mathbf{C} \simeq \mathbf{Q}_{\ell}$ , we obtain the smooth local Hecke  $\overline{\mathbf{Q}}_{\ell}$ -correspondence, which does not depend on the choice of the isomorphism j.

(iii) N disappears when one considers continuous  $\overline{\mathbf{Q}}_{\ell}$ -representations of  $W_F$  instead of smooth  $\overline{\mathbf{Q}}_{\ell}$ -representations. The pairs  $(\rho, N)$  are in bijection

$$(\rho, N) \leftrightarrow \sigma$$

with the *n*-dimensional  $\ell$ -adic representations  $\sigma$  of  $W_F$  with a semi-simple action of the Frobenius. The reason is that the kernel of the natural morphism  $t: I_F \to \mathbf{Z}_{\ell}$  is a profinite group prime to  $\ell$ . There is a nilpotent endomorphism N of the space of  $\sigma$ such that  $\sigma(?) = \exp(t(?)N)$  on a subgroup of finite index of  $I_F$  [8].

(iv) The *n*-dimensional  $\ell$ -adic representation  $\sigma$  of  $W_F$  in (iii) extends by continuity to an  $\ell$ -adic representation of  $\operatorname{Gal}_F$  if and only if  $\rho$  has a bounded image (i.e. the values of determinants of the irreducible components of  $\rho$  are units) [8].

(v)  $\rho$  has a bounded image if and only if  $\pi = LL(\rho, N)$  is integral [10, §1.4]; moreover all stable lattices in  $\pi$  are commensurable [11, Theorem 1].

Our task is to show that the completion with respect to a stable lattice gives a bijection between the isomorphism classes of b) and of c).

The beginning of the proof is valid for any locally profinite group G, with a countable fundamental system of neighborhoods of the unit, consisting of open profinite groups of pro-order not divisible by  $\ell$  (Section 2). We prove (Theorem 2.12) that the completion and the smooth part induce equivalences of categories between the category  $\mathscr{M}_{\ell}(G)^{\mathrm{adm}}$  of admissible smooth  $\ell$ -adic representations of G equipped with a commensurability class of lattices, and the category  $\mathscr{B}_{\ell}(G)^{\mathrm{adm}}$  of admissible Banach unitary  $\ell$ -adic representations of G.

Then we consider the group of rational points  $G_F$  of any reductive connected group over a local non Archimedean field F of residual characteristic  $p \neq \ell$  (Section 3). We prove (Theorem 3.6) that the completion and the smooth part induce equivalences of categories between the category  $\operatorname{Mod}_{\overline{\mathbf{Q}}_{\ell}}^{\operatorname{int},\mathrm{fl}}(G_F)$  of integral smooth  $\overline{\mathbf{Q}}_{\ell}$ -representations of  $G_F$  of finite length and the category  $\mathscr{B}_{\ell}(G_F)^{\operatorname{adm},\mathrm{fl}}$  of admissible Banach unitary  $\ell$ -adic representations of topological finite length of  $G_F$ . We deduce the wanted bijection between the isomorphism classes of b) and c) by restricting to irreducible representations and choosing  $G_F = GL(n, F)$ .

A natural question was raised by the referee: Is a topologically irreducible Banach unitary  $\ell$ -adic representation of  $G_F$  always admissible ? L. Clozel noticed that the examples of B. Diarra [5, th. 4] (van Rooj), give examples of topologically irreducible representations  $V \in \mathscr{B}_E(GL(1, F))$  where any non zero intertwining operator is bijective, which are not admissible.

## 2.

**2.1. The two categories.** — Let  $\ell \neq p$  be two distinct prime numbers, let  $E/\mathbf{Q}_{\ell}$  be a finite extension of ring of integers  $O_E$ , of uniformizer  $p_E$ , and of residual field  $k_E$ , and let G be a topological group admitting a *countable* fundamental system of neighborhoods of the unit consisting of open  $pro-\ell'$ -subgroups (profinite subgroups of pro-order *prime to*  $\ell$ ).

After having recalled some definitions and properties concerning the representations of the group G on E-vector spaces, we will introduce the two categories of representations  $\mathscr{M}_E(G)$  and  $\mathscr{B}_E(G)$  which will be compared in this paper.

Let  $\operatorname{Mod}_E$  be the category of E-vector spaces and let  $M \in \operatorname{Mod}_E$  non zero. A line in M is a subspace of dimension 1. A lattice L in M is a  $O_E$ -submodule of M which contains no line and contains a basis of M over E. Note that a quotient of a lattice may contain a line. When the dimension of M over E is countable, a lattice L in Mis a free  $O_E$ -submodule of M generated by a basis of M over E [9, I Appendice C.5]. Two lattices L, L' in M are commensurable when there exists an element  $a \in O_E$  such that  $aL \subset L', aL' \subset L$ . We denote by [L] the commensurability class of L.

**Remark 2.1.** — An  $O_E$ -submodule L of  $M \in Mod_E$  is a lattice in M if and only if any non zero element  $m \in M$  satisfies the two conditions:

a) there exists an integer  $n \in \mathbf{N}$  such that  $\ell^n m$  belongs to L,

b) there exists an integer  $n \in \mathbf{N}$  such that  $\ell^{-n}m$  does not belong to L.

Two lattices L, L' in M are commensurable if and only if there exists an integer  $n \in \mathbf{N}$  such that  $\ell^n L \subset L', \ \ell^n L' \subset L$ .

A representation (= a linear action) of G on M is called *admissible* when  $\dim_E M^H < \infty$ , for any open pro- $\ell'$ -subgroup H of G, where  $M^H \in \operatorname{Mod}_E$  is the subspace of H-invariant vectors of M. The representation M is called *irreducible* when  $M \neq 0$  and 0 and M are the only G-stable subspaces of M, finitely generated when M is a finitely generated EG-module, of finite length when there exists a finite G-stable filtration  $0 \subset M_1 \subset \cdots \subset M_n = M$  with *irreducible quotients*. The length of the filtration and the isomorphism classes of the quotients, up to the order, do not depend on the choice of the filtration.

A lattice L in the representation of G on M will always be a G-stable lattice in M; the lattice will be called *finitely generated* when it is a finitely generated  $O_EG$ -module. A representation of G on M containing a lattice is called *integral* (we do not suppose that the lattice is  $O_E$ -free as in [9]). There exist finitely generated lattices in a finitely generated integral representation; they form a commensurability class, and any lattice contains a finitely generated lattice.

A continuous E-representation of G is a topological Hausdorff E-vector space M equipped with a continuous action of G, i.e. such that the map  $(g, v) \to gv : G \times M \to M$  is continuous. It is called *topologically irreducible* when  $M \neq 0$  and 0 and M are the only closed G-stable subspaces of M. It is called of *finite topological length* when

 $\mathbf{5}$ 

there exists a finite filtration by G-stable closed subspaces  $0 \subset M_1 \subset \cdots \subset M_n = M$ with topologically irreducible quotients.

The category  $\mathscr{C}_E(G)$  of continuous representations of G on topological Hausdorff complete E-vector spaces with continuous G-equivariant E-linear morphisms, called intertwinning operators, contains the subcategory  $\operatorname{Mod}_E(G)$  of smooth representations and the subcategory  $\mathscr{B}_E(G)$  of Banach unitary representations, defined below. We indicate by the upper index adm or fl or adm, fl or int or int, fl the full subcategories representations which are admissible or of finite topological length or admissible and of finite topological length or integral or integral and of finite topological length. Example:  $\mathscr{C}_E(G)^{adm}$ ,  $\operatorname{Mod}_E(G)^{adm}$ ,  $\mathscr{B}_E(G)^{adm}$  for admissible representations.

A representation of G on an E-vector space W is smooth when the stabilizer in G of any vector of W is open; this is simply a continuous representation of G on W when W is equipped with the discrete topology. The category  $Mod_E(G)$  of smooth E-representations of G, with morphisms the G-equivariant E-linear maps, is a full subcategory of  $\mathscr{C}_E(G)$ .

A Banach unitary E-representation V of G is a Hausdorff complete topological E-vector space with a topology given by a norm, equipped with a continuous action of G which respects the norm. A unit ball of V is  $L = \{v \in V : ||v|| \le 1\}$  for some norm  $v \mapsto ||v||$  on V defining the topology [Sch I.3, III]; it is a lattice in V. The unit balls of two norms on V giving the same topology are commensurable.

An *E*-linear map  $f : V_1 \to V_2$  between two Banach *E*-vector spaces  $V_1, V_2$  is continuous if and only if there exists some non zero  $a \in E$  such that  $f(L_1) \subset af(L_2)$ for some unit balls  $L_1, L_2$  of  $V_1, V_2$  [Sch I.3.1]. The topology quotient topology on the image of f is the topology induced by  $V_2$  if and only if  $f(L_1)$  and  $L_2 \cap f(V_1)$  are commensurable (this does not depend on the choice of the unit balls  $L_1, L_2$ ). When f is continuous and bijective, the inverse of f is continuous [Sch I.8.7].

We will compare  $\mathscr{B}_E(G)$  with the category  $\mathscr{M}_E(G)$  of smooth *E*-representations *W* of *G* equipped with a commensurability class [L] of lattices; a morphism  $(W, [L]) \rightarrow$ (W', [L']) is a morphism  $f: W \rightarrow W'$  in  $\operatorname{Mod}_E(G)$  such that  $f(L) \subset aL'$  for some  $a \in E$ . The pair (W, [L]) is called admissible or of finite length when *W* is admissible or of finite length, and  $\mathscr{M}_E(G)^{\operatorname{adm}}$  or  $\mathscr{M}_E(G)^{fl}$  is the full subcategory of admissible or of finite length pairs in  $\mathscr{M}_E(G)$ .

**2.2. The two functors.** — We introduce two natural functors in opposite directions between the categories  $\mathscr{M}_E(G)$  and  $\mathscr{B}_E(G)$ .

There is the natural functor  $\mathscr{C}_E(G) \to \operatorname{Mod}_E(G)$  sending  $M \in \mathscr{C}_E(G)$  to its *smooth* part

$$M^{\infty} := \cup_H M^H,$$

for all open pro- $\ell'$ -subgroups H of G. When  $V \in \mathscr{B}_E(G)$  is a Banach unitary representation of G, the smooth part  $L^{\infty} = V^{\infty} \cap L$  of a unit ball L of V is a lattice of  $V^{\infty}$ . Two unit balls of V are commensurable and their smooth parts are commensurable, hence  $(V^{\infty}, [L^{\infty}]) \in \mathscr{M}_E(G)$  is well defined. A continuous morphism  $f: V_1 \to V_2$  of Banach unitary E-representations of G with unit balls  $L_1, L_2$ , restricts to a morphism  $f^\infty:(V_1^\infty,[L_1^\infty])\to (V_2^\infty,[L_2^\infty]).$  We get a functor

$$\mathscr{B}_E(G) \to \mathscr{M}_E(G).$$

In the opposite direction there is the natural functor

 $\mathscr{M}_E(G) \to \mathscr{B}_E(G)$ 

sending (W, [L]) to the *completion* of W for the L-adic topology [Sch 7.5]:

$$\hat{W}_L := \varprojlim_n W/\ell^n L \simeq E \otimes_{O_E} \hat{L} , \ \hat{L} := \varprojlim_n L/\ell^n L.$$

Any element  $v \in \hat{W}_L$  is written

(1) 
$$v = (w_n + \ell^n L)_n, w_n \in W, w_{n+1} \in w_n + \ell^n L,$$

for all  $n \in \mathbf{N}$ . The lattice  $\hat{L}$  is a unit ball of  $\hat{W}_L$  for the gauge norm  $||v|| = \inf_{a \in E, v \in a\hat{L}} |a|$ . The completions of W defined by two commensurable lattices of W are the same. The group G acts naturally on  $\hat{W}_L$ , for  $g \in G$  and v as above,

$$gv = (gw_n + \ell^n L)_{n \in \mathbf{N}},$$

and  $\hat{W}_L$  is a Banach unitary *E*-representation of *G* of unit ball  $\hat{L}$ , well defined by (W, [L]). A morphism  $f: (W, [L]) \to (W', [L'])$  in  $\mathscr{M}_E(G)$  extends by continuity to an intertwinning operator  $\hat{f}: \hat{W}_L \to \hat{W}'_{L'}$ .

**Remark 2.2.** — The map  $W \mapsto \hat{W}_L$  sending w to  $(w + \ell^n L)_{n \in \mathbb{N}}$  is injective, because L contains no line. We will identify W with its image in  $\hat{W}_L$ .

**2.3.** — To study the two functors, smooth part and completion, between  $\mathcal{M}_E(G)$  and  $\mathcal{B}_E(G)$ , the key point is the exactness of the *H*-invariants functor.

**Proposition 2.3**. — Let H be any open  $pro-\ell'$ -subgroup of G. The H-invariants functor

$$M \mapsto M^H : \mathscr{C}_E(G) \to \operatorname{Mod}_E$$

is exact.

*Proof.* — This is well known for the subcategory  $\operatorname{Mod}_E(G)$  of smooth representations in  $\mathscr{C}_E(G)$ . The exactness results from the existence of a Haar  $O_E$ -measure dg on Gsuch that the volume  $\operatorname{vol}(H, dg)$  of H is a unit in  $O_E$ . The function  $e_H$  equal to  $\operatorname{vol}(H, dg)^{-1}$  on H and 0 on G - H, is an idempotent in the convolution algebra  $C_c^{\infty}(G; O_E)$  of locally constant compactly supported functions  $G \to O_E$ , for the Haar measure dg. The idempotent  $e_H$  acts on  $M \in \mathscr{C}_E(G)$ , as follows. One chooses a decreasing sequence of normal subgroups  $H_n$  of H of finite index such that  $\cap_{n \in \mathbb{N}} H_n$ is trivial, and a system of representatives  $X_n$  in H of  $H/H_n$ . The continuity of the action of G on M implies that the sequence

$$v_n = [H:H_n]^{-1} \sum_{g \in X_n} gv$$

7

converges to a unique element  $e_H * v$  in the Hausdorff complete space M. This element  $e_H * v$  does not depend on the choice of  $(H_n, X_n)_{n \in \mathbb{N}}$  and clearly  $v \mapsto e_H * v$  is a linear projector  $M \to M^H$  of its *H*-invariants.

**Corollary 2.4.** — The smooth part functor  $M \mapsto M^{\infty} : \mathscr{C}_E(G) \to \operatorname{Mod}_E(G)$  is exact.

**Proposition 2.5.** — A Banach unitary E-representation V of G is equal to the closure of its smooth part  $V^{\infty}$ .

*Proof.* — Let v be an arbitrary element of V and let L be a unit ball of V. For any integer  $n \ge 1$ , there is an open pro- $\ell'$ -subgroup  $H_n$  of G such that  $H_n v \subset v + \ell^n L$ , by the continuity of the action of G on V. The element  $e_{H_n} * v$  is fixed by  $H_n$  and belongs to  $v + \ell^n L$ . The element  $(e_{H_n} * v + \ell^n L)_{n \in \mathbb{N}}$  belongs to the closure of  $V^{\infty}$  and is equal to v.

**Corollary 2.6.** — The smooth part functor  $\mathscr{B}_E(G) \to \mathscr{M}_E(G)$  is fully faithful.

*Proof.* — For i = 1, 2, let  $V_i \in \mathscr{B}_E(G)$  with unit ball  $L_i$ . The embedding  $V_i^{\infty} \to (V_i^{\infty})_{L_i^{\infty}}$  extends by continuity to an isomorphism  $\tau_i : V_i \to (V_i^{\infty})_{L_i^{\infty}}$  in  $\mathscr{B}_E(G)$  by the Proposition 2.5 and its proof. We deduce that arbitrary intertwinning operators  $\phi : (V_1^{\infty}, [L_1^{\infty}]) \to (V_2^{\infty}, [L_2^{\infty}])$  and  $f : V_1 \to V_2$  satisfy

$$\phi = (\tau_2^{-1} \hat{\phi} \tau_1)^{\infty} , \quad f = \tau_2^{-1} (f^{\infty}) \tau_1 .$$

We show that the completion commutes with the *H*-invariants.

**Proposition 2.7.** — Let V be the completion of an integral smooth E-representation W of G with respect to a lattice L, and let H be an open  $\text{pro-}\ell'$ -subgroup of G. The H-invariants  $V^H$  of V is equal to the closure of  $W^H$  in V,

$$V^H = \overline{W^H}.$$

*Proof.* — For X = W, L or V, we have  $e_H * X = X^H$ . Let  $v = (w_n + \ell^n L)_{n \in \mathbb{N}}$ be an element of V as in (1). Then  $e_H * w_{n+1} \in e_H * w_n + \ell^n L$ , and  $e_H * v = (e_H * w_n + \ell^n L)_{n \in \mathbb{N}}$ .

**Corollary 2.8.** — An admissible smooth E-representation of G with a commensurability class of lattices is equal to the smooth part of its completion.

*Proof.* — When the representation W is admissible, the *E*-vector space  $W^H$  is finite dimensional and already complete, hence  $V^H = W^H$  in the Proposition 2.7.

It is clear that the functor smooth part respects admissible representations, the corollary shows that the completion respects also admissible representations.

**Theorem 2.9.** — The smooth part and completion are inverse equivalences of categories between  $\mathscr{M}_E^{\mathrm{adm}}(G)$  and  $\mathscr{B}_E^{\mathrm{adm}}(G)$ .

*Proof.* — Proposition 2.5, Corollaries 2.6, 2.8.

In particular, the smooth part and the completion induce inverse equivalences of categories between admissible and of finite topological length representations  $\mathscr{M}_{E}^{\mathrm{adm,fl}}(G)$  and  $\mathscr{B}_{E}^{\mathrm{adm,fl}}(G)$ .

We consider now  $\ell$ -adic representations of G. For any finite extensions  $E'/E/\mathbf{Q}_{\ell}$  contained in a fixed algebraic closure  $\overline{\mathbf{Q}}_{\ell}$ , the scalar extension  $s_{E/E'}$  from E to E'

 $\mathscr{C}_E(G) \to \mathscr{C}_{E'}(G)$ 

sends  $M \in \mathscr{C}_E(G)$  to  $M_{E'} := E' \otimes_E M = \bigoplus(e_i \otimes M)$ , for a finite basis  $(e_i)$  of the *E*-vector space E', with the topology induced by M (independent of the choice of the basis) and a morphism  $f: M \to M'$  in  $\mathscr{C}_E(G)$  to  $\mathrm{id}_{E'} \otimes f$ . The inductive limit

$$\mathscr{C}_{\ell}(G) := \lim_{s_{E'/E}} \mathscr{C}_{E}(G)$$

is the category of  $\ell$ -adic representations of G. The scalar extension respects smooth representations, and the inductive limit

$$\operatorname{Mod}_{\ell}(G) := \lim_{s_{E'/E}} \operatorname{Mod}_{E}(G)$$

is the category of smooth  $\ell$ -adic representations of G, which is a (not full) subcategory of the classical category  $\operatorname{Mod}_{\overline{\mathbf{Q}}_{\ell}}(G)$  of smooth  $\overline{\mathbf{Q}}_{\ell}$ -representations of G.

Let R/E be any extension contained in  $\overline{\mathbf{Q}}_{\ell}$  and let  $O_R$  be the ring of integers in R. As the dimension of R/E is countable,  $O_R$  is an  $O_E$ -free module [9, Appendice C, C.4]. We denote by  $L_{O_R} := O_R \otimes_{O_E} L$ , the scalar extension from  $O_E$  to  $O_R$  of an  $O_E$ -module L.

**Lemma 2.10.** — Let  $M \in \mathscr{C}_E(G)$  equipped with a lattice L.

(i) The H-invariants commute with the scalar extension,  $e_H M_R = (e_H M)_R$ ,  $e_H L_{O_R} = (e_H L)_{O_R}$  for any pro- $\ell'$ -subgroup H of G (this is true for any extension R/E of fields of characteristic different from p). In particular, M is admissible if and only if  $M_R$  is admissible.

(ii) The intersection  $L = M \cap L'$  of a lattice L' of  $M_R$  with M is a lattice in M. In particular, if  $M_R$  is integral then M is integral.

(iii) The scalar extension  $L_{O_R}$  of a lattice L in M is a lattice in  $M_R$ . In particular, if M is integral then  $M_R$  is integral.

(iv) Two lattices L, L' of M are commensurable if and only if their scalar extensions  $L_{O_R}, L'_{O_R}$  are commensurable.

*Proof.* — (i) is clear. The other properties are clear using the Remark 2.1 and  $L_{O_R} = \bigoplus_i (e_i \otimes L)$  for a basis  $(e_i)$  of the free  $O_E$ -module  $O_R$ .

**Lemma 2.11.** — The scalar extension  $s_{E'/E}$  from E to a finite extension E' commutes with the smooth part functor  $\mathscr{C}_E(G) \to \operatorname{Mod}_E(G)$  and with the smooth part and completion functors between  $\mathscr{B}_E(G)$  and  $\mathscr{M}_E(G)$ .

*Proof.* — We choose a basis  $(e_i)$  of the free  $O_E$ -module  $O_{E'}$ . The scalar extension  $V_{E'} = \bigoplus_i (e_i \otimes V)$  of the completion V of  $(W, [L]) \in \mathscr{M}_E(G)$  is clearly the completion of the scalar extension  $(W_{E'} = \bigoplus_i (e_i \otimes W), [L_{O_{E'}} = \bigoplus_i (e_i \otimes L)])$  of (W, [L]). The

scalar extension of the *H*-invariants of  $V \in \mathscr{B}_E(G)$  is the *H*-invariants of the scalar extension  $V_{E'}$  (Lemma 2.10).

As the scalar extension  $s_{E'/E}$  from E to a finite extension E' respects admissibility, lattices, commensurability of lattices, Banach spaces (Lemma 2.10), the inductive limit over  $s_{E'/E}$  for all finite extensions  $E'/E/\mathbf{Q}_{\ell}$  contained in  $\overline{\mathbf{Q}}_{\ell}$ , defines the categories

a)  $\mathscr{C}_{\ell}(G)^{\mathrm{adm}}$  of admissible  $\ell$ -adic representations of G,

b)  $\operatorname{Mod}_{\ell}(G)^{\operatorname{int}}$  of integral smooth  $\ell$ -adic representations,

b)  $\mathscr{M}_{\ell}(G)$  of smooth  $\ell$ -adic representations of G equipped with a commensurability class of lattices,

c)  $\mathscr{B}_{\ell}(G)$  of Banach unitary  $\ell$ -adic representations of G.

We define the completion and smooth part functors between  $\mathcal{M}_{\ell}(G)$  and  $\mathcal{B}_{\ell}(G)$  using the Lemma 2.11.

**Theorem 2.12.** — The completion and smooth part functors induce equivalence of categories between the categories  $\mathscr{M}_{\ell}(G)^{\mathrm{adm}}$  and  $\mathscr{B}_{\ell}(G)^{\mathrm{adm}}$ .

Proof. — Theorem 2.9.

Let  $G_F$  be the group of rational points of a connected reductive group over a local non Archimedean field F of residual characteristic p. The group  $G_F$  is a locally pro-p-group. As before  $E/\mathbf{Q}_{\ell}$  is a finite extension contained in  $\overline{\mathbf{Q}}_{\ell}$  and  $\ell \neq p$ .

3.

**Proposition 3.1.** — Let  $R/R_o$  be any extension of fields of characteristic different from p. Then  $W \in Mod_{R_o}(G_F)$  has finite length if and only if  $W_R \in Mod_R(G_F)$  has finite length.

*Proof.* — [9, II.4.3.c].

**Proposition 3.2.** — Any finite length smooth representation W of  $G_F$  over a field of characteristic different from p is admissible.

*Proof.* — This is proved in [9, II.2.8] when the field is algebraically closed. The scalar extension is not sensitive to admissibility and finite length (Lemma 2.10, Proposition 3.1) for any extension of fields of characteristic different from p.

**Proposition 3.3.** — The lattices in an integral finite length representation  $W \in Mod_E(G_F)$  are commensurable (hence finitely generated).

*Proof.* — This is proved [11, th.1] when the field is  $\overline{Q}_{\ell}$ . The scalar extension is not sensitive to integrality, commensurability of lattices, and finite length, (Proposition 2.10, Proposition 3.1).

**Remark 3.4**. — One cannot replace "finite length" by "admissible" in the Proposition 3.3.

**Lemma 3.5.** — The category of smooth  $\overline{\mathbf{Q}}_{\ell}$ -representation of  $G_F$  of finite length is equal to the category of smooth  $\ell$ -adic representations of  $G_F$  of finite length,

$$\operatorname{Mod}_{\overline{\mathbf{Q}}_{\ell}}(G_F)^{fl} \simeq \operatorname{Mod}_{\ell}(G_F)^{fl}.$$

Proof. — Let  $W \in \text{Mod}_{\overline{\mathbf{Q}}_{\ell}}(G_F)^{fl}$ . There exists a finite extension  $E/\mathbf{Q}_{\ell}$  and  $W_E \in \text{Mod}_E(G_F)^{fl}$  such that W is the scalar extension of  $W_E$ . When W is irreducible, this is proved in [9, II.4.7]. In general, let H be an open pro-*p*-subgroup of  $G_F$  such that the length of W is equal to the length of the module  $e_H W$  over the Hecke algebra  $\text{End}_{\overline{\mathbf{Q}}_{\ell}G_F} \overline{\mathbf{Q}}_{\ell}[G_F/H]$ . Let  $(w_i)$  be a finite  $\overline{\mathbf{Q}}_{\ell}$ -basis of  $e_H W$ . The convolution algebra  $\text{End}_{\overline{\mathbf{Q}}_{\ell}G_F} \overline{\mathbf{Z}}_{\ell}[G_F/H]$  is finitely generated [9, II.2.13], and the dimension of  $e_H W$  over  $\overline{\mathbf{Q}}_{\ell}$  is finite. Hence there exists a finite extension  $E/\mathbf{Q}_{\ell}$  such that the *E*-vector space  $\bigoplus_i Ew_i$  in  $e_H W$  is stable by the Hecke algebra  $\text{End}_{EG_F} E[G_F/H]$ . The *E*-representation *U* of  $G_F$  generated by  $(w_i)$  in *W* satisfies  $e_H U = \bigoplus_i Ew_i$ . The scalar extension  $\overline{\mathbf{Q}}_{\ell} \otimes_E U$  is equal to *W* because it is a subrepresentation of *W* with the same *H*-invariants. By the Proposition 3.1,  $W_E$  has finite length.

Let E, E' be two finite extensions. Let  $W_E \in Mod_E(G_F)^{fl}, W_{E'} \in Mod_{E'}(G_F)^{fl}$ , let

$$f: \overline{\mathbf{Q}}_{\ell} \otimes_E W_E \to \overline{\mathbf{Q}}_{\ell} \otimes_{E'} W_{E'}$$

be a  $\overline{\mathbf{Q}}_{\ell}G_{F}$ -morphism between their scalar extensions to  $\overline{\mathbf{Q}}_{\ell}$ . There exists a finite extension E'' containing E, E' such that f is defined on E'', i.e. induces a  $E''G_{F}$ -morphism  $f_{E''}: E'' \otimes_{E} W_{E} \to E'' \otimes_{E'} W_{E'}$  between their scalar extensions to E'' [9, proof of II.4.7].

The scalar extension  $s_{E'/E}$  for smooth representations of  $G_F$  respects finite length (Proposition 3.1) and the category  $Mod_{\ell}(G_F)^{fl}$  of smooth  $\ell$ -adic representations of  $G_F$  of finite length is well defined, contained in the category  $Mod_{\ell}(G_F)^{adm}$  of admissible smooth  $\ell$ -adic representations of  $G_F$  (Proposition 3.2). The category  $Mod_{\ell}(G_F)^{int,fl}$  of integral smooth  $\ell$ -adic representations of  $G_F$  of finite length is equivalent by the forgetful functor composite with the completion and the smooth part to the category  $\mathscr{B}_{\ell}(G_F)^{adm,fl}$  of Banach unitary  $\ell$ -adic representations which are admissible and of finite length.

**Theorem 3.6.** — The completion and the smooth part define equivalence of categories between  $\operatorname{Mod}_{\overline{\mathbf{Q}}_{\ell}}^{\operatorname{int}, \operatorname{fl}}(G_F)$  and  $\mathscr{B}_{\ell}(G_F)^{\operatorname{adm}, \operatorname{fl}}$ .

In particular, they give bijections between the irreducible integral smooth  $\mathbf{Q}_{\ell}$ -representations of  $G_F$  and the topologically irreducible admissible Banach unitary  $\ell$ -adic representations of  $G_F$ .

For  $G_F = GL(n, F)$ , we deduce the  $\ell$ -adic local Langlands correspondence for GL(n, F), given in the introduction.

A very natural question (asked by the referee) for a Banach unitary  $\ell$ -adic representation V of  $G_F$  (notations of the Sections 2 and 3) is: does V topologically irreducible imply V admissible? The answer is no.