

Johannes Sjöstrand

WEYL LAW FOR
SEMI-CLASSICAL
RESONANCES WITH
RANDOMLY PERTURBED
POTENTIALS

MÉMOIRES DE LA SMF 136

Société Mathématique de France 2014

Comité de rédaction

Jean BARGE	Charles FAVRE
Emmanuel BREUILLARD	Daniel HUYBRECHTS
Gérard BESSON	Yves LE JAN
Antoine CHAMBERT-LOIR	Laure SAINT-RAYMOND
Jean-François DAT	Wilhem SCHLAG
Jean-Marc DELORT	

Raphaël KRIKORIAN (dir.)

Diffusion

Maison de la SMF Case 916 - Luminy 13288 Marseille Cedex 9 France smf@smf.univ-mrs.fr	Hindustan Book Agency O-131, The Shopping Mall Arjun Marg, DLF Phase 1 Gurgaon 122002, Haryana Inde	AMS P.O. Box 6248 Providence RI 02940 USA www.ams.org
---	---	---

Tarifs

Vente au numéro : 35 € (\$52)

Abonnement Europe : 300 € hors Europe : 334 € (\$501)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Mémoires de la SMF
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris Cedex 05, France
Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96
revues@smf.ens.fr • <http://smf.emath.fr/>

© Société Mathématique de France 2014

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 1629-4939

ISBN 978-2-85629-780-3

Directeur de la publication : Marc PEIGNÉ

MÉMOIRES DE LA SMF 136

**WEYL LAW FOR
SEMI-CLASSICAL
RESONANCES WITH
RANDOMLY PERTURBED
POTENTIALS**

Johannes Sjöstrand

Société Mathématique de France 2014
Publié avec le concours du Centre National de la Recherche Scientifique

Johannes Sjöstrand

Institut de Mathématiques de Bourgogne, Université de Bourgogne, UMR
5584 du CNRS, 9 avenue Alain Savary – BP 47870, 21078 Dijon CEDEX.

Url : johannes.sjostrand@u-bourgogne.fr

2000 Mathematics Subject Classification. — 81U99, 35P20, 35P25.

Key words and phrases. — Resonance, Weyl law, Random.

Ce travail a bénéficié d'une aide de l'Agence Nationale de la Recherche portant les références JC05-52556 et ANR-08-BLAN-0228-01 ainsi que d'une bourse FABER du conseil régional de Bourgogne.

WEYL LAW FOR SEMI-CLASSICAL RESONANCES WITH RANDOMLY PERTURBED POTENTIALS

Johannes Sjöstrand

Abstract. — We consider semi-classical Schrödinger operators with potentials supported in a bounded strictly convex subset \mathcal{O} of \mathbb{R}^n with smooth boundary. Letting h denote the semi-classical parameter, we consider classes of small random perturbations and show that with probability very close to 1, the number of resonances in rectangles $[a, b] - i[0, ch^{\frac{2}{3}}[$, is equal to the number of eigenvalues in $[a, b]$ of the Dirichlet realization of the unperturbed operator in \mathcal{O} up to a small remainder.

Résumé (Loi de Weyl pour des résonances semi-classiques associées aux potentiels avec perturbations aléatoires)

On considère des opérateurs de Schrödinger dont les potentiels ont leur supports dans un ensemble strictement convexe à bord lisse $\mathcal{O} \Subset \mathbb{R}^n$. En désignant par h le paramètre semi-classique, nous considérons des classes de petites perturbations aléatoires et montrons qu'avec une probabilité très proche de 1, le nombre de résonances dans des rectangles $[a, b] - i[0, ch^{\frac{2}{3}}[$ est égal (à un petit reste près) au nombre de valeurs propres dans $[a, b]$ de la réalisation de Dirichlet de l'opérateur dans \mathcal{O} .

CONTENTS

1. Introduction	1
Acknowledgements	5
2. The result	7
3. Some elements of the proof	13
4. Grushin problems and determinants	17
4.1. Gaussian elimination	17
4.2. Generalized determinants for holomorphic Fredholm families	18
4.3. Extension to meromorphic families	21
4.4. Determinants via traces	25
4.5. Addendum	29
5. Complex dilations	31
5.1. Complex dilations and symmetry	31
5.2. Dilations and convex sets	33
6. Semi-Classical Sobolev spaces	43
7. Reductions to \mathcal{O} and to $\partial\mathcal{O}$	47
8. Some ODE preparations	53
8.1. Nullsolutions and factorizations of 2nd order ODEs	53
8.2. Simple turning point analysis	54
8.3. The exterior ODE	61
9. Parametrix for the exterior Dirichlet problem	69
10. Exterior Poisson operator and DN map	75

11. The interior DN map	81
12. Some determinants	93
13. Upper bounds on the basic determinant	97
14. Some estimates for P_{out}	111
15. Perturbation matrices and their singular values	117
16. End of the construction	121
17. End of the proof of Theorem 2.2 and proof of Proposition 2.4	129
A. WKB estimates on an interval	135
Bibliography	141

CHAPTER 1

INTRODUCTION

There is now a very large literature about the distribution of scattering poles (resonances) often using methods from non-self-adjoint spectral theory and microlocal analysis, including many results about upper and lower bounds on the density of resonances. See for instance [34], [6] and the references given there. Less is known about actual asymptotics for the number of resonances in various domains. In this paper we shall give such a result for the semi-classical Schrödinger operator

$$(1.1) \quad P = -h^2\Delta + V(x),$$

on \mathbb{R}^n where $V \in L^\infty(\mathbb{R}^n; \mathbb{R})$ has compact support.

Recall that the resonances or scattering poles of the operator (1.1) can be defined as the poles of the meromorphic extension of the resolvent

$$(P - z)^{-1} : C_0^\infty(\mathbb{R}^n) \longrightarrow H_{\text{loc}}^2(\mathbb{R}^n)$$

across the positive real axis, to the logarithmic covering space of $\mathbb{C} \setminus \{0\}$ when n is even and to the double covering when n is odd. Alternatively we can continue $(P - k^2)^{-1}$ from the upper half-plane across $\mathbb{R} \setminus \{0\}$ which gives a meromorphic function on \mathbb{C} when n is odd. Using the second definition, we can introduce the number $N(r)$ of resonances in the disc $D(0, r)$ when n is odd.

In one dimension and for $h = 1$, M. Zworski [38] showed that if $[a, b]$ is the convex hull of the support of V , then

$$(1.2) \quad N(r) = \frac{2(b-a)}{\pi}r + o(r), \quad r \rightarrow \infty,$$

which is 2 times the asymptotic number of eigenvalues $\leq r^2$ of the Dirichlet realization of $-\Delta + V$ on $[a, b]$, the factor 2 being explained by the fact that

the resonances are symmetric around the imaginary axis. He also showed that most of these concentrate to narrow sectors around the real axis. This extended an earlier result of T. Regge [20]. Subsequently, B. Simon [21] gave a different proof, inspired by the work of R. Froese [12], who got similar results for potentials that do not necessarily have compact support but are very small near infinity. See also the recent works [8], [7], [10] about Weyl and non-Weyl asymptotics for graphs.

In higher odd dimensions, M. Zworski [40] considered the case of radial potentials of the form

$$V(x) = f(|x|)$$

with support in $\overline{B(0, a)}$ where $f \in C^2([0, a])$, $a > 0$, $f(a) \neq 0$ and obtained a Weyl type asymptotics (still with $h = 1$),

$$(1.3) \quad N(r) = K_n a^n r^n + o(r^n), \quad r \rightarrow +\infty,$$

where $K_n > 0$. Recall also that Zworski [39] gave an upper bound in the non-radial case with the correct power of r and using his analysis, P. Stefanov [34], gave an explicit formula for the constant $K_n a^n$ in the radial case and showed that the right hand side of (1.3) is up to $o(r^n)$ the sum of 2 times the number of eigenvalues $\leq r^2$ for the interior Dirichlet problem in the ball $B(0, a)$ and the number of scattering poles for the exterior Dirichlet Laplacian in $\mathbb{R}^n \setminus B(0, a)$. (See also G. Vodev [35].) He also showed (as a corollary of a more general result for operators with black box) that if we drop the radially assumption and only assume that $V \in L^\infty(\mathbb{R}^n; \mathbb{R})$ has its support in $\overline{B(0, a)}$, then we have the upper bound

$$(1.4) \quad N(r) \leq K_n a^n r^n + o(r^n), \quad r \rightarrow +\infty.$$

T. Christiansen [6] introduced the set \mathfrak{M}_a of L^∞ potentials V with support in $\overline{B(0, a)}$ for which we have (1.3) and gave the leading asymptotics, of the form $C r^n$, for the number of resonances in sectors in the lower half-plane intersected with the disc $D(0, r)$. These formulas were implicit in [40], [34] in the case of the radial potentials considered there. In particular, when considering smaller and smaller sectors adjacent to \mathbb{R}_+ or \mathbb{R}_- we can see, using Lemma 3.3 of [6] and some wellknown formulas for the Γ function and the volume of the unit ball, that the constant C converges to the one we get in the leading Weyl asymptotics for the number of Dirichlet eigenvalues for the Laplacian in $B(0, a)$. In the theorems 1.2, 1.3 of the same paper the author gives interesting extensions “for most values of z ” to the case of potentials $V(x, z)$ depending holomorphically on a parameter z with $\text{supp } V(\cdot, z) \subset \overline{B(0, a)}$ such