COMPRESSIBLE NAVIER-STOKES EQUATIONS A. Novotný, R. Danchin, M. Perepelitsa Edited by D. Bresch



Panoramas et Synthèses

Numéro 50

SOCIÉTÉ MATHÉMATIQUE DE FRANCE Publié avec le concours du Centre national de la recherche scientifique

Comité de rédaction

Nicolas BERGERON Serge CANTAT Anne-Laure DALIBARD Tien-Cuong DINH Arnaud GUILLIN Marc HINDRY Pascal MASSART Ariane MÉZARD Hervé PAJOT

Bertrand RÉMY (dir.)

Diffusion

Maison de la SMF Case 916 - Luminy 13288 Marseille Cedex 9 France smf@smf.univ-mrs.fr AMS P.O. Box 6248 Providence RI 02940 USA www.ams.org EDP Sciences 17, avenue du Hoggar 91944 Les Ulis Cedex A France www.edpsciences.com

Tarifs

Vente au numéro : $40 \in (\$60)$

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Panoramas et Synthèses Société Mathématique de France Institut Henri Poincaré, 11, rue Pierre et Marie Curie 75231 Paris Cedex 05, France Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96 revues@smf.ens.fr • http://smf.emath.fr/

© Société Mathématique de France 2016

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 1272-3835

ISBN 978-2-85629-847-3

Directeur de la publication : Stéphane Seuret

PANORAMAS ET SYNTHÈSES 50

COMPRESSIBLE NAVIER-STOKES EQUATIONS

A. Novotný, R. Danchin, M. Perepelitsa

Edited by D. Bresch

Société mathématique de France Publié avec le concours du Centre national de la recherche scientifique $Didier\ Bresch$ LAMA, équipe EDPs² Bat. Le Chablais, Campus scientifique 73376 Le Bourget du Lac

Raphaël Danchin Université Paris-est, LAMA UMR 8050 and Institut Universitaire de France 61, avenue du Général de Gaulle, 94010 Créteil Cedex, France *E-mail :* danchin@univ-paris12.fr

Antonin Novotný IMATH, Université du Sud Toulon-Var, BP 132, 839 57 La Garde, France

Misha Perepelitsa University of Houston, PGH 631, 4800 Calhoun Rd., Houston, TX, USA *E-mail*: misha@math.uh.edu

2000 Mathematics Subject Classification. — 35Q30, 76N10, 35Q35.

Keywords and phrases. — Compressible Navier-Stokes equations, Navier-Stokes-Fourier system, Leray type solutions, relative entropy methods, Littlewood-Paley theory, surfaces of discontinuity for compressible flows.

TOPICS ON COMPRESSIBLE NAVIER-STOKES EQUATIONS

A. Novotný, R. Danchin & M. Perepelitsa

Abstract. — This edition of *Panoramas & Synthèses* follows the session États de la Recherche : "Topics on compressible Navier-Stokes equations" that was organized from 21 to 25 May 2012 in the Laboratoire de Mathématiques - UMR5127 CNRS, 73376 Le Bourget du Lac, France. This national training session has been the opportunity to gather four internationally well known specialists (namely R. Danchin, B. Desjardins, A. Novotny, M. Perepetlisa) allowing to present the major actual mathematical developments related to the well-posedness character problem for the compressible Navier-Stokes equations to non-subject specialists. For the sake of unity, we have decided to collect in this special issue only the contributions dedicated to the non-degenerate viscosities case, hoping by this way to present a self-contained contribution on the subject : global weak-solutions à la Leray, intermediate solutions à la Hoff and strong solutions in critical spaces à la Fujita-Kato.

Résumé. — Autour du système de Navier-Stokes compressible. Ce volume de *Panora-mas & Synthèses* fait suite à la session des États de la Recherche « Topics on compressible Navier-Stokes equations », qui a eu lieu du 21 au 25 mai 2012 au Laboratoire de Mathématiques - UMR5127 CNRS, 73376 Le Bourget du Lac, France.

Lors de cette école, quatre spécialistes de renommée internationale (R. Danchin, B. Desjardins, A. Novotny, M. Perepelitsa) présentèrent à une audience de nonspécialistes l'état de l'art et les principaux enjeux contemporains concernant le caractère bien posé des équations de Navier-Stokes compressible. Par souci d'unité, seules les contributions portant sur le cas de viscosités non dégénérées seront présentées dans ce volume. Nous espérons ainsi proposer un texte auto-contenu sur ce sujet : solutions globales à la Leray, solutions intermédiaires à la Hoff, et solutions fortes dans des espaces critiques à la Fujita-Kato.

TABLE OF CONTENTS

Did	DIER BRESCH — Topics on compressible Navier-Stokes equations : non-	:
1	Letre duction	1X :
1.	Come comments on the contributions	IX
2.	Some comments on the contributions	х
An	FONIN NOVOTNÝ — Lecture Notes on the Navier-Stokes-Fourier system :	1
1	Introduction	1
1. 9	Definitions and main results	0
2. 2	Weak compactness of the set of weak colutions	19
ე. ⊿	Dreaf of Theorem 2.2. Deleting entropy inequality	10
4. E	Weak strong uniqueness	20
э.	Defenences	31 40
	References	40
RAI	RAPHAËL DANCHIN — Fourier analysis methods for compressible flows	
	Introduction	43
1.	The Fourier analysis toolbox	45
2.	Solving the compressible Navier-Stokes equations in critical spaces	62
3.	Partially parabolic or dissipative linear PDEs	80
4.	Global results	92
	Appendix	101
	References	103
Mis	SHA PEREPELITSA — Near-equilibrium weak solutions of the Navier-Stokes	
	equations for compressible flows	107
1.	Properties of the solutions at the interface of discontinuity	107
2.	Existence results	112
3.	Proof of Theorem 1	114
4.	Dynamics of surfaces of discontinuity	120
5.	Dynamics of surfaces of discontinuity with corner singularities	125
	Appendix	130
	References	134

ABSTRACTS

Lecture Notes on the Navier-Stokes-Fourier system: weak solutions, relative entropy inequality, weak strong uniqueness

 Αντονίν Νονοτνή
 1

We investigate three issues in the theory of the Navier-Stokes-Fourier system describing the motion of a compressible, viscous and heat conducting fluid: weak compactness of the family of weak solutions (that is the main ingredient in the proof of the existence of weak solutions), relative entropy inequality and weakstrong uniqueness principle.

Fourier analysis methods for compressible flows RAPHAËL DANCHIN 43

Since the 80's, Fourier analysis methods have known a growing importance in the study of linear and nonlinear PDE's. In particular, after the seminal paper by J.-M. Bony [Ann. Sci. École Norm. Sup. 14 (1981)], the use of Littlewood-Paley decomposition and paradifferential calculus brought a number of new results whenever those equations are considered in the whole space or the torus.

We here aim at giving a survey of recent advances obtained by Fourier analysis methods in the context of compressible fluid mechanics. For simplicity, we shall focus on the *barotropic* compressible Navier-Stokes equations.

 Near-equilibrium weak solutions of the Navier-Stokes equations for compressible flows

 MISHA PEREPELITSA
 107

In these notes we review the theory of weak solutions of the Navier-Stokes equations for compressible flows in near equilibrium flow regime. The study of solutions of this type was initiated by D. Hoff [*Arch. Rational Mech. Anal.* **114** (1991)] with the motivation of studying the dynamics of an interface of discontinuity of solutions.

We demonstrate three results. The first is the existence of weak solutions with uniformly bounded density for the typical initial-boundary value problems. The second result improves the regularity of such weak solutions in the case when the density is piecewise Hölder continuous with a jump discontinuity across a $C^{1+\alpha}$ hypersurface. We show that the flow generated by the velocity u transports the discontinuity surface and preserve its regularity.

In the last result, we consider the fluid flows in which the density, at time t = 0, is piecewise smooth and the interface of discontinuity is a surface with a cornertype singularity. We show that solutions in Hoff's regularity class are well suited for the analysis of this problem and describe the phenomenon of an instantaneous change of the geometry of the interface at the point of the singularity.

TOPICS ON COMPRESSIBLE NAVIER-STOKES EQUATIONS: NON-DEGENERATE VISCOSITIES

Didier Bresch

Dedicated to the memory of Alexander Kazhikov.

1. Introduction

Professor A.V. Kazhikhov was one of the very distinguished analysts of our time. He made fundamental contributions on the mathematical theory of fluid mechanics, in particular he achieved outstanding results on the compressible Navier-Stokes system that, had, and still have, a significant influence on this field.

This special issue of Panoramas et Synthèses (SMF), dedicated to the memory of the Professor A.V. Kazhikhov, consists of three mathematical contributions dealing with compressible flows PDEs. They correspond to mini-courses respectively given by R. Danchin, A. Novtony, M. Perepetlisa with a constant or non-degenerate viscosities assumption. A contribution with degenerate viscosities (depending on the density) was presented by B. Desjardins during the session États de la Recherche but is not included in the volume because it does not really belongs to the same framework.

The main objective of this issue is to help the reader understand various mathematical tools that have been developed recently to deal with the problem of well posedness for the compressible Navier-Stokes equations with constant or at least nondegenerate viscosities. The three contributions focus on different kind of solutions, namely:

- Global weak solutions (regularity given by the energy estimates),
- Intermediate but not regular (discontinuous density and velocity with (low) regularity),
- Scaling invariant strong solutions (critical regularity).

In order to prepare the reader to these three main contributions, let us present in the following section different tools and main ingredients used by the authors. We hope by this section to motivate and help the reader start reading the contributions by showing them nice properties contained in the compressible Navier-Stokes equations. We start with the contribution by A. Novotny because it corresponds to the weakest regularity. We then continue with the contribution by M. Perepetlisa which concerns

intermediate regularity, namely discontinuous density. Then we end the issue with the contribution by R. Danchin which focuses on critical spaces regularity. For each contributions, we will present some properties which are included in some sense in the contributions and mathematically justified in a more general setting sometimes.

Interested readers are invited to read the extensive exposition which can be found in the monographs by P.-L. Lions, E. Feireisl, A. Novotny and I. Straskraba, S. Benzoni and D. Serre, H.Bahouri, J.-Y. Chemin and R. Danchin and papers by D. Hoff *et al.*

2. Some comments on the contributions

The contribution by A. Novotny. – It concerns the existence of global weak solutions for the compressible Navier-Stokes-Fourier system in the spirit of the result obtained in 1933 by J. Leray for the incompressible Navier-Stokes equations, the existence of suitable weak solutions using the concept of relative entropies, the weak strong uniqueness property in the class of weak solutions. The main ingredients are energy estimates, weak limit, compactness of the temperature, div-curl Lemma, parametrized Young measures, effective flux, commutators, oscillations defect measure and renormalized continuity equation, relative entropy type inequality, weak-strong uniqueness techniques.

Global existence of weak solutions à la Leray. – Concerning the global in time existence result for the heat conducting Navier-Stokes equations, viscosities and conductivity coefficients are assumed to depend on the temperature (not on the density) and to be non-degenerate. In the barotropic case, namely in the non-dependent temperature case, viscosities are assumed to be constant and isotropic (same viscosities in all the directions: no eddy viscosities): see the book by P.-L. Lions and the one by A. Novotny and I. Strakraba. Extending the result to constant eddy viscosities is an interesting open problem with applications in geophysics.

For the heat conducting Navier-Stokes equations, dependency with respect to the temperature is possible because of the parabolic behavior of the equation satisfied by the temperature. Isotropy of the viscosities is necessary to get the weak compacted property on the effective flux $F = (2\mu + \lambda) \operatorname{div} u - p$. The regularity on the temperature helps to control commutators.

The form of the pressure is also strongly used to derive estimates on the density and find weak limit properties: the pressure is composed of two parts (a cold part which does not depend on the temperature but depends on the density and a part which depends on the density and on the temperature). It remains then to characterize weak limit of some difficult nonlinear terms: mainly the weak limit of the pressure term, viscous term. For that the authors show strong convergence of temperature. This point involves the treatment of the entropy production rate as a Radon measure and a convenient use of the compensated compactness lemma in combination with the theory of parametrized Young measures. Then it is needed to get strong convergence of densities. As for the barotropic case, weak compactness on the effective viscous flux is the key point and established using Riesz transform, commutators estimates à la Meyer through div-curl Lemma, boundedness of the oscillation defect measure for the sequence of densities, renormalized solution to the continuity equation and show the oscillations in the density sequence do not increase in time.

Some heuristic calculations. – For the reader's convenience, let us explain heuristically the main steps to prove global existence of weak solutions in the barotropic case with constant viscosities and power γ pressure law. This will help in order to understand the Navier-Stokes-Fourier analysis in the contribution by A. Novtony. In that case, the compressible Navier-Stokes equation with constant viscosities reads

(1)
$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + \nabla p(\rho) = \rho f, \end{cases}$$

with $p(\rho) = a\rho^{\gamma}$. For simplicity, we assume to be in a Lipschitz bounded domain Ω with homogeneous Dirichlet boundary conditions on the velocity. When ρ and u are regular and satisfied the continuity equation, for all $b \in \mathcal{C}([0, +\infty))$, it is clear, multiplying the mas equation by $b'(\rho)$, that (ρ, u) satisfy also the renormalized continuity equation (this terminology coming from the transport theory by R.J. DiPerna and P.-L. Lions). This equation reads

$$\partial_t b(\rho) + \operatorname{div}(b(\rho)u) + (b'(\rho)\rho - b(\rho))\operatorname{div} u = 0.$$

For $T \in (0, +\infty)$, f, ρ_0 , m_0 satisfying some technical assumptions, we say that a couple (ρ, u) is a weak renormalized solution with bounded energy if it possesses the following properties:

- $\rho \in L^{\infty}(0,T;L^{\gamma}(\Omega)) \cap \mathscr{C}^{0}([0,T],L^{\gamma}_{\text{weak}}(\Omega)), \ \rho \geq 0 \ a.e. \text{ in } (0,T) \times \Omega, \ \rho|_{t=0} = \rho_{0}$ a.e. in Ω ,
- $\begin{array}{l} \ u \in L^{2}(0,T;H^{1}_{0}(\Omega)); \ \rho |u|^{2} \in L^{\infty}(0,T;L^{1}(\Omega)) \ \text{and} \ \rho u \in \ \mathcal{C}^{0}([0,T];L^{2\gamma/(\gamma+1)}_{\text{weak}}(\Omega)), \\ (\rho u)|_{t=0} = m_{0} \ a.e. \ \Omega; \end{array}$
- the solution extended by zero is solution of the mass and momentum equations in $\mathscr{D}'((0,T) \times \mathbb{R}^d)$;
- for almost all $\tau \in (0,T)$, (ρ, u) satisfies the energy inequality

$$E(\rho, u)(\tau) + \int_0^\tau \int_\Omega (\mu |\nabla u|^2 + (\lambda + \mu) |\operatorname{div} u|^2) \le E_0 + \int_0^t \int_\Omega \rho f \cdot u$$

In this inequality, $E(\rho, u)(\tau) = \int_{\Omega} (\rho |u|^2/2 + a\rho^{\gamma}/(\gamma - 1))(\tau)$ denotes the total energy at time τ and $E_0 = \int_{\Omega} |m_0|^2/2\rho_0 + a\rho_0^{\gamma}/(\gamma - 1))$ denotes the initial total energy;

 $-b(\rho)$ satisfies the renormalized equation for b with some increasing properties.

The theory developed by P.-L. Lions to prove the global existence of renormalized weak solution with bounded energy asks for some limitation on the adiabatic constant γ , namely $\gamma > 3d/(d+2)$. Note that E. Feireisl *et al.* have generalized this approach in order to cover the range $\gamma > 3/2$ in dimension 3 and more generally $\gamma > d/2$ where *d* is the space dimension. We will try here to help the readers to understand

the different lines of the proof in the "simplest" case, namely the one studied by P.-L. Lions. As usual in PDEs, the method asks for the compactness of a bounded sequence of solutions satisfying uniformly the energy estimates. The construction of such an approximate sequence is not the subject of this heuristic part (see the book by A. Novotny and I. Straskraba for details).

We present the proof due to P.-L. Lions and indicate quickly at the end how it was improved by E. Feireisl *et al.*. Let us note that using some appropriate multiplicator, for $\gamma > d/2$, it is possible to prove the following key estimate on the density

$$\int_0^\infty dt \int_\Omega \rho^{\gamma+\theta} \leq C(R,T) \text{ for } \theta = \frac{2}{d}\gamma - 1.$$

This observation is due to P.-L. Lions. To prove such an estimate, we need the fact that (ρ_n, u_n) is a sequence solution of the renormalized equation in which $b(s) = s^{\theta}$. Let us observe also that the regularity asked for the domain is linked to the use of the Bogovski operator as proposed by E. Feireisl *et al.* Using the energy estimate and the extra integrability property proved on the density, it is is then possible to precise some convergence and to show that

$$\begin{split} \rho_n &\to \rho \text{ in } \mathcal{C}^0([0,T]; L^{\gamma}_{\text{weak}}(\Omega)) \\ \rho_n^{\gamma} &\to \overline{\rho^{\gamma}} \text{ in } L^{(\gamma+\theta)/\gamma}((0,T) \times \Omega) \\ \rho_n u_n &\to \rho u \text{ in } \mathcal{C}^0([0,T]; L^{2\gamma/(\gamma+1)}(\Omega)) \\ \rho u_n^i u_n^j &\to \rho u u^j \text{ in } \mathcal{D}'((0,T) \times \Omega) \text{ for } i, j = 1, 2, 3. \end{split}$$

Let us note the convergence in some nonlinear terms which comes from the strong convergence of ρ_n deduced from the uniform estimate on $\partial_t \rho_n$ given by the mass equation and of the strong convergence of $\sqrt{\rho_n u_n}$ deduced form the uniform estimate on $\partial_t (\rho_n u_n)$ given by the momentum equation (which provides strong convergence in negative sobolev spaces) and by the convergence of the term $\int_0^T \int_\Omega \rho_n |u_n|^2$ writting $\int_0^T \int_\Omega \rho_n |u_n|^2 = \int_0^T \langle \rho_n u_n, u_n \rangle_{H^{-1}(\Omega) \times H^1_0(\Omega)}$. Consequently, the prolongation by zero in $(0, T) \times R^3 / \Omega$ of the functions $\rho, u, \overline{\rho^{\gamma}}$ denoted again $\rho, u, \overline{\rho^{\gamma}}$ satisfy the equations

(2)
$$\begin{cases} \partial_t \rho + \operatorname{div}(\rho u) = 0, \\ \partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) - \mu \Delta u - (\lambda + \mu) \nabla \operatorname{div} u + a \nabla \overline{\rho^{\gamma}} = \rho f. \end{cases}$$

The difficulty consists in proving that (ρ, u) is a renormalized weak solution with bounded energy and is mainly linked to the proof that $\overline{\rho^{\gamma}} = \rho^{\gamma}$ a.e. in $((0,T) \times \Omega$. This asks for a kind of compactness on the density sequence which is not available from the estimates only. This is an observation by P.-L. Lions which will fill the gap: The sequence $\{a\rho_n^{\gamma} - \lambda + 2\mu \operatorname{div} u_n\}_{n \in N^*}$, usually called viscous effective flux possesses a kind of weak compactness. This property was previously identified in one space dimension by D. Hoff and D. Serre. More precisely, we have the following property

xiii

for all function $b \in \mathcal{C}^1([0, +\infty))$ satisfying some increasing properties at infinity

$$\lim_{n \to +\infty} \int_0^T \int_\Omega (a\rho_n^{\gamma} - (2\mu + \lambda) \mathrm{div} u_n) b(\rho_n) \varphi dx dt = \int_0^T \int_\Omega (a\overline{\rho^{\gamma}} - (2\mu + \lambda) \mathrm{div} u) \overline{b(\rho)} \varphi dx dt$$

where the overline quantities design the weak limit of the corresponding quantities and $\varphi \in \mathcal{D}((0,T) \times \Omega)$. In higher dimension, the proof by P.-L. Lions is based on harmonic analysis due to R. Coifman and Y. Meyer and takes into account the observations by D. Serre made in the one-dimensional case. The proof by E. Feireisl is based on divcurl Lemma introduced by F. Murat and L Tartar. Let us explain how P.-L. Lions concluded about the strong convergence on the density sequences using this weak compactness property on the effective flux. To simplify the computations, we will assume $\gamma \geq 3d/(d+2)$ and in that case due to the extra integrability on the density, we get that $\rho_n \in L^2((0,T) \times \Omega)$. In that case, the transport theory by R.J. DiPerna and P.-L. Lions applies to the continuity equation in order to get the validity of the renormalized equations. Then choose $b(s) = s \ln s$ in the renormalized formulation for ρ_n and ρ and do the difference of the two equations. Then pass to the limit $n \to +\infty$ and use the identity of weak compactness on the effective flux to replace terms with divergence of velocity by terms with density. This allows to get an evolution equation on the amplitude of the oscillation for the sequence of density. It reads

$$\partial_t (\overline{\rho \ln \rho} - \rho \ln \rho) + \operatorname{div}((\overline{\rho \ln \rho} - \rho \ln \rho)u) = \frac{a}{2\mu + \lambda} (\overline{\rho^{\gamma}} \rho - \overline{\rho^{\gamma+1}}).$$

The formal integration of this equation on $(0, T) \times \Omega$, the monotonicity of the pressure $p(\rho) = a\rho^{\gamma}$ and the strict-convexity of the function $s \mapsto s \ln s$, $s \ge 0$ implies that $\rho \ln \rho = \rho \ln \rho$ if initially it is the case. In other terms, the weak convergence commute with a strictly convex function and therefore it gives the strong convergence of the densities. This achieves the proof of the Theorem in the case $\gamma \ge 3d/(d+2)$. The proof of E. Feireisl works even if the density is not a priori square integrable. For that E. Feireisl observes that it is possible to control the amplitude of the possible oscillations on the density in a norm L^p with p > 2 allowing to use an effective flux property with some troncature. The existence result is then also valid for $\gamma > d/2$: See the book by A. Novotny and I. Straskraba. For the heat conducting Navier-Stokes equations, the tools are more complicated but the steps endowed in the barotropic case are included in: Estimates and weak limits (step 1, step 4, step 5, step 6), Strong convergence of densities (Section 3) of course with not straightforward adaptations.

Remark. – If we have oscillation-concentration on the initial density sequence, the result is more complex because $\overline{\rho_0 \ln \rho_0} \neq \rho_0 \ln \rho_0$. We refer the reader to the recent paper by D. Bresch and M. Hillairet (2013) where they justify the derivation of viscous and compressible multi-component flow equations if initially the initial density sequence is uniformly bounded with corresponding Young measures which are linear convex combination of m Dirac measures.

Relative entropy and suitable weak solutions. – Since the pioneering work of C. Dafermos, relative entropy methods have become a crucial and widely used tool in the