

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

TYPE THEORY AND COEFFICIENT SYSTEMS ON THE BUILDING

Paul Broussous & Peter Schneider

**Tome 145
Fascicule 1**

2017

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 97-159

Le *Bulletin de la Société Mathématique de France* est un périodique
trimestriel de la Société Mathématique de France.

Fascicule 1, tome 145, mars 2017

Comité de rédaction

Christine BACHOC	Laurent MANIVEL
Emmanuel BREUILLARD	Julien MARCHÉ
Yann BUGEAUD	Kieran O'GRADY
Jean-François DAT	Emmanuel RUSS
Charles FAVRE	Christophe SABOT
Marc HERZLICH	Wilhelm SCHLAG
Raphaël KRIKORIAN	

Pascal HUBERT (Dir.)

Diffusion

Maison de la SMF - Case 916 - Luminy - 13288 Marseille Cedex 9 - France
christian.smf@cirm-math.fr

Hindustan Book Agency	AMS
O-131, The Shopping Mall	P.O. Box 6248
Arjun Marg, DLF Phase 1	Providence RI 02940
Gurgaon 122002, Haryana	USA
Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

bullsmf@ihp.fr • smf.emath.fr

© *Société Mathématique de France* 2017

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Stéphane SEURET

TYPE THEORY AND COEFFICIENT SYSTEMS ON THE BUILDING

BY PAUL BROUSSOUS & PETER SCHNEIDER

ABSTRACT. — Let F be a non-archimedean local field and G be the group $GL(N, F)$ for some integer $N \geq 2$. Let π be a smooth complex representation of G lying in the Bernstein block $\mathcal{B}(\pi)$ of some simple type in the sense of Bushnell and Kutzko [10]. Refining the approach of the second author and U. Stuhler in [18], we canonically attach to π a subset X_π of the Bruhat-Tits building X of G , as well as a G -equivariant coefficient system $\mathcal{C}[\pi]$ on X_π . Roughly speaking the coefficient system is obtained by taking isotypic components of π according to some representations constructed from the Bushnell and Kutzko type of π . We conjecture that when π has central character, the augmented chain complex associate to $\mathcal{C}[\pi]$ is a projective resolution of π in the category $\mathcal{B}(\pi)$. Moreover we reduce this conjecture to a technical lemma of representation theoretic nature. We prove this lemma when π is an irreducible discrete series of G . Following closely [19], we then attach to any irreducible discrete series π of G an explicit pseudo-coefficient f_π and obtain a Lefschetz type formula for the value of the Harish-Chandra character of π at a regular elliptic element. In contrast to that of [19], this formula allows explicit character value computations.

Texte reçu le 15 février 2015, modifié le 7 avril 2016, accepté le 7 avril 2016.

PAUL BROUSSOUS, Laboratoire de Mathématiques, Téléport 2 - BP 30179,
Bd Marie et Pierre Curie, 86962 Futuroscope Chasseneuil Cedex, France •
E-mail : Paul.Broussous@math.univ-poitiers.fr

PETER SCHNEIDER, Universität Münster, Mathematisches Institut, Einsteinstr. 62, 48291
Münster, Germany • *E-mail* : pschnei@uni-muenster.de

RÉSUMÉ (*Théorie des types et système de coefficients sur l'immeuble*). — Soient F un corps local non archimédien et G le groupe $\mathrm{GL}(N, F)$, pour un entier $N \geq 2$. Soit π une représentation lisse complexe de G appartenant au bloc de Bernstein $\mathcal{B}(\pi)$ d'un type simple au sens de Bushnell et Kutzko [10]. En affinant l'approche que proposent le second auteur et U. Stuhler dans [18], nous attachons canoniquement à π un sous-ensemble X_π de l'immeuble de Bruhat-Tits X de G , ainsi qu'un système de coefficients G -équivariant $\mathcal{C}[\pi]$ sur X_π . Grossièrement parlant, le système de coefficients est construit en prenant des composantes isotypiques de π selon des représentations construites à partir du type de Bushnell et Kutzko de π . Nous conjecturons que lorsque π possède un caractère central, le complexe de chaînes augmenté associé à $\mathcal{C}[\pi]$ est une résolution de π dans la catégorie $\mathcal{B}(\pi)$. De plus nous réduisons cette conjecture à un lemme technique en théorie des représentations. Nous démontrons ce lemme lorsque π est une représentation irréductible de la série discrète de G . Ensuite, suivant de près [19], nous attachons à toute représentation irréductible π de la série discrète de G un pseudo-coefficient explicite f_π et obtenons une formule de type Lefschetz pour la valeur du caractère de Harish-Chandra de π en un élément elliptique régulier. Contrairement à celle obtenue dans [19], notre formule permet des calculs explicites.

Introduction

Let F be a non-archimedean local field, and for some integer $N \geq 2$, let G denote the locally compact group $\mathrm{GL}(N, F)$ and X its Bruhat-Tits building. The aim of this work is to refine the construction of [18] (also see [19]) to attach to certain representations of G new equivariant coefficient systems on the Bruhat-Tits building. These representations belong to the Bernstein blocks of the category of smooth complex representations of G corresponding to *simple types* in the sense of Bushnell and Kutzko [10]. Let (π, \mathcal{V}) be a smooth complex representation of G . In [18] an equivariant coefficient system $\mathcal{C}[\pi]$ is constructed by attaching to each simplex σ of X the space of vectors fixed by a certain congruence subgroup of level e of the parahoric subgroup of G fixing σ . Here the integer e is such that \mathcal{V} is generated as a G -module by its vectors fixed by the principal congruence subgroup of level e of some maximal compact subgroup of G . In [18] it is proved that the augmented chain complex $C_\bullet(X, \mathcal{C}[\pi]) \rightarrow \mathcal{V}$ of X with coefficients in $\mathcal{C}[\pi]$ is exact. If one moreover assumes that (π, \mathcal{V}) admits a central character χ , then $C_\bullet(X, \mathcal{C}[\pi]) \rightarrow \mathcal{V}$ is a projective resolution of (π, \mathcal{V}) in the category of smooth representations of G with central character χ . In [3], the first author gave another proof of this fact for Iwahori-spherical representations. In [19], the second author and U. Stuhler draw some important consequences concerning the harmonic analysis on G as well as the homological algebra of the category of smooth representations of G . Among other things they prove that these projective resolutions give rise to pseudo-coefficients for discrete series representations (generalizing the pseudo-coefficient constructed by Kottwitz in [15] for the Steinberg representation) as

well as a Lefschetz type character formula for the Harish-Chandra character of any smooth representation. Note that if the construction of [18] is restricted to the group G , [19] gives a generalization to any connected reductive F -group \mathbb{G} and most of its results are valid without restriction on G (but sometimes F is assumed to have characteristic 0, and $\mathbb{G}(F)$ to have compact center).

If the construction and results of [18], [19] have important theoretic consequences, they do not allow explicit calculations. Indeed in general the coefficient system $\mathcal{C}(\pi)$ cannot be explicitly computed (except may be in the *level 0 case*, but this is nowhere written). Indeed the only explicit way to be given an irreducible smooth representation of G is to specify its Bushnell and Kutzko type. This is why it is natural to seek for a refinement of [18] based on Bushnell and Kutzko theory.

In this paper, for technical reasons, we restrict to representations belonging to Bernstein blocks of G attached to simple types. These Bernstein blocks are exactly those containing discrete series representations. We fix a simple type (J, λ) and denote by $\mathcal{R}_{(J, \lambda)}(G)$ the category of smooth representations of G that are generated by their λ -isotypic component. We fix a smooth representation (π, \mathcal{V}) of G lying in $\mathcal{R}_{(J, \lambda)}(G)$. To the datum (J, λ) , in a non canonical way, one may associate a field extension E/F of degree dividing N whose multiplicative group E^\times is embedded in G . The centralizer G_E of E^\times in G is isomorphic to $\mathrm{GL}(N/[E : F], E)$. Using a result of the first author and B. Lemaire [5], we may view the Bruhat-Tits building X_E of G_E as being embedded in X in a G_E -equivariant way. We show that *hidden* in the properties of *Heisenberg representations* constructed in [10]§ (5.1) and in the *mobility* of simple characters established in *loc. cit.* § (3.6), there is a *geometric structure* allowing to attach to π a G_E -equivariant coefficient system $\mathcal{C}_E[\pi]$ on the first barycentric subdivision $\mathrm{sd}(X_E)$ of X_E . More precisely, in a non canonical way, we attach to (J, λ) a collection of pairs $(J^1(\sigma, \tau), \eta(\sigma, \tau))_{\sigma \subset \tau}$, where σ and τ run over the simplices of X_E satisfying $\sigma \subset \tau$. Here $J^1(\sigma, \tau)$ is some compact open subgroup of G and $\eta(\sigma, \tau)$ a Heisenberg representation of $J^1(\sigma, \tau)$ as considered in *loc. cit.* (5.1.14) (but Bushnell and Kutzko do not use this language nor this notation). Moreover the collection $(J^1(\sigma, \tau), \eta(\sigma, \tau))_{\sigma \subset \tau}$ is G_E -equivariant. Exploiting the compatibility relations among the various $\eta(\sigma, \tau)$ proved in *loc. cit.* § (5.1), and by taking isotypic components of \mathcal{V} according to the Heisenberg representations $\eta(\sigma, \tau)$, we construct our equivariant coefficient system $\mathcal{C}_E[\pi]$.

We then show that the subset $X[E]$ of X obtained by taking the union of the $g.X_E$, where g runs over G , has the structure of a G -simplicial complex containing X_E as a subcomplex. We naturally attach to $\mathcal{C}_E[\pi]$ a G -equivariant coefficient system $\mathcal{C}[\pi]$ on the first barycentric subdivision of $X[E]$ and show that it actually derives from a coefficient complex on $X[E]$, still denoted by $\mathcal{C}[\pi]$. We prove that the simplicial complex $X[E]$ and the coefficient system $\mathcal{C}[\pi]$ are actually independent of any choice made in their construction: these are