

## BANACH $\ell$ -ADIC REPRESENTATIONS OF $p$ -ADIC GROUPS

by

Marie-France Vignéras

---

**Abstract.** — Let  $p \neq \ell$  be two distinct prime numbers, let  $F$  be a  $p$ -adic field and let  $E$  be an  $\ell$ -adic field. We prove that the smooth part and the completion are inverse equivalences of categories between the category of admissible Banach unitary  $E$ -representations of  $GL(n, F)$  and the category of admissible smooth  $E$ -representations of  $GL(n, F)$  equipped with a commensurability class of lattices. We formulate the  $\ell$ -adic local Langlands correspondence as a canonical bijection between the  $n$ -dimensional  $\ell$ -adic representations of the absolute Galois group  $\text{Gal}_F$  and the topologically irreducible admissible Banach unitary  $\ell$ -adic representations of  $GL(n, F)$ .

**Résumé (Représentations  $\ell$ -adiques de groupes  $p$ -adiques).** — Soient  $p \neq \ell$  deux nombres premiers distincts, soit  $F$  un corps  $p$ -adique et soit  $E$  un corps  $\ell$ -adique. Nous démontrons que la partie lisse et la complétion définissent des équivalences de catégories inverses l'une de l'autre entre la catégorie des représentations admissibles de Banach unitaires de  $GL(n, F)$  sur  $E$  et la catégorie des représentations lisses admissibles de  $GL(n, F)$  sur  $E$  munies d'une classe de commensurabilité de réseaux. Nous formulons la correspondance de Langlands locale  $\ell$ -adique comme une bijection canonique entre les représentations  $\ell$ -adiques de dimension  $n$  du groupe de Galois absolu  $\text{Gal}_F$  et les représentations topologiquement irréductibles admissibles de Banach unitaires  $\ell$ -adiques de  $GL(n, F)$ .

### 1. Introduction

Let  $p$  be a prime number, let  $F$  be a finite extension of  $\mathbf{Q}_p$  or a field of Laurent series  $k((T))$  over a finite field  $k$  of characteristic  $p$ , let  $\bar{F}$  be an algebraic closure of  $F$  and let  $n$  be an integer  $\geq 1$ .

For any topological field  $C$ , the continuous representations of  $GL(n, F)$  on topological vector spaces over  $C$  are interesting for their applications in arithmetic, geometry or physics, via the theory of  $L$ -functions associated to automorphic representations. When  $C$  varies, the theories of  $C$ -representations of  $GL(n, F)$  present simultaneously

---

**2010 Mathematics Subject Classification.** — 11F70, 11F80, 11F85.

**Key words and phrases.** — Banach  $\ell$ -adic representation, reductive  $p$ -adic group, admissible representation, smooth representation, lattice.

strong similarities and strong different features but the Langlands insight, when  $C$  is the complex field, to use the smooth complex representations of  $\text{Gal}_F = \text{Gal}(\overline{F}/F)$  as a classifying scheme, seems to extend to other fields.

Why moving the coefficient field  $C$ ? There are many reasons.

1) The representations of  $\text{Gal}_F$  appearing naturally are not smooth complex. In the étale cohomology of proper smooth algebraic varieties, they are continuous  $\ell$ -adic representations on finite dimensional vector spaces  $V$  over finite extensions  $E/\mathbf{Q}_\ell$ , for a prime number  $\ell$ . By a reduction of a stable  $O_E$ -lattice of  $V$ , they give smooth mod  $\ell$ -representations over the residual field of  $E$ .

2) The local Langlands correspondence for  $GL(n, F)$ , over any algebraically closed field  $R$  of characteristic different from  $p$ , is a bijection

$$\pi \leftrightarrow (\rho, N)$$

between the equivalence classes of the smooth irreducible  $R$ -representations  $\pi$  of  $GL(n, F)$  and of the pairs  $(\rho, N)$  where  $\rho$  is a  $n$ -dimensional smooth semi-simple  $R$ -representation of the Weil group  $W_F$  and  $N$  a nilpotent endomorphism of the space of  $\rho$  such that  $\rho(w)N = N|w|\rho(w)$  where  $|?|$  is the unramified  $R$ -character of  $W_F$  sending a geometric Frobenius to  $q$ , the order of the residual field of  $F$ .

Our purpose is to obtain a local Langlands correspondence for continuous  $\ell$ -adic representations.

**Theorem 1.** — *Let  $\ell$  be a prime number different from  $p$ . The  $\ell$ -adic local Langlands correspondence for  $GL(n, F)$  is a canonical bijection between the equivalence classes of*

- a)  *$n$ -dimensional continuous  $\ell$ -adic representations of  $\text{Gal}_F$  with a semi-simple action of the Frobenius,*
- b) *topologically irreducible admissible Banach unitary  $\ell$ -adic representations of  $GL(n, F)$ .*

This theorem<sup>(1)</sup> is motivated by the fascinating work and conjectures of Christophe Breuil on the  $p$ -adic local Langlands correspondence, where topologically irreducible admissible Banach unitary  $p$ -adic representations of  $GL(2, \mathbf{Q}_p)$  appear naturally.

With the existing literature, one translates the local Langlands complex correspondence for  $GL(n, F)$  into a canonical bijection between the isomorphism classes of a) and of

- c) *Irreducible smooth  $\overline{\mathbf{Q}}_\ell$ -representations of  $GL(n, F)$  with a stable lattice.*

Indeed, as is well known,

(i) The smooth complex local Langlands correspondence  $LL(\rho, N)$  twisted by a suitable unramified character,

$$(\rho, N) \leftrightarrow LL(\rho, N) \otimes |\det?|^{-(n-1)/2},$$

---

<sup>(1)</sup> Proved in a letter to Breuil in september 2003, and announced in the Emmy Noether lectures 2005 of Goettingen.

called the smooth complex local Hecke correspondence, is  $\text{Aut } \mathbf{C}$ -equivariant [H prop.6].

(ii) Transporting the correspondence (i) with an algebraic isomorphism  $j : \mathbf{C} \simeq \overline{\mathbf{Q}}_\ell$ , we obtain the smooth local Hecke  $\overline{\mathbf{Q}}_\ell$ -correspondence, which does not depend on the choice of the isomorphism  $j$ .

(iii)  $N$  disappears when one considers continuous  $\overline{\mathbf{Q}}_\ell$ -representations of  $W_F$  instead of smooth  $\overline{\mathbf{Q}}_\ell$ -representations. The pairs  $(\rho, N)$  are in bijection

$$(\rho, N) \leftrightarrow \sigma$$

with the  $n$ -dimensional  $\ell$ -adic representations  $\sigma$  of  $W_F$  with a semi-simple action of the Frobenius. The reason is that the kernel of the natural morphism  $t : I_F \rightarrow \mathbf{Z}_\ell$  is a profinite group prime to  $\ell$ . There is a nilpotent endomorphism  $N$  of the space of  $\sigma$  such that  $\sigma(?) = \exp(t(?)N)$  on a subgroup of finite index of  $I_F$  [8].

(iv) The  $n$ -dimensional  $\ell$ -adic representation  $\sigma$  of  $W_F$  in (iii) extends by continuity to an  $\ell$ -adic representation of  $\text{Gal}_F$  if and only if  $\rho$  has a bounded image (i.e. the values of determinants of the irreducible components of  $\rho$  are units) [8].

(v)  $\rho$  has a bounded image if and only if  $\pi = LL(\rho, N)$  is integral [10, §1.4]; moreover all stable lattices in  $\pi$  are commensurable [11, Theorem 1].

Our task is to show that the completion with respect to a stable lattice gives a bijection between the isomorphism classes of b) and of c).

The beginning of the proof is valid for any locally profinite group  $G$ , with a countable fundamental system of neighborhoods of the unit, consisting of open profinite groups of pro-order not divisible by  $\ell$  (Section 2). We prove (Theorem 2.12) that the completion and the smooth part induce equivalences of categories between the category  $\mathcal{M}_\ell(G)^{\text{adm}}$  of admissible smooth  $\ell$ -adic representations of  $G$  equipped with a commensurability class of lattices, and the category  $\mathcal{B}_\ell(G)^{\text{adm}}$  of admissible Banach unitary  $\ell$ -adic representations of  $G$ .

Then we consider the group of rational points  $G_F$  of any reductive connected group over a local non Archimedean field  $F$  of residual characteristic  $p \neq \ell$  (Section 3). We prove (Theorem 3.6) that the completion and the smooth part induce equivalences of categories between the category  $\text{Mod}_{\overline{\mathbf{Q}}_\ell}^{\text{int, fl}}(G_F)$  of integral smooth  $\overline{\mathbf{Q}}_\ell$ -representations of  $G_F$  of finite length and the category  $\mathcal{B}_\ell(G_F)^{\text{adm, fl}}$  of admissible Banach unitary  $\ell$ -adic representations of topological finite length of  $G_F$ . We deduce the wanted bijection between the isomorphism classes of b) and c) by restricting to irreducible representations and choosing  $G_F = GL(n, F)$ .

A natural question was raised by the referee: Is a topologically irreducible Banach unitary  $\ell$ -adic representation of  $G_F$  always admissible? L. Clozel noticed that the examples of B. Diarra [5, th. 4] (van Rooj), give examples of topologically irreducible representations  $V \in \mathcal{B}_E(GL(1, F))$  where any non zero intertwining operator is bijective, which are *not admissible*.

## 2.

**2.1. The two categories.** — Let  $\ell \neq p$  be two distinct prime numbers, let  $E/\mathbf{Q}_\ell$  be a finite extension of ring of integers  $O_E$ , of uniformizer  $p_E$ , and of residual field  $k_E$ , and let  $G$  be a topological group admitting a *countable* fundamental system of neighborhoods of the unit consisting of open *pro- $\ell'$* -subgroups (profinite subgroups of pro-order *prime to  $\ell$* ).

After having recalled some definitions and properties concerning the representations of the group  $G$  on  $E$ -vector spaces, we will introduce the two categories of representations  $\mathcal{M}_E(G)$  and  $\mathcal{B}_E(G)$  which will be compared in this paper.

Let  $\text{Mod}_E$  be the category of  $E$ -vector spaces and let  $M \in \text{Mod}_E$  non zero. A *line* in  $M$  is a subspace of dimension 1. A *lattice*  $L$  in  $M$  is a  $O_E$ -submodule of  $M$  which contains *no line* and contains a basis of  $M$  over  $E$ . Note that a quotient of a lattice may contain a line. When the dimension of  $M$  over  $E$  is *countable*, a lattice  $L$  in  $M$  is a *free*  $O_E$ -submodule of  $M$  generated by a basis of  $M$  over  $E$  [9, I Appendice C.5]. Two lattices  $L, L'$  in  $M$  are *commensurable* when there exists an element  $a \in O_E$  such that  $aL \subset L', aL' \subset L$ . We denote by  $[L]$  the commensurability class of  $L$ .

**Remark 2.1.** — An  $O_E$ -submodule  $L$  of  $M \in \text{Mod}_E$  is a lattice in  $M$  if and only if any non zero element  $m \in M$  satisfies the two conditions:

- a) there exists an integer  $n \in \mathbf{N}$  such that  $\ell^n m$  belongs to  $L$ ,
- b) there exists an integer  $n \in \mathbf{N}$  such that  $\ell^{-n} m$  does not belong to  $L$ .

Two lattices  $L, L'$  in  $M$  are commensurable if and only if there exists an integer  $n \in \mathbf{N}$  such that  $\ell^n L \subset L', \ell^n L' \subset L$ .

A representation (= a linear action) of  $G$  on  $M$  is called *admissible* when  $\dim_E M^H < \infty$ , for any open pro- $\ell'$ -subgroup  $H$  of  $G$ , where  $M^H \in \text{Mod}_E$  is the subspace of  $H$ -invariant vectors of  $M$ . The representation  $M$  is called *irreducible* when  $M \neq 0$  and  $0$  and  $M$  are the only  $G$ -stable subspaces of  $M$ , *finitely generated* when  $M$  is a finitely generated  $EG$ -module, *of finite length* when there exists a finite  $G$ -stable filtration  $0 \subset M_1 \subset \dots \subset M_n = M$  with *irreducible quotients*. The length of the filtration and the isomorphism classes of the quotients, up to the order, do not depend on the choice of the filtration.

A *lattice*  $L$  in the *representation* of  $G$  on  $M$  will always be a  $G$ -stable lattice in  $M$ ; the lattice will be called *finitely generated* when it is a finitely generated  $O_E G$ -module. A representation of  $G$  on  $M$  containing a lattice is called *integral* (we do not suppose that the lattice is  $O_E$ -free as in [9]). There exist finitely generated lattices in a finitely generated integral representation; they form a commensurability class, and any lattice contains a finitely generated lattice.

A *continuous*  $E$ -representation of  $G$  is a topological Hausdorff  $E$ -vector space  $M$  equipped with a continuous action of  $G$ , i.e. such that the map  $(g, v) \rightarrow gv : G \times M \rightarrow M$  is continuous. It is called *topologically irreducible* when  $M \neq 0$  and  $0$  and  $M$  are the only *closed*  $G$ -stable subspaces of  $M$ . It is called of *finite topological length* when

there exists a finite filtration by  $G$ -stable *closed* subspaces  $0 \subset M_1 \subset \cdots \subset M_n = M$  with *topologically irreducible quotients*.

The category  $\mathcal{C}_E(G)$  of continuous representations of  $G$  on topological Hausdorff *complete*  $E$ -vector spaces with continuous  $G$ -equivariant  $E$ -linear morphisms, called *intertwining operators*, contains the subcategory  $\text{Mod}_E(G)$  of smooth representations and the subcategory  $\mathcal{B}_E(G)$  of Banach unitary representations, defined below. We indicate by the upper index *adm* or *fl* or *adm*, *fl* or *int* or *int*, *fl* the full subcategories representations which are admissible or of finite topological length or admissible and of finite topological length or integral or integral and of finite topological length. Example:  $\mathcal{C}_E(G)^{\text{adm}}$ ,  $\text{Mod}_E(G)^{\text{adm}}$ ,  $\mathcal{B}_E(G)^{\text{adm}}$  for admissible representations.

A representation of  $G$  on an  $E$ -vector space  $W$  is *smooth* when the stabilizer in  $G$  of any vector of  $W$  is open; this is simply a continuous representation of  $G$  on  $W$  when  $W$  is equipped with the discrete topology. The category  $\text{Mod}_E(G)$  of smooth  $E$ -representations of  $G$ , with morphisms the  $G$ -equivariant  $E$ -linear maps, is a full subcategory of  $\mathcal{C}_E(G)$ .

A *Banach unitary*  $E$ -representation  $V$  of  $G$  is a Hausdorff complete topological  $E$ -vector space with a topology given by a norm, equipped with a continuous action of  $G$  which respects the norm. A unit ball of  $V$  is  $L = \{v \in V : \|v\| \leq 1\}$  for some norm  $v \mapsto \|v\|$  on  $V$  defining the topology [Sch I.3, III]; it is a lattice in  $V$ . The unit balls of two norms on  $V$  giving the same topology are commensurable.

An  $E$ -linear map  $f : V_1 \rightarrow V_2$  between two Banach  $E$ -vector spaces  $V_1, V_2$  is continuous if and only if there exists some non zero  $a \in E$  such that  $f(L_1) \subset af(L_2)$  for some unit balls  $L_1, L_2$  of  $V_1, V_2$  [Sch I.3.1]. The topology quotient topology on the image of  $f$  is the topology induced by  $V_2$  if and only if  $f(L_1)$  and  $L_2 \cap f(V_1)$  are commensurable (this does not depend on the choice of the unit balls  $L_1, L_2$ ). When  $f$  is continuous and bijective, the inverse of  $f$  is continuous [Sch I.8.7].

We will compare  $\mathcal{B}_E(G)$  with the category  $\mathcal{M}_E(G)$  of smooth  $E$ -representations  $W$  of  $G$  equipped with a commensurability class  $[L]$  of lattices; a morphism  $(W, [L]) \rightarrow (W', [L'])$  is a morphism  $f : W \rightarrow W'$  in  $\text{Mod}_E(G)$  such that  $f(L) \subset aL'$  for some  $a \in E$ . The pair  $(W, [L])$  is called admissible or of finite length when  $W$  is admissible or of finite length, and  $\mathcal{M}_E(G)^{\text{adm}}$  or  $\mathcal{M}_E(G)^{\text{fl}}$  is the full subcategory of admissible or of finite length pairs in  $\mathcal{M}_E(G)$ .

**2.2. The two functors.** — We introduce two natural functors in opposite directions between the categories  $\mathcal{M}_E(G)$  and  $\mathcal{B}_E(G)$ .

There is the natural functor  $\mathcal{C}_E(G) \rightarrow \text{Mod}_E(G)$  sending  $M \in \mathcal{C}_E(G)$  to its *smooth part*

$$M^\infty := \cup_H M^H,$$

for all open pro- $\ell'$ -subgroups  $H$  of  $G$ . When  $V \in \mathcal{B}_E(G)$  is a Banach unitary representation of  $G$ , the smooth part  $L^\infty = V^\infty \cap L$  of a unit ball  $L$  of  $V$  is a lattice of  $V^\infty$ . Two unit balls of  $V$  are commensurable and their smooth parts are commensurable, hence  $(V^\infty, [L^\infty]) \in \mathcal{M}_E(G)$  is well defined. A continuous morphism  $f : V_1 \rightarrow V_2$  of Banach unitary  $E$ -representations of  $G$  with unit balls  $L_1, L_2$ , restricts to a morphism

$f^\infty : (V_1^\infty, [L_1^\infty]) \rightarrow (V_2^\infty, [L_2^\infty])$ . We get a functor

$$\mathcal{B}_E(G) \rightarrow \mathcal{M}_E(G).$$

In the opposite direction there is the natural functor

$$\mathcal{M}_E(G) \rightarrow \mathcal{B}_E(G)$$

sending  $(W, [L])$  to the *completion* of  $W$  for the  $L$ -adic topology [Sch 7.5]:

$$\hat{W}_L := \varprojlim_n W/\ell^n L \simeq E \otimes_{O_E} \hat{L}, \quad \hat{L} := \varprojlim_n L/\ell^n L.$$

Any element  $v \in \hat{W}_L$  is written

$$(1) \quad v = (w_n + \ell^n L)_n, \quad w_n \in W, \quad w_{n+1} \in w_n + \ell^n L,$$

for all  $n \in \mathbf{N}$ . The lattice  $\hat{L}$  is a unit ball of  $\hat{W}_L$  for the gauge norm  $\|v\| = \inf_{a \in E, v \in a\hat{L}} |a|$ . The completions of  $W$  defined by two commensurable lattices of  $W$  are the same. The group  $G$  acts naturally on  $\hat{W}_L$ , for  $g \in G$  and  $v$  as above,

$$gv = (gw_n + \ell^n L)_{n \in \mathbf{N}},$$

and  $\hat{W}_L$  is a Banach unitary  $E$ -representation of  $G$  of unit ball  $\hat{L}$ , well defined by  $(W, [L])$ . A morphism  $f : (W, [L]) \rightarrow (W', [L'])$  in  $\mathcal{M}_E(G)$  extends by continuity to an intertwining operator  $\hat{f} : \hat{W}_L \rightarrow \hat{W}'_L$ .

**Remark 2.2.** — *The map  $W \mapsto \hat{W}_L$  sending  $w$  to  $(w + \ell^n L)_{n \in \mathbf{N}}$  is injective, because  $L$  contains no line. We will identify  $W$  with its image in  $\hat{W}_L$ .*

**2.3.** — To study the two functors, smooth part and completion, between  $\mathcal{M}_E(G)$  and  $\mathcal{B}_E(G)$ , the key point is the exactness of the  $H$ -invariants functor.

**Proposition 2.3.** — *Let  $H$  be any open pro- $\ell'$ -subgroup of  $G$ . The  $H$ -invariants functor*

$$M \mapsto M^H : \mathcal{C}_E(G) \rightarrow \text{Mod}_E$$

*is exact.*

*Proof.* — This is well known for the subcategory  $\text{Mod}_E(G)$  of smooth representations in  $\mathcal{C}_E(G)$ . The exactness results from the existence of a Haar  $O_E$ -measure  $dg$  on  $G$  such that the volume  $\text{vol}(H, dg)$  of  $H$  is a unit in  $O_E$ . The function  $e_H$  equal to  $\text{vol}(H, dg)^{-1}$  on  $H$  and 0 on  $G - H$ , is an idempotent in the convolution algebra  $C_c^\infty(G; O_E)$  of locally constant compactly supported functions  $G \rightarrow O_E$ , for the Haar measure  $dg$ . The idempotent  $e_H$  acts on  $M \in \mathcal{C}_E(G)$ , as follows. One chooses a decreasing sequence of normal subgroups  $H_n$  of  $H$  of finite index such that  $\bigcap_{n \in \mathbf{N}} H_n$  is trivial, and a system of representatives  $X_n$  in  $H$  of  $H/H_n$ . The continuity of the action of  $G$  on  $M$  implies that the sequence

$$v_n = [H : H_n]^{-1} \sum_{g \in X_n} gv$$

converges to a unique element  $e_H * v$  in the Hausdorff complete space  $M$ . This element  $e_H * v$  does not depend on the choice of  $(H_n, X_n)_{n \in \mathbb{N}}$  and clearly  $v \mapsto e_H * v$  is a linear projector  $M \rightarrow M^H$  of its  $H$ -invariants.  $\square$

**Corollary 2.4.** — *The smooth part functor  $M \mapsto M^\infty : \mathcal{C}_E(G) \rightarrow \text{Mod}_E(G)$  is exact.*

**Proposition 2.5.** — *A Banach unitary  $E$ -representation  $V$  of  $G$  is equal to the closure of its smooth part  $V^\infty$ .*

*Proof.* — Let  $v$  be an arbitrary element of  $V$  and let  $L$  be a unit ball of  $V$ . For any integer  $n \geq 1$ , there is an open pro- $\ell'$ -subgroup  $H_n$  of  $G$  such that  $H_n v \subset v + \ell^n L$ , by the continuity of the action of  $G$  on  $V$ . The element  $e_{H_n} * v$  is fixed by  $H_n$  and belongs to  $v + \ell^n L$ . The element  $(e_{H_n} * v + \ell^n L)_{n \in \mathbb{N}}$  belongs to the closure of  $V^\infty$  and is equal to  $v$ .  $\square$

**Corollary 2.6.** — *The smooth part functor  $\mathcal{B}_E(G) \rightarrow \mathcal{M}_E(G)$  is fully faithful.*

*Proof.* — For  $i = 1, 2$ , let  $V_i \in \mathcal{B}_E(G)$  with unit ball  $L_i$ . The embedding  $V_i^\infty \rightarrow (V_i^\infty)_{L_i}^\wedge$  extends by continuity to an isomorphism  $\tau_i : V_i \rightarrow (V_i^\infty)_{L_i}^\wedge$  in  $\mathcal{B}_E(G)$  by the Proposition 2.5 and its proof. We deduce that arbitrary intertwining operators  $\phi : (V_1^\infty, [L_1^\infty]) \rightarrow (V_2^\infty, [L_2^\infty])$  and  $f : V_1 \rightarrow V_2$  satisfy

$$\phi = (\tau_2^{-1} \hat{\phi} \tau_1)^\infty \quad , \quad f = \tau_2^{-1} (f^\infty) \tau_1 \quad . \quad \square$$

We show that the completion commutes with the  $H$ -invariants.

**Proposition 2.7.** — *Let  $V$  be the completion of an integral smooth  $E$ -representation  $W$  of  $G$  with respect to a lattice  $L$ , and let  $H$  be an open pro- $\ell'$ -subgroup of  $G$ . The  $H$ -invariants  $V^H$  of  $V$  is equal to the closure of  $W^H$  in  $V$ ,*

$$V^H = \overline{W^H}.$$

*Proof.* — For  $X = W, L$  or  $V$ , we have  $e_H * X = X^H$ . Let  $v = (w_n + \ell^n L)_{n \in \mathbb{N}}$  be an element of  $V$  as in (1). Then  $e_H * w_{n+1} \in e_H * w_n + \ell^n L$ , and  $e_H * v = (e_H * w_n + \ell^n L)_{n \in \mathbb{N}}$ .  $\square$

**Corollary 2.8.** — *An admissible smooth  $E$ -representation of  $G$  with a commensurability class of lattices is equal to the smooth part of its completion.*

*Proof.* — When the representation  $W$  is admissible, the  $E$ -vector space  $W^H$  is finite dimensional and already complete, hence  $V^H = W^H$  in the Proposition 2.7.  $\square$

It is clear that the functor smooth part respects admissible representations, the corollary shows that the completion respects also admissible representations.

**Theorem 2.9.** — *The smooth part and completion are inverse equivalences of categories between  $\mathcal{M}_E^{\text{adm}}(G)$  and  $\mathcal{B}_E^{\text{adm}}(G)$ .*

*Proof.* — Proposition 2.5, Corollaries 2.6, 2.8.  $\square$

In particular, the smooth part and the completion induce inverse equivalences of categories between admissible and of finite topological length representations  $\mathcal{M}_E^{\text{adm,fl}}(G)$  and  $\mathcal{B}_E^{\text{adm,fl}}(G)$ .

We consider now  $\ell$ -adic representations of  $G$ . For any *finite* extensions  $E'/E/\mathbf{Q}_\ell$  contained in a fixed algebraic closure  $\overline{\mathbf{Q}}_\ell$ , the scalar extension  $s_{E'/E}$  from  $E$  to  $E'$

$$\mathcal{C}_E(G) \rightarrow \mathcal{C}_{E'}(G)$$

sends  $M \in \mathcal{C}_E(G)$  to  $M_{E'} := E' \otimes_E M = \oplus(e_i \otimes M)$ , for a finite basis  $(e_i)$  of the  $E$ -vector space  $E'$ , with the topology induced by  $M$  (independent of the choice of the basis) and a morphism  $f : M \rightarrow M'$  in  $\mathcal{C}_E(G)$  to  $\text{id}_{E'} \otimes f$ . The inductive limit

$$\mathcal{C}_\ell(G) := \lim_{s_{E'/E}} \mathcal{C}_E(G)$$

is the category of  $\ell$ -adic representations of  $G$ . The scalar extension respects smooth representations, and the inductive limit

$$\text{Mod}_\ell(G) := \lim_{s_{E'/E}} \text{Mod}_E(G)$$

is the category of *smooth*  $\ell$ -adic representations of  $G$ , which is a (not full) subcategory of the classical category  $\text{Mod}_{\overline{\mathbf{Q}}_\ell}(G)$  of smooth  $\overline{\mathbf{Q}}_\ell$ -representations of  $G$ .

Let  $R/E$  be any extension contained in  $\overline{\mathbf{Q}}_\ell$  and let  $O_R$  be the ring of integers in  $R$ . As the dimension of  $R/E$  is countable,  $O_R$  is an  $O_E$ -free module [9, Appendice C, C.4]. We denote by  $L_{O_R} := O_R \otimes_{O_E} L$ , the scalar extension from  $O_E$  to  $O_R$  of an  $O_E$ -module  $L$ .

**Lemma 2.10.** — *Let  $M \in \mathcal{C}_E(G)$  equipped with a lattice  $L$ .*

(i) *The  $H$ -invariants commute with the scalar extension,  $e_H M_R = (e_H M)_R$ ,  $e_H L_{O_R} = (e_H L)_{O_R}$  for any pro- $\ell'$ -subgroup  $H$  of  $G$  (this is true for any extension  $R/E$  of fields of characteristic different from  $p$ ). In particular,  $M$  is admissible if and only if  $M_R$  is admissible.*

(ii) *The intersection  $L = M \cap L'$  of a lattice  $L'$  of  $M_R$  with  $M$  is a lattice in  $M$ . In particular, if  $M_R$  is integral then  $M$  is integral.*

(iii) *The scalar extension  $L_{O_R}$  of a lattice  $L$  in  $M$  is a lattice in  $M_R$ . In particular, if  $M$  is integral then  $M_R$  is integral.*

(iv) *Two lattices  $L, L'$  of  $M$  are commensurable if and only if their scalar extensions  $L_{O_R}, L'_{O_R}$  are commensurable.*

*Proof.* — (i) is clear. The other properties are clear using the Remark 2.1 and  $L_{O_R} = \oplus_i(e_i \otimes L)$  for a basis  $(e_i)$  of the free  $O_E$ -module  $O_R$ .  $\square$

**Lemma 2.11.** — *The scalar extension  $s_{E'/E}$  from  $E$  to a finite extension  $E'$  commutes with the smooth part functor  $\mathcal{C}_E(G) \rightarrow \text{Mod}_E(G)$  and with the smooth part and completion functors between  $\mathcal{B}_E(G)$  and  $\mathcal{M}_E(G)$ .*

*Proof.* — We choose a basis  $(e_i)$  of the free  $O_E$ -module  $O_{E'}$ . The scalar extension  $V_{E'} = \oplus_i(e_i \otimes V)$  of the completion  $V$  of  $(W, [L]) \in \mathcal{M}_E(G)$  is clearly the completion of the scalar extension  $(W_{E'} = \oplus_i(e_i \otimes W), [L_{O_{E'}} = \oplus_i(e_i \otimes L)])$  of  $(W, [L])$ . The



scalar extension of the  $H$ -invariants of  $V \in \mathcal{B}_E(G)$  is the  $H$ -invariants of the scalar extension  $V_{E'}$  (Lemma 2.10).  $\square$

As the scalar extension  $s_{E'/E}$  from  $E$  to a finite extension  $E'$  respects admissibility, lattices, commensurability of lattices, Banach spaces (Lemma 2.10), the inductive limit over  $s_{E'/E}$  for all finite extensions  $E'/E/\mathbf{Q}_\ell$  contained in  $\overline{\mathbf{Q}_\ell}$ , defines the categories

- a)  $\mathcal{C}_\ell(G)^{\text{adm}}$  of admissible  $\ell$ -adic representations of  $G$ ,
- b)  $\text{Mod}_\ell(G)^{\text{int}}$  of integral smooth  $\ell$ -adic representations,
- b)  $\mathcal{M}_\ell(G)$  of smooth  $\ell$ -adic representations of  $G$  equipped with a commensurability class of lattices,
- c)  $\mathcal{B}_\ell(G)$  of Banach unitary  $\ell$ -adic representations of  $G$ .

We define the completion and smooth part functors between  $\mathcal{M}_\ell(G)$  and  $\mathcal{B}_\ell(G)$  using the Lemma 2.11.

**Theorem 2.12.** — *The completion and smooth part functors induce equivalence of categories between the categories  $\mathcal{M}_\ell(G)^{\text{adm}}$  and  $\mathcal{B}_\ell(G)^{\text{adm}}$ .*

*Proof.* — Theorem 2.9.  $\square$

### 3.

Let  $G_F$  be the group of rational points of a connected reductive group over a local non Archimedean field  $F$  of residual characteristic  $p$ . The group  $G_F$  is a locally pro- $p$ -group. As before  $E/\mathbf{Q}_\ell$  is a finite extension contained in  $\overline{\mathbf{Q}_\ell}$  and  $\ell \neq p$ .

**Proposition 3.1.** — *Let  $R/R_o$  be any extension of fields of characteristic different from  $p$ . Then  $W \in \text{Mod}_{R_o}(G_F)$  has finite length if and only if  $W_R \in \text{Mod}_R(G_F)$  has finite length.*

*Proof.* — [9, II.4.3.c].  $\square$

**Proposition 3.2.** — *Any finite length smooth representation  $W$  of  $G_F$  over a field of characteristic different from  $p$  is admissible.*

*Proof.* — This is proved in [9, II.2.8] when the field is algebraically closed. The scalar extension is not sensitive to admissibility and finite length (Lemma 2.10, Proposition 3.1) for any extension of fields of characteristic different from  $p$ .  $\square$

**Proposition 3.3.** — *The lattices in an integral finite length representation  $W \in \text{Mod}_E(G_F)$  are commensurable (hence finitely generated).*

*Proof.* — This is proved [11, th.1] when the field is  $\overline{\mathbf{Q}_\ell}$ . The scalar extension is not sensitive to integrality, commensurability of lattices, and finite length, (Proposition 2.10, Proposition 3.1).  $\square$

**Remark 3.4.** — *One cannot replace “finite length” by “admissible” in the Proposition 3.3.*

**Lemma 3.5.** — *The category of smooth  $\overline{\mathbf{Q}}_\ell$ -representation of  $G_F$  of finite length is equal to the category of smooth  $\ell$ -adic representations of  $G_F$  of finite length,*

$$\mathrm{Mod}_{\overline{\mathbf{Q}}_\ell}(G_F)^{fl} \simeq \mathrm{Mod}_\ell(G_F)^{fl}.$$

*Proof.* — Let  $W \in \mathrm{Mod}_{\overline{\mathbf{Q}}_\ell}(G_F)^{fl}$ . There exists a finite extension  $E/\mathbf{Q}_\ell$  and  $W_E \in \mathrm{Mod}_E(G_F)^{fl}$  such that  $W$  is the scalar extension of  $W_E$ . When  $W$  is irreducible, this is proved in [9, II.4.7]. In general, let  $H$  be an open pro- $p$ -subgroup of  $G_F$  such that the length of  $W$  is equal to the length of the module  $e_H W$  over the Hecke algebra  $\mathrm{End}_{\overline{\mathbf{Q}}_\ell G_F} \overline{\mathbf{Q}}_\ell[G_F/H]$ . Let  $(w_i)$  be a finite  $\overline{\mathbf{Q}}_\ell$ -basis of  $e_H W$ . The convolution algebra  $\mathrm{End}_{\mathbf{Z}_\ell G_F} \mathbf{Z}_\ell[G_F/H]$  is finitely generated [9, II.2.13], and the dimension of  $e_H W$  over  $\overline{\mathbf{Q}}_\ell$  is finite. Hence there exists a finite extension  $E/\mathbf{Q}_\ell$  such that the  $E$ -vector space  $\oplus_i E w_i$  in  $e_H W$  is stable by the Hecke algebra  $\mathrm{End}_{EG_F} E[G_F/H]$ . The  $E$ -representation  $U$  of  $G_F$  generated by  $(w_i)$  in  $W$  satisfies  $e_H U = \oplus_i E w_i$ . The scalar extension  $\overline{\mathbf{Q}}_\ell \otimes_E U$  is equal to  $W$  because it is a subrepresentation of  $W$  with the same  $H$ -invariants. By the Proposition 3.1,  $W_E$  has finite length.

Let  $E, E'$  be two finite extensions. Let  $W_E \in \mathrm{Mod}_E(G_F)^{fl}$ ,  $W_{E'} \in \mathrm{Mod}_{E'}(G_F)^{fl}$ , let

$$f : \overline{\mathbf{Q}}_\ell \otimes_E W_E \rightarrow \overline{\mathbf{Q}}_\ell \otimes_{E'} W_{E'}$$

be a  $\overline{\mathbf{Q}}_\ell G_F$ -morphism between their scalar extensions to  $\overline{\mathbf{Q}}_\ell$ . There exists a finite extension  $E''$  containing  $E, E'$  such that  $f$  is defined on  $E''$ , i.e. induces a  $E'' G_F$ -morphism  $f_{E''} : E'' \otimes_E W_E \rightarrow E'' \otimes_{E'} W_{E'}$  between their scalar extensions to  $E''$  [9, proof of II.4.7].  $\square$

The scalar extension  $s_{E'/E}$  for smooth representations of  $G_F$  respects finite length (Proposition 3.1) and the category  $\mathrm{Mod}_\ell(G_F)^{fl}$  of smooth  $\ell$ -adic representations of  $G_F$  of finite length is well defined, contained in the category  $\mathrm{Mod}_\ell(G_F)^{\mathrm{adm}}$  of admissible smooth  $\ell$ -adic representations of  $G_F$  (Proposition 3.2). The category  $\mathrm{Mod}_\ell(G_F)^{\mathrm{int}, \mathrm{fl}}$  of integral smooth  $\ell$ -adic representations of  $G_F$  of finite length is equivalent by the forgetful functor composite with the completion and the smooth part to the category  $\mathcal{B}_\ell(G_F)^{\mathrm{adm}, \mathrm{fl}}$  of Banach unitary  $\ell$ -adic representations which are admissible and of finite length.

**Theorem 3.6.** — *The completion and the smooth part define equivalence of categories between  $\mathrm{Mod}_{\overline{\mathbf{Q}}_\ell}^{\mathrm{int}, \mathrm{fl}}(G_F)$  and  $\mathcal{B}_\ell(G_F)^{\mathrm{adm}, \mathrm{fl}}$ .*

*In particular, they give bijections between the irreducible integral smooth  $\overline{\mathbf{Q}}_\ell$ -representations of  $G_F$  and the topologically irreducible admissible Banach unitary  $\ell$ -adic representations of  $G_F$ .*

For  $G_F = GL(n, F)$ , we deduce the  $\ell$ -adic local Langlands correspondence for  $GL(n, F)$ , given in the introduction.

A very natural question (asked by the referee) for a Banach unitary  $\ell$ -adic representation  $V$  of  $G_F$  (notations of the Sections 2 and 3) is: does  $V$  topologically irreducible imply  $V$  admissible? The answer is no.