

*quatrième série - tome 49      fascicule 2      mars-avril 2016*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Hélène ESNAULT & Olivier WITTENBERG

*On the cycle class map for zero-cycles over local fields*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> janvier 2016

N. ANANTHARAMAN I. GALLAGHER  
P. BERNARD B. KLEINER  
E. BREUILLARD E. KOWALSKI  
R. CERF M. MUSTAȚĂ  
A. CHAMBERT-LOIR L. SALOFF-COSTE

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

### Édition / *Publication*

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

### Abonnements / *Subscriptions*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

### Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

---

© 2016 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

# ON THE CYCLE CLASS MAP FOR ZERO-CYCLES OVER LOCAL FIELDS

BY HÉLÈNE ESNAULT AND OLIVIER WITTENBERG  
WITH AN APPENDIX BY SPENCER BLOCH

---

**ABSTRACT.** – We study the Chow group of 0-cycles of smooth projective varieties over local and strictly local fields. We prove in particular the injectivity of the cycle class map to integral  $\ell$ -adic cohomology for a large class of surfaces with positive geometric genus, over local fields of residue characteristic  $\neq \ell$ . The same statement holds for semistable  $K3$  surfaces defined over  $\mathbf{C}((t))$ , but does not hold in general for surfaces over strictly local fields.

**RÉSUMÉ.** – Nous étudions le groupe de Chow des 0-cycles des variétés projectives et lisses sur les corps locaux et strictement locaux. Nous prouvons en particulier l'injectivité de l'application classe de cycle vers la cohomologie  $\ell$ -adique entière pour de nombreuses surfaces de genre géométrique non nul, sur les corps locaux de caractéristique résiduelle  $\neq \ell$ . Le même énoncé vaut pour les surfaces  $K3$  semi-stables définies sur  $\mathbf{C}((t))$ , mais ne vaut pas en général pour les surfaces sur les corps strictement locaux.

## 1. Introduction

Let  $X$  be a smooth projective variety over a field  $K$ , let  $\mathrm{CH}_0(X)$  denote the Chow group of 0-cycles on  $X$  up to rational equivalence and let  $A_0(X) \subset \mathrm{CH}_0(X)$  be the subgroup of cycle classes of degree 0.

When  $K$  is algebraically closed, the group  $A_0(X)$  is divisible and its structure as an abelian group is, conjecturally, rather well understood, thanks to Roitman's theorem and to the Bloch-Beilinson-Murre conjectures. A central tool for the study of  $A_0(X)$  over other types of fields is the cycle class map

$$\psi : \mathrm{CH}_0(X)/n\mathrm{CH}_0(X) \rightarrow H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d))$$

to étale cohomology, where  $n$  denotes an integer invertible in  $K$  and  $d = \dim(X)$ . The group  $H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d))$  is easier to understand, thanks to the Hochschild-Serre spectral

---

The first author is supported by the the ERC Advanced Grant 226257, the Chaire d'excellence 2011 of the Fondation Sciences Mathématiques de Paris and the Einstein Foundation.

sequence; for instance, if  $K$  has cohomological dimension  $\leq 1$  and  $X$  is simply connected, then  $H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d)) = \mathbf{Z}/n\mathbf{Z}$  and  $\psi$  may be interpreted as the degree map.

According to one of the main results of higher-dimensional unramified class field theory, due to Kato and Saito [43], if  $K$  is a finite field, the group  $A_0(X)$  is finite and  $\psi$  is an isomorphism. More recently, Saito and Sato [69] have shown that if  $K$  is the quotient field of an excellent Henselian discrete valuation ring with finite or separably closed residue field, the group  $A_0(X)$  is the direct sum of a finite group of order prime to  $p$  and a group divisible by all integers prime to  $p$  (see also [11, Théorème 3.25]). In this case, however, the map  $\psi$  need not be either injective or surjective. What Saito and Sato prove, instead, is the bijectivity of the analogous cycle class map for cycles of dimension 1 on regular models of  $X$  over the ring of integers of  $K$  (see [69, Theorem 1.16]).

Following a method initiated by Bloch [4], one may approach the torsion subgroup of  $A_0(X)$ , as well as the kernel of  $\psi$ , with the help of algebraic K-theory, when  $X$  is a surface. We refer to [10] for a detailed account of this circle of ideas. Strong results were obtained in this way for rational surfaces, and more generally for surfaces with geometric genus zero, over number fields,  $p$ -adic fields, and fields of characteristic 0 and cohomological dimension 1 (see [5], [6], [16], [9], [14], [68]). We note that over algebraically closed fields of characteristic 0, surfaces with geometric genus zero are those surfaces for which the Chow group  $A_0(X)$  should be representable, according to Bloch's conjecture (see [5, § 1]).

The first theorem of this paper establishes the injectivity of  $\psi$  for a large class of surfaces over local fields, when  $n$  is divisible enough and prime to the residue characteristic, without any assumption on the geometric genus. In principle, this theorem should be applicable to all simply connected surfaces, a generality in which the injectivity of  $\psi$  may not have been expected. Before we state it, we set up some notation. Let  $\mathcal{X}$  be a regular proper flat scheme over an excellent Henselian discrete valuation ring  $R$ . Let  $X = \mathcal{X} \otimes_R K$  and  $A = \mathcal{X} \otimes_R k$  denote the generic fiber and the special fiber, respectively. We assume the reduced special fiber  $A_{\text{red}}$  has simple normal crossings, and write  $\text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell = \varprojlim \text{CH}_0(X)/\ell^n \text{CH}_0(X)$ .

**THEOREM 1** (Theorem 3.1 and Remark 3.2). – *Assume the residue field  $k$  is finite and  $X$  is a surface whose Albanese variety has potentially good reduction. If the irreducible components of  $A$  satisfy the Tate conjecture, then for any  $\ell$  invertible in  $k$ , the cycle class map*

$$(1.1) \quad \text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_{\text{ét}}^4(X, \mathbf{Z}_\ell(2))$$

*is injective. Equivalently, the natural pairing  $\text{CH}_0(X) \times \text{Br}(X) \rightarrow \mathbf{Q}/\mathbf{Z}$  is non-degenerate on the left modulo the maximal  $\ell$ -divisible subgroup of  $\text{CH}_0(X)$ .*

The assumption on the irreducible components of  $A$  holds as soon as  $X$  has geometric genus zero, as well as in many examples of nontrivial degenerations of surfaces with nonzero geometric genus (see § 3). Theorem 1 is due to Saito [68] when  $X$  is a surface with geometric genus zero over a  $p$ -adic field. An example of Parimala and Suresh [62] shows that the assumption on the Albanese variety cannot be removed. Finally, we note that Theorem 1 may be viewed as a higher-dimensional generalization of Lichtenbaum-Tate duality for curves, according to which the natural pairing  $\text{CH}_0(X) \times \text{Br}(X) \rightarrow \mathbf{Q}/\mathbf{Z}$  is non-degenerate if  $X$  is a smooth proper curve over a  $p$ -adic field (see [54]).

Our starting point for the proof of Theorem 1 is the theorem of Saito and Sato alluded to above about the cycle class map for 1-cycles on  $\mathcal{X}$  [69, Theorem 1.16], which allows us to express the kernel of  $\psi$  purely in terms of the scheme  $A$  and of the cohomology of  $X$ , when  $k$  is either finite or separably closed (Theorem 2.1 and Theorem 2.2). The dimension of  $X$  plays no role in this part of the argument; an application to the study of 0-cycles on a cubic threefold over  $\mathbf{Q}_p$  may be found in Example 2.12. Theorem 1 is then obtained by analyzing the various cohomology groups which appear in the resulting expression for  $\text{Ker}(\psi)$ . More precisely, in the situation of Theorem 1, we prove the stronger assertion that the 1-dimensional cycle class map  $\psi_{1,A} : \text{CH}_1(A) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_A^4(\mathcal{X}, \mathbf{Z}_\ell(2))$  to integral  $\ell$ -adic étale homology is surjective. This provides a geometric explanation for the assumption that the Albanese variety of  $X$  have potentially good reduction, a condition which first appeared in [68] and which turns out to be essential for the surjectivity of  $\psi_{1,A}$  to hold (see Lemma 3.7 and § 4.3).

When the residue field  $k$  is separably closed instead of finite, the arguments used in the proof of Theorem 1 fail in several places. They still lead to the following statement, which may also be deduced from results of Colliot-Thélène and Raskind [13] (see § 4 for comments on this point).

**THEOREM 2** (Theorem 4.1 and Remark 4.3). – *Assume  $k$  is separably closed and  $K$  has characteristic 0. If  $X$  is a surface with geometric genus zero, then for any  $\ell$  invertible in  $k$ , the cycle class map*

$$(1.2) \quad \text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_{\text{ét}}^4(X, \mathbf{Z}_\ell(2))$$

*is injective. If in addition  $X$  is simply connected, then  $A_0(X)$  is divisible by  $\ell$  and the unramified cohomology group  $H_{\text{nr}}^3(X, \mathbf{Q}_\ell/\mathbf{Z}_\ell(2))$  vanishes.*

This leaves open the question of the injectivity of the cycle class map (1.2) when  $k$  is separably closed and  $X$  is a surface with positive geometric genus over  $K$ . In this situation, the 1-dimensional cycle class map  $\psi_{1,A}$  is far from being surjective. Building on the work of Kulikov, Persson, Pinkham [51] [64], and of Miranda and Morrison [59], we nevertheless give a positive answer for semistable K3 surfaces over  $\mathbf{C}((t))$ .

**THEOREM 3** (Theorem 5.1). – *Let  $X$  be a K3 surface over  $\mathbf{C}((t))$ . If  $X$  has semistable reduction, the group  $A_0(X)$  is divisible.*

The proof of Theorem 3 hinges on the precise knowledge of the combinatorial structure of a degeneration of  $X$ . It would go through over the maximal unramified extension of a  $p$ -adic field, as far as prime-to- $p$  divisibility is concerned, if similar knowledge were available. This is in marked contrast with the situation over  $p$ -adic fields, where such knowledge is not necessary for the proof of Theorem 1.

In the final section of this paper, with the help of Ogg-Shafarevich theory and of a construction due to Persson, we show that the hope for a statement analogous to Theorem 1 over the quotient field of a strictly Henselian excellent discrete valuation ring is in fact too optimistic, even over the maximal unramified extension of a  $p$ -adic field.