

*quatrième série - tome 52      fascicule 1      janvier-février 2019*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Jean-Baptiste TEYSSIER

*Skeletons and moduli of Stokes torsors*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Patrick BERNARD

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE

de 1883 à 1888 par H. DEBRAY

de 1889 à 1900 par C. HERMITE

de 1901 à 1917 par G. DARBOUX

de 1918 à 1941 par É. PICARD

de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> mars 2019

P. BERNARD

D. HARARI

S. BOUCKSOM

A. NEVES

R. CERF

J. SZEFTTEL

G. CHENEVIER

S. VŨ NGỌC

Y. DE CORNULIER

A. WIENHARD

A. DUCROS

G. WILLIAMSON

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,

45, rue d'Ulm, 75230 Paris Cedex 05, France.

Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.

[annales@ens.fr](mailto:annales@ens.fr)

---

## Édition et abonnements / *Publication and subscriptions*

Société Mathématique de France

Case 916 - Luminy

13288 Marseille Cedex 09

Tél. : (33) 04 91 26 74 64

Fax : (33) 04 91 41 17 51

email : [abonnements@smf.emath.fr](mailto:abonnements@smf.emath.fr)

### Tarifs

Abonnement électronique : 420 euros.

Abonnement avec supplément papier :

Europe : 551 €. Hors Europe : 620 € (\$ 930). Vente au numéro : 77 €.

---

© 2019 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Stéphane Seuret

Périodicité : 6 n<sup>os</sup> / an

## SKELETONS AND MODULI OF STOKES TORSORS

BY JEAN-BAPTISTE TEYSSIER

---

**ABSTRACT.** – We prove an analog for Stokes torsors of Deligne’s skeleton conjecture and deduce from it the representability of the functor of relative Stokes torsors by an affine scheme of finite type over  $C$ . This provides, in characteristic 0, a local analog of the existence of a coarse moduli for skeletons with bounded ramification, due to Deligne. As an application, we use the geometry of this moduli to derive quite strong finiteness results for integrable systems of differential equations in several variables which did not have any analog in one variable.

**RÉSUMÉ.** – Nous prouvons une variante pour les toseurs de Stokes de la conjecture des squelettes de Deligne, et en déduisons la représentabilité du foncteur des toseurs de Stokes relatifs par un schéma affine de type fini sur  $C$ . Cela fournit, en caractéristique 0, un analogue local de l’existence d’un espace de modules grossier pour les squelettes à ramification bornée, dû à Deligne. À titre d’application, nous utilisons la géométrie de cet espace de modules pour en déduire de nouveaux résultats de finitude sur les systèmes intégrables d’équations différentielles à plusieurs variables qui n’avaient pas d’analogue à une variable.

Consider the following linear differential equation  $(E)$  with polynomial coefficients:

$$p_n \frac{d^n f}{dz^n} + p_{n-1} \frac{d^{n-1} f}{dz^{n-1}} + \cdots + p_1 \frac{df}{dz} + p_0 f = 0.$$

If  $p_n(0) \neq 0$ , Cauchy theorem asserts that a holomorphic solution to  $(E)$  defined on a small disk around 0 is equivalent to the values of its  $n$  first derivatives at 0. If  $p_n(0) = 0$ , holomorphic solutions to  $(E)$  on a small disk around 0 may always be zero. Nonetheless,  $(E)$  may have formal power series solutions. The *Main asymptotic development theorem* [26], due to Hukuhara and Turrittin asserts that for a direction  $\theta$  emanating from 0, and for a formal power series solution  $f$ , there is a sector  $\mathcal{S}_\theta$  containing  $\theta$  such that  $f$  can be “lifted” in a certain sense to a holomorphic solution  $f_\theta$  of  $(E)$  on  $\mathcal{S}_\theta$ . We say that  $f_\theta$  is asymptotic to  $f$  at 0. If  $f_\theta$  is analytically continued around 0 into a solution  $\tilde{f}_\theta$  of  $(E)$  on the sector  $\mathcal{S}_{\theta'}$ , where  $\theta' \neq \theta$ , it may be that the asymptotic development of  $\tilde{f}_\theta$  at 0 is not  $f$  any more. This is the *Stokes phenomenon*. As a general principle, the study of  $(E)$  amounts to the study of its “formal type” and the study of how asymptotic developments of solutions jump via

analytic continuation around 0. To organize these informations, it is traditional to adopt a linear algebra point of view.

The equation (E) can be seen as a *differential module*, i.e., a finite dimensional vector space  $\mathcal{N}$  over the field  $C\{z\}[z^{-1}]$  of convergent Laurent series, endowed with a  $C$ -linear endomorphism  $\nabla : \mathcal{N} \rightarrow \mathcal{N}$  satisfying the Leibniz rule. In this language, solutions of (E) correspond to elements of  $\text{Ker } \nabla$  (also called *flat sections of  $\nabla$* ). Furthermore, a differential equation with same “formal type” as (E) corresponds to a differential module  $\mathcal{M}$  with an isomorphism of formal differential modules  $\text{iso} : \mathcal{M}_{\hat{0}} \rightarrow \mathcal{N}_{\hat{0}}$ . Since  $\text{iso}$  can be seen as a formal flat section of the differential module  $\text{Hom}({}_{\mathcal{O}}\mathcal{M}, \mathcal{N})$ , the main asymptotic development theorem applies to it. The lifts of  $\text{iso}$  to sectors thus produce a cocycle  $\gamma := (\text{iso}_{\theta} \text{iso}_{\theta'}^{-1})_{\theta, \theta' \in S^1}$  with value into the sheaf of sectorial automorphisms of  $\mathcal{N}$  which are asymptotic to  $\text{Id}$  at 0. This is the *Stokes sheaf* of  $\mathcal{N}$ , denoted by  $\text{St}_{\mathcal{N}}(C)$ . A fundamental result of Malgrange [17] and Sibuya [25] implies that  $(\mathcal{M}, \text{iso})$  is determined by the torsor under  $\text{St}_{\mathcal{N}}(C)$  associated to the cocycle  $\gamma$ . Hence, *Stokes torsors encode in an algebraic way analytic data and classifying differential equations amounts to studying Stokes torsors*. As a result, the study of Stokes torsors is meaningful.

In higher dimension, the role played by differential modules is played by *good meromorphic connections*. We will take such a connection  $\mathcal{N}$  defined around  $0 \in C^n$  to be of the shape

$$(0.0.1) \quad \mathcal{E}^{a_1} \otimes \mathcal{R}_{a_1} \oplus \cdots \oplus \mathcal{E}^{a_d} \otimes \mathcal{R}_{a_d},$$

where the  $a_i$  are meromorphic functions with poles contained in a normal crossing divisor  $D$ , where  $\mathcal{E}^{a_i}$  stands for the rank one connection  $(\mathcal{O}_{C^n,0}(*D), d - da_i)$ , and where the  $\mathcal{R}_{a_i}$  are regular connections. Note that from works of Kedlaya [12, 13] and Mochizuki [20, 22], every meromorphic connection is (up to ramification) formally isomorphic at each point to a connection of the form (0.0.1) at the cost of blowing-up enough the pole locus. If  $(r_i, \theta_i)_{i=1, \dots, n}$  are the usual polar coordinates on  $C^n$ , the Stokes sheaf  $\text{St}_{\mathcal{N}}$  of  $\mathcal{N}$  is a sheaf of complex unipotent algebraic groups over the torus  $T := (S^1)^n$  defined by  $r_1 = \cdots = r_n = 0$ .

By a  $\mathcal{N}$ -marked connection, we mean the data  $({}_{\mathcal{O}}\mathcal{M}, \nabla, \text{iso})$  of a meromorphic connection  $(\mathcal{M}, \nabla)$  around 0 endowed with an isomorphism of formal connections at the origin  $\text{iso} : \mathcal{M}_{\hat{0}} \rightarrow \mathcal{N}_{\hat{0}}$ . As in dimension 1, Mochizuki [21, 22] showed that  $\mathcal{N}$ -marked connections are determined by their associated Stokes torsor, so we consider them as elements in  $H^1(T, \text{St}_{\mathcal{N}}(C))$ .

Since  $\text{St}_{\mathcal{N}}$  is a sheaf of complex algebraic groups, its sheaf of  $R$ -points  $\text{St}_{\mathcal{N}}(R)$  is a well-defined sheaf of groups on  $T$  for any commutative  $C$ -algebra  $R$ . Consequently, one can consider the functor of relative Stokes torsors  $R \rightarrow H^1(T, \text{St}_{\mathcal{N}}(R))$ , denoted by  $H^1(T, \text{St}_{\mathcal{N}})$ . Following a strategy designed by Deligne, Babbitt and Varadarajan [2] proved that  $H^1(S^1, \text{St}_{\mathcal{N}})$  is representable by an affine space. Hence in dimension 1, the set of torsors under  $\text{St}_{\mathcal{N}}$  has a structure of a complex algebraic variety.

The interest of this result is to *provide a framework in which questions related to differential equations can be treated with the apparatus of algebraic geometry*. This might look like a wish rather than a documented fact since the local theory of linear differential equations is fully understood in dimension 1 by means of analysis. In dimension  $\geq 2$  however, new phenomena appear and this geometric perspective seems relevant. As we will show, the representability of  $H^1(T, \text{St}_{\mathcal{N}})$  in any dimension implies for differential equations quite strong finiteness

results which have no counterparts in dimension 1 and which seem out of reach with former technology. Thus, we prove the following:

**THEOREM 1.** – *The functor  $H^1(T, \text{St}_{\mathcal{N}})$  is representable by an affine scheme of finite type over  $C$ .*

Before explaining how the proof relates to Deligne’s skeleton conjecture, let us describe two applications to finiteness results.

Suppose that  $\mathcal{N}$  is *very good*, that is, for functions  $a_i, a_j$  appearing in (0.0.1) with  $a_i \neq a_j$ , the difference  $a_i - a_j$  has poles along all the components of the divisor  $D$  along which  $\mathcal{N}$  is localized. Let  $V$  be a manifold containing 0 and let us denote by  $\mathcal{N}_V$  the restriction of the connection  $\mathcal{N}$  to  $V$ . We prove the following:

**THEOREM 2.** – *If  $V$  is transverse to every irreducible component of  $D$ , there is only a finite number of equivalence classes of  $\mathcal{N}$ -marked connections with given restriction to  $V$ . Furthermore, this number depends only on  $\mathcal{N}$  and on  $V$ .*

This theorem looks like a weak differential version of Lefschetz’s theorem. A differential Lefschetz theorem would assert that for a generic choice of  $V$ ,  $\mathcal{N}$ -marked connections are determined by their restriction to  $V$ . It is a hope of the author that such a question is approachable by geometric means using the morphism of schemes

$$(0.0.2) \quad \text{res}_V : H^1(T, \text{St}_{\mathcal{N}}) \longrightarrow H^1(T', \text{St}_{\mathcal{N}_V})$$

induced by the restriction to  $V$ .

To give flesh to this intuition, let us indicate how geometry enters the proof of Theorem 2. Since unramified morphisms of finite type have finite fibers, it is enough to show that  $\mathcal{N}$ -marked connections lie in the unramified locus of  $\text{res}_V$ , which is the locus where the tangent map of  $\text{res}_V$  is injective. We show in 5.2 a canonical identification

$$(0.0.3) \quad T_{(\mathcal{M}, \nabla, \text{iso})} H^1(T, \text{St}_{\mathcal{N}}) \simeq \mathcal{H}^1(\text{Sol End } \mathcal{M})_0,$$

where the left-hand side denotes the tangent space of  $H^1(T, \text{St}_{\mathcal{N}})$  at  $(\mathcal{M}, \nabla, \text{iso})$  and where  $\mathcal{H}^1 \text{Sol}$  denotes the first cohomology sheaf of the solution complex of a  $\mathcal{D}$ -module. Note that the left-hand side of (0.0.3) is *algebraic*, whereas the right-hand side is *transcendental*. From (0.0.3) we deduce a similar transcendental interpretation for the kernel  $\text{Ker } T_{(\mathcal{M}, \nabla, \text{iso})} \text{res}_V$  and prove its vanishing using a perversity theorem due to Mebkhout [18].

Using an invariance theorem due to Sabbah [24], we further prove the following rigidity result:

**THEOREM 3.** – *Suppose that  $D$  has at least two components and that  $\mathcal{N}$  is very general. Then there is only a finite number of equivalence classes of  $\mathcal{N}$ -marked connections.*

In this statement, very general means that  $\mathcal{N}$  is very good and that the residues of each regular constituent contributing to  $\mathcal{N}$  in (0.0.1) lie away from a countable union of strict Zariski closed subsets of the affine space.

Let us finally explain roughly the proof of Theorem 1. The main idea is to import and prove a conjecture from the field of Galois representations. Let  $X$  be a smooth variety over a finite field of characteristic  $p > 0$ , and let  $\ell \neq p$  be a prime number. To any  $\ell$ -adic local system  $\mathcal{F}$  on  $X$  up to semi-simplification, one can associate its *skeleton*  $\text{sk } \mathcal{F}$ , that is