

quatrième série - tome 49 fascicule 2 mars-avril 2016

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Hélène ESNAULT & Olivier WITTENBERG

On the cycle class map for zero-cycles over local fields

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2016

N. ANANTHARAMAN I. GALLAGHER
P. BERNARD B. KLEINER
E. BREUILLARD E. KOWALSKI
R. CERF M. MUSTAȚĂ
A. CHAMBERT-LOIR L. SALOFF-COSTE

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition / *Publication*

Société Mathématique de France
Institut Henri Poincaré
11, rue Pierre et Marie Curie
75231 Paris Cedex 05
Tél. : (33) 01 44 27 67 99
Fax : (33) 01 40 46 90 96

Abonnements / *Subscriptions*

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 09
Fax : (33) 04 91 41 17 51
email : smf@smf.univ-mrs.fr

Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

© 2016 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ON THE CYCLE CLASS MAP FOR ZERO-CYCLES OVER LOCAL FIELDS

BY HÉLÈNE ESNAULT AND OLIVIER WITTENBERG
WITH AN APPENDIX BY SPENCER BLOCH

ABSTRACT. – We study the Chow group of 0-cycles of smooth projective varieties over local and strictly local fields. We prove in particular the injectivity of the cycle class map to integral ℓ -adic cohomology for a large class of surfaces with positive geometric genus, over local fields of residue characteristic $\neq \ell$. The same statement holds for semistable $K3$ surfaces defined over $\mathbf{C}((t))$, but does not hold in general for surfaces over strictly local fields.

RÉSUMÉ. – Nous étudions le groupe de Chow des 0-cycles des variétés projectives et lisses sur les corps locaux et strictement locaux. Nous prouvons en particulier l'injectivité de l'application classe de cycle vers la cohomologie ℓ -adique entière pour de nombreuses surfaces de genre géométrique non nul, sur les corps locaux de caractéristique résiduelle $\neq \ell$. Le même énoncé vaut pour les surfaces $K3$ semi-stables définies sur $\mathbf{C}((t))$, mais ne vaut pas en général pour les surfaces sur les corps strictement locaux.

1. Introduction

Let X be a smooth projective variety over a field K , let $\mathrm{CH}_0(X)$ denote the Chow group of 0-cycles on X up to rational equivalence and let $A_0(X) \subset \mathrm{CH}_0(X)$ be the subgroup of cycle classes of degree 0.

When K is algebraically closed, the group $A_0(X)$ is divisible and its structure as an abelian group is, conjecturally, rather well understood, thanks to Roitman's theorem and to the Bloch-Beilinson-Murre conjectures. A central tool for the study of $A_0(X)$ over other types of fields is the cycle class map

$$\psi : \mathrm{CH}_0(X)/n\mathrm{CH}_0(X) \rightarrow H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d))$$

to étale cohomology, where n denotes an integer invertible in K and $d = \dim(X)$. The group $H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d))$ is easier to understand, thanks to the Hochschild-Serre spectral

The first author is supported by the the ERC Advanced Grant 226257, the Chaire d'excellence 2011 of the Fondation Sciences Mathématiques de Paris and the Einstein Foundation.

sequence; for instance, if K has cohomological dimension ≤ 1 and X is simply connected, then $H_{\text{ét}}^{2d}(X, \mathbf{Z}/n\mathbf{Z}(d)) = \mathbf{Z}/n\mathbf{Z}$ and ψ may be interpreted as the degree map.

According to one of the main results of higher-dimensional unramified class field theory, due to Kato and Saito [43], if K is a finite field, the group $A_0(X)$ is finite and ψ is an isomorphism. More recently, Saito and Sato [69] have shown that if K is the quotient field of an excellent Henselian discrete valuation ring with finite or separably closed residue field, the group $A_0(X)$ is the direct sum of a finite group of order prime to p and a group divisible by all integers prime to p (see also [11, Théorème 3.25]). In this case, however, the map ψ need not be either injective or surjective. What Saito and Sato prove, instead, is the bijectivity of the analogous cycle class map for cycles of dimension 1 on regular models of X over the ring of integers of K (see [69, Theorem 1.16]).

Following a method initiated by Bloch [4], one may approach the torsion subgroup of $A_0(X)$, as well as the kernel of ψ , with the help of algebraic K-theory, when X is a surface. We refer to [10] for a detailed account of this circle of ideas. Strong results were obtained in this way for rational surfaces, and more generally for surfaces with geometric genus zero, over number fields, p -adic fields, and fields of characteristic 0 and cohomological dimension 1 (see [5], [6], [16], [9], [14], [68]). We note that over algebraically closed fields of characteristic 0, surfaces with geometric genus zero are those surfaces for which the Chow group $A_0(X)$ should be representable, according to Bloch's conjecture (see [5, § 1]).

The first theorem of this paper establishes the injectivity of ψ for a large class of surfaces over local fields, when n is divisible enough and prime to the residue characteristic, without any assumption on the geometric genus. In principle, this theorem should be applicable to all simply connected surfaces, a generality in which the injectivity of ψ may not have been expected. Before we state it, we set up some notation. Let \mathcal{X} be a regular proper flat scheme over an excellent Henselian discrete valuation ring R . Let $X = \mathcal{X} \otimes_R K$ and $A = \mathcal{X} \otimes_R k$ denote the generic fiber and the special fiber, respectively. We assume the reduced special fiber A_{red} has simple normal crossings, and write $\text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell = \varprojlim \text{CH}_0(X)/\ell^n \text{CH}_0(X)$.

THEOREM 1 (Theorem 3.1 and Remark 3.2). – *Assume the residue field k is finite and X is a surface whose Albanese variety has potentially good reduction. If the irreducible components of A satisfy the Tate conjecture, then for any ℓ invertible in k , the cycle class map*

$$(1.1) \quad \text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_{\text{ét}}^4(X, \mathbf{Z}_\ell(2))$$

is injective. Equivalently, the natural pairing $\text{CH}_0(X) \times \text{Br}(X) \rightarrow \mathbf{Q}/\mathbf{Z}$ is non-degenerate on the left modulo the maximal ℓ -divisible subgroup of $\text{CH}_0(X)$.

The assumption on the irreducible components of A holds as soon as X has geometric genus zero, as well as in many examples of nontrivial degenerations of surfaces with nonzero geometric genus (see § 3). Theorem 1 is due to Saito [68] when X is a surface with geometric genus zero over a p -adic field. An example of Parimala and Suresh [62] shows that the assumption on the Albanese variety cannot be removed. Finally, we note that Theorem 1 may be viewed as a higher-dimensional generalization of Lichtenbaum-Tate duality for curves, according to which the natural pairing $\text{CH}_0(X) \times \text{Br}(X) \rightarrow \mathbf{Q}/\mathbf{Z}$ is non-degenerate if X is a smooth proper curve over a p -adic field (see [54]).

Our starting point for the proof of Theorem 1 is the theorem of Saito and Sato alluded to above about the cycle class map for 1-cycles on \mathcal{X} [69, Theorem 1.16], which allows us to express the kernel of ψ purely in terms of the scheme A and of the cohomology of X , when k is either finite or separably closed (Theorem 2.1 and Theorem 2.2). The dimension of X plays no role in this part of the argument; an application to the study of 0-cycles on a cubic threefold over \mathbf{Q}_p may be found in Example 2.12. Theorem 1 is then obtained by analyzing the various cohomology groups which appear in the resulting expression for $\text{Ker}(\psi)$. More precisely, in the situation of Theorem 1, we prove the stronger assertion that the 1-dimensional cycle class map $\psi_{1,A} : \text{CH}_1(A) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_A^4(\mathcal{X}, \mathbf{Z}_\ell(2))$ to integral ℓ -adic étale homology is surjective. This provides a geometric explanation for the assumption that the Albanese variety of X have potentially good reduction, a condition which first appeared in [68] and which turns out to be essential for the surjectivity of $\psi_{1,A}$ to hold (see Lemma 3.7 and § 4.3).

When the residue field k is separably closed instead of finite, the arguments used in the proof of Theorem 1 fail in several places. They still lead to the following statement, which may also be deduced from results of Colliot-Thélène and Raskind [13] (see § 4 for comments on this point).

THEOREM 2 (Theorem 4.1 and Remark 4.3). – *Assume k is separably closed and K has characteristic 0. If X is a surface with geometric genus zero, then for any ℓ invertible in k , the cycle class map*

$$(1.2) \quad \text{CH}_0(X) \widehat{\otimes} \mathbf{Z}_\ell \rightarrow H_{\text{ét}}^4(X, \mathbf{Z}_\ell(2))$$

is injective. If in addition X is simply connected, then $A_0(X)$ is divisible by ℓ and the unramified cohomology group $H_{\text{nr}}^3(X, \mathbf{Q}_\ell/\mathbf{Z}_\ell(2))$ vanishes.

This leaves open the question of the injectivity of the cycle class map (1.2) when k is separably closed and X is a surface with positive geometric genus over K . In this situation, the 1-dimensional cycle class map $\psi_{1,A}$ is far from being surjective. Building on the work of Kulikov, Persson, Pinkham [51] [64], and of Miranda and Morrison [59], we nevertheless give a positive answer for semistable K3 surfaces over $\mathbf{C}((t))$.

THEOREM 3 (Theorem 5.1). – *Let X be a K3 surface over $\mathbf{C}((t))$. If X has semistable reduction, the group $A_0(X)$ is divisible.*

The proof of Theorem 3 hinges on the precise knowledge of the combinatorial structure of a degeneration of X . It would go through over the maximal unramified extension of a p -adic field, as far as prime-to- p divisibility is concerned, if similar knowledge were available. This is in marked contrast with the situation over p -adic fields, where such knowledge is not necessary for the proof of Theorem 1.

In the final section of this paper, with the help of Ogg-Shafarevich theory and of a construction due to Persson, we show that the hope for a statement analogous to Theorem 1 over the quotient field of a strictly Henselian excellent discrete valuation ring is in fact too optimistic, even over the maximal unramified extension of a p -adic field.