

Astérisque

MOCHIZUKI TAKURO

**Kobayashi-Hitchin correspondence for tame
harmonic bundles and an application**

Astérisque, tome 309 (2006)

[<http://www.numdam.org/item?id=AST_2006__309__R1_0>](http://www.numdam.org/item?id=AST_2006__309__R1_0)

© Société mathématique de France, 2006, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

**KOBAYASHI-HITCHIN
CORRESPONDENCE FOR
TAME HARMONIC BUNDLES
AND AN APPLICATION**

Takuro Mochizuki

T. Mochizuki

Department of Mathematics, Kyoto University, Kyoto 606-8502, Japan.

E-mail : takuro@math.kyoto-u.ac.jp

2000 Mathematics Subject Classification. — 14J60, 53C07.

Key words and phrases. — Higgs bundle, harmonic bundle, Kobayashi-Hitchin correspondence, Hermitian-Einstein metric, Bogomolov-Gieseker inequality, flat bundle, variation of polarized Hodge structure, quasiprojective variety.

This work was prepared with the partial support of Ministry of Education, Culture, Sports, Science and Technology.

To my parents

KOBAYASHI-HITCHIN CORRESPONDENCE FOR TAME HARMONIC BUNDLES AND AN APPLICATION

Takuro Mochizuki

Abstract. — We establish the correspondence between tame harmonic bundles and μ_L -polystable parabolic Higgs bundles with trivial characteristic numbers. We also show the Bogomolov-Gieseker type inequality for μ_L -stable parabolic Higgs bundles.

Then we show that any local system on a smooth quasiprojective variety can be deformed to a variation of polarized Hodge structure. As a consequence, we can conclude that some kind of discrete groups cannot be a split quotient of the fundamental group of a smooth quasiprojective variety.

Résumé (La correspondance de Kobayashi-Hitchin pour les fibrés harmoniques modérés et une application)

Nous établissons la correspondance de Kobayashi-Hitchin entre les fibrés harmoniques modérés et fibrés de Higgs paraboliques μ_L -polystables dont les deux premiers nombres de Chern sont nuls. Ensuite, nous montrons que tout système local sur une variété quasi-projective lisse peut être déformé vers une variation de structure de Hodge polarisée. En conséquence, nous pouvons conclure que certains groupes discrets ne peuvent pas apparaître comme quotient scindé d'un groupe fondamental d'une variété quasi-projective lisse.

CONTENTS

1. Introduction	1
1.1. Background	1
1.2. Main Purpose	2
1.3. Additional Results	6
1.4. Outline	8
1.5. Acknowledgement	9
2. Preliminary	11
2.1. Notation and Words	11
2.2. Review of some Results of Simpson on Kobayashi-Hitchin Correspondence	12
2.3. Weitzenböck Formula	15
2.4. A Priori Estimate of Higgs Fields	16
2.5. Norm Estimate for Tame Harmonic Bundle in Two Dimensional Case	19
2.6. Preliminary from Elementary Calculus	20
2.7. Reflexive Sheaf	22
2.8. Moduli Spaces of Representations	23
3. Parabolic Higgs Bundle and Regular Filtered Higgs Bundle	25
3.1. Parabolic Higgs Bundle	25
3.2. Filtered Sheaf	31
3.3. Perturbation of Parabolic Structure	34
3.4. Mehta-Ramanathan Type Theorem	37
3.5. Adapted Metric and the Associated Parabolic Flat Higgs Bundle	42
3.6. Convergence	42
4. An Ordinary Metric for a Parabolic Higgs Bundle	45
4.1. Around the Intersection $D_i \cap D_j$	45
4.2. Around a Smooth Point of the Divisor	47
4.3. Global Ordinary Metric	53

5. Parabolic Higgs Bundle Associated to Tame Harmonic Bundle	59
5.1. Polystability and Uniqueness	59
5.2. Vanishing of Characteristic Numbers	60
6. Preliminary Correspondence and Bogomolov-Gieseker Inequality ..	65
6.1. Graded Semisimple Parabolic Higgs Bundles on Surface	65
6.2. Bogomolov-Gieseker Inequality	67
7. Construction of a Frame	69
7.1. A Priori Estimate of Higgs Field on a Punctured Disc	70
7.2. Construction of Local Holomorphic Frames	73
8. Some Convergence Results	79
8.1. Convergence of a Sequence of Tame Harmonic Bundles	79
8.2. Preparation for the Proof of Theorem 9.1	82
9. Existence of Adapted Pluri-Harmonic Metric	87
9.1. The Surface Case	87
9.2. The Higher Dimensional Case	92
10. Torus Action and the Deformation of Representations	95
10.1. Torus Action on the Moduli Space of Representations	95
10.2. Monodromy Group	99
G-Harmonic Bundle	103
A.1. G -Principal Bundles with Flat Structure or Holomorphic Structure	103
A.2. Definitions	105
A.3. Semisimplicity and Pluri-Harmonic Reduction	108
Bibliography	113

CHAPTER 1

INTRODUCTION

1.1. Background

1.1.1. Kobayashi-Hitchin correspondence. — We briefly recall some aspects of the so-called Kobayashi-Hitchin correspondence. (See the introduction of [38] for more detail.) In 1960's, M. S. Narasimhan and C. S. Seshadri proved the correspondence between irreducible flat unitary bundles and stable vector bundles with degree 0, on a compact Riemann surface ([47]). Clearly, it was desired to extend their result to the higher dimensional case and the non-flat case.

In early 1980's, S. Kobayashi introduced the Hermitian-Einstein condition for holomorphic bundles on Kahler manifolds ([30], [31]). He and M. Lübke ([37]) proved that the existence of Hermitian-Einstein metric implies the polystability of the underlying holomorphic bundle. S. K. Donaldson pioneered the way for the inverse problem ([12] and [13]). He attributed the problem to Kobayashi and N. Hitchin. The definitive result was given by K. Uhlenbeck, S. T. Yau and Donaldson ([64] and [14]). We also remark that V. Mehta and A. Ramanathan ([40]) proved the correspondence in the case where the Chern class is trivial, i.e., the correspondence of flat unitary bundles and stable vector bundles with trivial Chern classes.

On the other hand, it was quite fruitful to consider the correspondences for vector bundles with some additional structures like Higgs fields, which was initiated by Hitchin ([22]). He studied the Higgs bundles on a compact Riemann surface and the moduli spaces. His work has influenced various fields of mathematics. It involves a lot of subjects and ideas, and one of his results is the correspondence of the stability and the existence of Hermitian-Einstein metrics for Higgs bundles on a compact Riemann surface.

1.1.2. A part of C. Simpson's work. — C. Simpson studied the Higgs bundles over higher dimensional complex manifolds, influenced by the work of Hitchin, but motivated by his own subject: Variation of Polarized Hodge Structure. He made great