

*quatrième série - tome 53    fascicule 5    septembre-octobre 2020*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Chen JIANG

*Boundedness of  $\mathbb{Q}$ -Fano varieties with degrees and alpha-invariants  
bounded from below*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

**Responsable du comité de rédaction / *Editor-in-chief***

Patrick BERNARD

**Publication fondée en 1864 par Louis Pasteur**

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

**Comité de rédaction au 1<sup>er</sup> janvier 2020**

P. BERNARD      D. HARARI  
S. BOUCKSOM      A. NEVES  
G. CHENEVIER      J. SZEFTEL  
Y. DE CORNULIER      S. VŨ NGỌC  
A. DUCROS      A. WIENHARD  
G. GIACOMIN      G. WILLIAMSON

**Rédaction / *Editor***

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annales@ens.fr](mailto:annales@ens.fr)

---

**Édition et abonnements / *Publication and subscriptions***

Société Mathématique de France  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Tél. : (33) 04 91 26 74 64  
Fax : (33) 04 91 41 17 51  
email : [abonnements@smf.emath.fr](mailto:abonnements@smf.emath.fr)

**Tarifs**

Abonnement électronique : 428 euros.  
Abonnement avec supplément papier :  
Europe : 576 €. Hors Europe : 657 € (\$ 985). Vente au numéro : 77 €.

---

© 2020 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593 (print) 1873-2151 (electronic)

Directeur de la publication : Fabien Durand  
Périodicité : 6 n<sup>os</sup> / an

# BOUNDEDNESS OF $\mathbb{Q}$ -FANO VARIETIES WITH DEGREES AND ALPHA-INVARIANTS BOUNDED FROM BELOW

BY CHEN JIANG

---

**ABSTRACT.** – We show that  $\mathbb{Q}$ -Fano varieties of fixed dimension with anti-canonical degrees and alpha-invariants bounded from below form a bounded family. As a corollary,  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties of fixed dimension with anti-canonical degrees bounded from below form a bounded family.

**RÉSUMÉ.** – Nous démontrons que les variétés de  $\mathbb{Q}$ -Fano de dimension fixe dont les degrés anticanoniques et les alpha-invariants sont bornés inférieurement forment une famille bornée. En corollaire, les variétés de  $\mathbb{Q}$ -Fano  $\mathbb{K}$ -semistables de dimension fixe dont les degrés anticanoniques sont bornés inférieurement forment une famille bornée.

## 1. Introduction

Throughout the article, we work over an algebraically closed field of characteristic zero. A  $\mathbb{Q}$ -Fano variety is defined to be a normal projective variety  $X$  with at most klt singularities such that the anti-canonical divisor  $-K_X$  is an ample  $\mathbb{Q}$ -Cartier divisor.

When the base field is the complex number field, an interesting problem for  $\mathbb{Q}$ -Fano varieties is the existence of Kähler-Einstein metrics which is related to  $\mathbb{K}$ -(semi)stability of  $\mathbb{Q}$ -Fano varieties. It has been known that a Fano manifold  $X$  (i.e., a smooth  $\mathbb{Q}$ -Fano variety over  $\mathbb{C}$ ) admits Kähler-Einstein metrics if and only if  $X$  is  $K$ -polystable by the works [15, 42, 16, 17, 14, 40, 32, 33, 3] and [11, 12, 13, 43].  $\mathbb{K}$ -stability is stronger than  $\mathbb{K}$ -polystability, and  $\mathbb{K}$ -polystability is stronger than  $\mathbb{K}$ -semistability. Hence  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties are interesting for both differential geometers and algebraic geometers.

It also turned out that Kähler-Einstein metrics and  $\mathbb{K}$ -stability play crucial roles for the construction of nice moduli spaces of certain  $\mathbb{Q}$ -Fano varieties. For example, compact moduli spaces of smoothable Kähler-Einstein  $\mathbb{Q}$ -Fano varieties have been constructed (see [36] for dimension two case and [30, 39, 34] for higher dimensional case). In order to consider the

---

The author was supported by JSPS KAKENHI Grant Number JP16K17558 and World Premier International Research Center Initiative (WPI), MEXT, Japan.

moduli space of certain (singular)  $\mathbb{Q}$ -Fano varieties, the first step is to show the boundedness property, which is the motivation of this paper. We show the boundedness of  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties of fixed dimension with anti-canonical degrees bounded from below, which gives an affirmative answer to a question asked by Yuchen Liu during the AIM workshop “Stability and moduli spaces” in January 2017.

**THEOREM 1.1.** – *Fix a positive integer  $d$  and a real number  $\delta > 0$ . Then the set of  $d$ -dimensional  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties  $X$  with  $(-K_X)^d > \delta$  forms a bounded family.*

Note that the assumption that  $(-K_X)^d$  is bounded from below is necessary, by Example 1.4(2) later.

As mentioned before, one might have further applications of Theorem 1.1 such as constructing moduli spaces of  $d$ -dimensional  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties with bounded anti-canonical degrees. An interesting corollary of Theorem 1.1 is the discreteness of the anti-canonical degrees of  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties.

**COROLLARY 1.2.** – *Fix a positive integer  $d$ . Then the set of  $(-K_X)^d$  for  $d$ -dimensional  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties  $X$  is finite away from 0.*

Here a set  $\mathcal{P}$  of positive real numbers is *finite away from 0* if for any  $\delta > 0$ ,  $\mathcal{P} \cap (\delta, \infty)$  is a finite set. We remark that Corollary 1.2 might be related to the conjectural discreteness of minimal normalized volumes of klt singularities, cf. [31, Question 4.3].

The idea of proof of Theorem 1.1 comes from birational geometry. According to Minimal Model Program,  $\mathbb{Q}$ -Fano varieties form a fundamental class in birational geometry, and the boundedness property for  $\mathbb{Q}$ -Fano varieties is also interesting from the point view of birational geometry. For example, Kollár, Miyaoka, and Mori [26] proved that smooth Fano varieties form a bounded family. The most celebrated progress recently is the proof of Borisov-Alexeev-Borisov Conjecture due to Birkar [4, 5], which says that given a positive integer  $d$  and a real number  $\epsilon > 0$ , the set of  $\epsilon$ -lc  $\mathbb{Q}$ -Fano varieties of dimension  $d$  forms a bounded family.

In this paper, inspired by Birkar’s work, in order to show Theorem 1.1, we show the following theorem.

**THEOREM 1.3.** – *Fix a positive integer  $d$  and a real number  $\delta > 0$ . Then the set of  $d$ -dimensional  $\mathbb{Q}$ -Fano varieties  $X$  with  $(-K_X)^d > \delta$  and  $\alpha(X) > \delta$  forms a bounded family.*

Here  $\alpha(X)$  is the *alpha-invariant* of  $X$  defined by Tian [41] (see also [7]) in order to investigate the existence of Kähler-Einstein metrics on Fano manifolds. Recall that Fujita and Odaka [18, Theorem 3.5] proved that the alpha-invariant of a  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano variety of dimension  $d$  is always not less than  $1/(d+1)$ , so Theorem 1.3 implies Theorem 1.1 naturally. The advantage to consider Theorem 1.3 is that we can then apply methods from birational geometry, instead of dealing with  $\mathbb{K}$ -semistable  $\mathbb{Q}$ -Fano varieties.

The point of Theorem 1.3 is that we replace the  $\epsilon$ -lc condition in Borisov-Alexeev-Borisov Conjecture by the condition on lower bound of anti-canonical degrees and alpha-invariants, which are global invariants.

We remark that if one takes  $\delta = 1$ , then Theorem 1.3 is a consequence of [4, Theorem 1.3], which says that the set of *exceptional*  $\mathbb{Q}$ -Fano varieties (i.e.,  $\mathbb{Q}$ -Fano varieties  $X$  with  $\alpha(X) > 1$ ) of fixed dimension forms a bounded family. Note that in this case we do not even need to

assume  $(-K_X)^d$  is bounded from below. But in general we need to assume both  $(-K_X)^d$  and  $\alpha(X)$  are bounded from below, by the following examples.

EXAMPLE 1.4. – Fix a positive integer  $d$ .

1. Consider the weighted projective space  $X_n = \mathbb{P}(1^d, n)$  which is a  $\mathbb{Q}$ -Fano variety of dimension  $d$  with  $(-K_{X_n})^d = (n+d)^d/n > 1$ , but it is clear that  $\{X_n\}$  does not form a bounded family.
2. Consider  $Y_{8n+4} \subset \mathbb{P}(2, 2n+1, 2n+1, 4n+1)$ , a general weighted hypersurface of degree  $8n+4$ , which is a  $\mathbb{Q}$ -Fano variety of dimension 2 with  $\alpha(Y_{8n+4}) = 1$  (see [6, Corollary 1.12] or [22]), but it is clear that  $\{Y_{8n+4}\}$  does not form a bounded family. For more interesting examples of  $\mathbb{Q}$ -Fano varieties with  $\alpha \geq 1$ , we refer to [6, 9] in dimension 2 and [8, 10] in higher dimensions. Note that all examples with  $\alpha \geq 1$  are  $\mathbb{K}$ -semistable (in fact,  $\mathbb{K}$ -stable) by [35, Theorem 1.4] (or [41]).

By [4, Proposition 7.13] or [5, Theorem 2.15], Theorem 1.3 is a consequence of the following theorem.

THEOREM 1.5. – *Fix a positive integer  $d$  and a real number  $\delta > 0$ . Then there exists a positive integer  $m$  depending only on  $d$  and  $\delta$  such that if  $X$  is a  $d$ -dimensional  $\mathbb{Q}$ -Fano variety with  $(-K_X)^d > \delta$  and  $\alpha(X) > \delta$ , then  $| -mK_X |$  defines a birational map.*

To show Theorem 1.5, our main idea is to establish an inequality expressed in terms of the volume of  $-K_X|_G$  on a covering family of subvarieties  $G$  of  $X$  and  $(-K_X)^d, \alpha(X)$ , see Lemma 3.1.

As a variation of Theorem 1.3, we can also show the following theorem.

THEOREM 1.6. – *Fix a positive integer  $d$  and a real number  $\theta > 0$ . Then the set of  $d$ -dimensional  $\mathbb{Q}$ -Fano varieties  $X$  with  $\alpha(X)^d \cdot (-K_X)^d > \theta$  forms a bounded family.*

Logically, Theorem 1.3 is implied by Theorem 1.6. But we will show Theorem 1.3 first in order to make the explanation more clear.

REMARK 1.7. – Note that the invariant  $\alpha(X)^d \cdot (-K_X)^d$  appears naturally in birational geometry, see for example [25, Theorem 6.7.1]. It is not clear whether we can replace  $\alpha(X)^d \cdot (-K_X)^d$  in Theorem 1.6 by  $\alpha(X)^{d'} \cdot (-K_X)^d$  for some positive real number  $d' < d$ . At least  $d' \leq d-1$  is not sufficient to conclude the boundedness. For example, in Example 1.4(1),  $(-K_{X_n})^d = (n+d)^d/n$  and  $\alpha(X_n) = 1/(n+d)$  (for computation of alpha-invariants of toric varieties, see [1, 6.3]), hence  $\alpha(X_n)^{d-1} \cdot (-K_{X_n})^d > 1$ .

REMARK 1.8. – We remark that the proof of both Theorems 1.3 and 1.6 works under the weaker assumption that  $X$  is a *weak  $\mathbb{Q}$ -Fano variety* (i.e.,  $X$  has at most klt singularities and  $-K_X$  is nef and big), see also Remark 2.5. But it is not clear yet whether the log Fano pair versions hold or not.