

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

ON THE MOTION OF A SMALL BODY IMMERSED IN A TWO-DIMENSIONAL INCOMPRESSIBLE PERFECT FLUID

Olivier Glass & Christophe Lacave & Franck Sueur

Tome 142

Fascicule 3

2014

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 489-536

Le *Bulletin de la Société Mathématique de France* est un
périodique trimestriel de la Société Mathématique de France.

Fascicule 3, tome 142, septembre 2014

Comité de rédaction

Jean BARGE	Daniel HUYBRECHTS
Gérard BESSON	Yves LE JAN
Emmanuel BREUILLARD	Julien MARCHÉ
Antoine CHAMBERT-LOIR	Laure SAINT-RAYMOND
Jean-François DAT	Wilhelm SCHLAG
Charles FAVRE	
Raphaël KRIKORIAN (dir.)	

Diffusion

Maison de la SMF	Hindustan Book Agency	AMS
Case 916 - Luminy	O-131, The Shopping Mall	P.O. Box 6248
13288 Marseille Cedex 9	Arjun Marg, DLF Phase 1	Providence RI 02940
France	Gurgaon 122002, Haryana	USA
smf@smf.univ-mrs.fr	Inde	www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement Europe : 300 €, hors Europe : 334 € (\$ 519)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat : Nathalie Christiaën

Bulletin de la Société Mathématique de France

Société Mathématique de France

Institut Henri Poincaré, 11, rue Pierre et Marie Curie

75231 Paris Cedex 05, France

Tél : (33) 01 44 27 67 99 • Fax : (33) 01 40 46 90 96

revues@smf.ens.fr • <http://smf.emath.fr/>

© *Société Mathématique de France* 2014

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484

Directeur de la publication : Marc PEIGNÉ

ON THE MOTION OF A SMALL BODY IMMERSSED IN A TWO-DIMENSIONAL INCOMPRESSIBLE PERFECT FLUID

BY OLIVIER GLASS, CHRISTOPHE LACAVE & FRANCK SUEUR

ABSTRACT. — In this paper we prove that the motion of a solid body in a two dimensional incompressible perfect fluid converges, when the body shrinks to a point with fixed mass and circulation, to a variant of the vortex-wave system where the vortex, placed in the point occupied by the shrunk body, is accelerated by a lift force similar to the Kutta-Joukowski force of the irrotational theory.

RÉSUMÉ (*Sur le mouvement d'un petit corps solide immergé dans un fluide parfait incompressible en deux dimensions*)

Dans cet article nous prouvons que le mouvement d'un petit corps solide immergé dans un fluide parfait incompressible en deux dimensions converge, quand le corps se rétrécit en un point, avec sa masse et sa circulation fixées, vers une variante du système « Euler+point vortex » où le vortex, placé au point où le solide a rétréci, est accéléré par un terme de force de portance similaire à la force de Kutta-Joukowski de la théorie irrotationnelle.

Texte reçu le 16 janvier 2012 et accepté le 30 mars 2012.

OLIVIER GLASS, Ceremade, Université Paris-Dauphine, Place du Maréchal de Lattre de Tassigny, 75775 Paris Cedex 16, France

CHRISTOPHE LACAVE, Univ Paris Diderot, Sorbonne Paris Cité, Institut de Mathématiques de Jussieu-Paris Rive Gauche, UMR 7586, CNRS, Sorbonne Universités, UPMC Univ Paris 06, 75013, Paris, France.

FRANCK SUEUR, Franck Sueur, Institut de Mathématiques de Bordeaux, UMR CNRS 5251, Université de Bordeaux, 351 cours de la Libération, 33405 Talence Cedex, France.

2010 Mathematics Subject Classification. — 76B99, 70E99.

Key words and phrases. — Fluid-solid interactions, incompressible perfect fluid, vortex-wave system, Kutta-Joukowski force.

1. Introduction

In this paper we consider the motion of a small solid body in a planar ideal fluid, and the limit behaviour of the system as the solid body is reduced to a point.

Let us first describe the equations when the solid has a fixed size. Let \mathcal{S}_0 be a closed, bounded, connected and simply connected subset of the plane with smooth boundary. We assume that the body initially occupies the domain \mathcal{S}_0 and rigidly moves so that at time t it occupies an isometric domain denoted by $\mathcal{S}(t)$. We denote $\mathcal{F}(t) := \mathbb{R}^2 \setminus \mathcal{S}(t)$ the domain occupied by the fluid at time t starting from the initial domain $\mathcal{F}_0 := \mathbb{R}^2 \setminus \mathcal{S}_0$.

The equations modelling the dynamics of the system then read

$$\begin{aligned}
 (1) \quad & \frac{\partial u}{\partial t} + (u \cdot \nabla)u + \nabla p = 0 \text{ for } x \in \mathcal{F}(t), \\
 (2) \quad & \operatorname{div} u = 0 \text{ for } x \in \mathcal{F}(t), \\
 (3) \quad & u \cdot n = u_{\mathcal{S}} \cdot n \text{ for } x \in \partial \mathcal{S}(t), \\
 (4) \quad & \lim_{|x| \rightarrow \infty} |u| = 0, \\
 (5) \quad & mh''(t) = \int_{\partial \mathcal{S}(t)} pn \, ds, \\
 (6) \quad & \mathcal{J}r'(t) = \int_{\partial \mathcal{S}(t)} (x - h(t))^\perp \cdot pn \, ds, \\
 (7) \quad & u|_{t=0} = u_0 \text{ for } x \in \mathcal{F}_0, \\
 (8) \quad & h(0) = h_0, \quad h'(0) = \ell_0, \quad r(0) = r_0.
 \end{aligned}$$

Here $u = (u_1, u_2)$ and p denote the velocity and pressure fields, m and \mathcal{J} denote respectively the mass and the moment of inertia of the body while the fluid is supposed to be homogeneous of density 1, to simplify the notations. When $x = (x_1, x_2)$ the notation x^\perp stands for $x^\perp = (-x_2, x_1)$, n denotes the unit normal vector pointing outside the fluid, $h'(t)$ is the velocity of the center of mass $h(t)$ of the body and $r(t)$ denotes the angular velocity of the rigid body. Finally we denote by $u_{\mathcal{S}}$ the velocity of the body:

$$(9) \quad u_{\mathcal{S}}(t, x) = h'(t) + r(t)(x - h(t))^\perp.$$

Since $\mathcal{S}(t)$ is the position occupied by a rigid body there exists a rotation matrix

$$(10) \quad Q(t) := \begin{bmatrix} \cos \theta(t) & -\sin \theta(t) \\ \sin \theta(t) & \cos \theta(t) \end{bmatrix},$$

such that the position $\eta(t, x) \in \mathcal{S}(t)$ at the time t of the point fixed to the body with an initial position x is

$$(11) \quad \eta(t, x) := h(t) + Q(t)(x - h_0).$$

The angle θ satisfies

$$\theta'(t) = r(t),$$

and we choose $\theta(t)$ such that $\theta(0) = 0$.

The Equations (1) and (2) are the incompressible Euler equations, the condition (3) means that the boundary is impermeable and the Equations (5) and (6) are the Newton's balance law for linear and angular momenta.

For the study of ideal flow, an important quantity is the vorticity $w := \text{curl } u = \partial_1 u_2 - \partial_2 u_1$, satisfying the transport equation:

$$(12) \quad \frac{\partial w}{\partial t} + (u \cdot \nabla)w = 0 \text{ for } x \in \mathcal{F}(t).$$

One has the following result concerning the Cauchy problem for the above system, the initial position of the solid being given. This result describes weak solutions, extending results concerning the fluid alone. It considers the case when vorticity belongs in L^p as in DiPerna and Majda [7] and includes weak solutions with bounded vorticity as in Yudovich [13].

THEOREM 1. — *Let $p \in (2, +\infty]$. For any $u_0 \in C^0(\overline{\mathcal{F}_0}; \mathbb{R}^2)$, $(\ell_0, r_0) \in \mathbb{R}^2 \times \mathbb{R}$, such that:*

$$(13) \quad \text{div } u_0 = 0 \text{ in } \mathcal{F}_0 \text{ and } u_0 \cdot n = (\ell_0 + r_0(x - h_0)^\perp) \cdot n \text{ on } \partial \mathcal{S}_0,$$

$$(14) \quad w_0 := \text{curl } u_0 \in L^p_c(\overline{\mathcal{F}_0}),$$

$$\lim_{|x| \rightarrow +\infty} u_0(x) = 0,$$

there exists a solution (h', r, u) of (1)–(8) in $C^1(\mathbb{R}^+; \mathbb{R}^2 \times \mathbb{R}) \times C^0(\mathbb{R}^+, W^{1,p}(\mathcal{F}(t)))$ with $\partial_t u, \nabla p \in L^\infty_{loc}(\mathbb{R}^+, L^q(\mathcal{F}(t)))$ for any $q \in (1, p]$ when $p < \infty$ and in $C^1(\mathbb{R}^+; \mathbb{R}^2 \times \mathbb{R}) \times L^\infty_{loc}(\mathbb{R}^+, \mathcal{LL}(\mathcal{F}(t)))$ with $\partial_t u, \nabla p \in L^\infty_{loc}(\mathbb{R}^+, L^q(\mathcal{F}(t)))$ for any $q \in (1, +\infty)$ when $p = \infty$.

Moreover such a solution satisfies that for all $t > 0$, $w(t) := \text{curl}(u(t)) \in L^p_c(\overline{\mathcal{F}(t)})$, it is energy-conserving in the sense of Proposition 4 and $\|w(t, \cdot)\|_{L^q(\mathcal{F}(t))}$ (for any $q \in [1, p]$), $\int_{\mathcal{F}(t)} w(t, x) dx$ and $\int_{\partial \mathcal{S}(t)} u \cdot \tau ds$ are preserved over time.

Finally when $p = \infty$, the solution is unique.

This result is proven in [10]. For the sake of self-containedness, we give a short proof of it in appendix. The notation $\mathcal{LL}(\Omega)$ refers to the space of log-Lipschitz functions on Ω , that is the set of functions $f \in L^\infty(\Omega)$ such that

$$(15) \quad \|f\|_{\mathcal{LL}(\Omega)} := \|f\|_{L^\infty(\Omega)} + \sup_{x \neq y} \frac{|f(x) - f(y)|}{|(x - y)(1 + \ln^- |x - y|)} < +\infty.$$