

*quatrième série - tome 48      fascicule 1      janvier-février 2015*

*ANNALES  
SCIENTIFIQUES  
de  
L'ÉCOLE  
NORMALE  
SUPÉRIEURE*

Curtis T. McMULLEN

*Cascades in the dynamics of measured foliations*

---

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

# Annales Scientifiques de l'École Normale Supérieure

---

Publiées avec le concours du Centre National de la Recherche Scientifique

## Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

### Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE  
de 1883 à 1888 par H. DEBRAY  
de 1889 à 1900 par C. HERMITE  
de 1901 à 1917 par G. DARBOUX  
de 1918 à 1941 par É. PICARD  
de 1942 à 1967 par P. MONTEL

### Comité de rédaction au 1<sup>er</sup> janvier 2015

N. ANANTHARAMAN B. KLEINER  
E. BREUILLARD E. KOWALSKI  
R. CERF P. LE CALVEZ  
A. CHAMBERT-LOIR M. MUSTAȚĂ  
I. GALLAGHER L. SALOFF-COSTE

## Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,  
45, rue d'Ulm, 75230 Paris Cedex 05, France.  
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.  
[annaales@ens.fr](mailto:annaales@ens.fr)

---

### Édition / *Publication*

Société Mathématique de France  
Institut Henri Poincaré  
11, rue Pierre et Marie Curie  
75231 Paris Cedex 05  
Tél. : (33) 01 44 27 67 99  
Fax : (33) 01 40 46 90 96

### Abonnements / *Subscriptions*

Maison de la SMF  
Case 916 - Luminy  
13288 Marseille Cedex 09  
Fax : (33) 04 91 41 17 51  
email : [smf@smf.univ-mrs.fr](mailto:smf@smf.univ-mrs.fr)

### Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

---

© 2015 Société Mathématique de France, Paris

En application de la loi du 1<sup>er</sup> juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

*All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.*

---

ISSN 0012-9593

Directeur de la publication : Marc Peigné  
Périodicité : 6 n<sup>os</sup> / an

# CASCADES IN THE DYNAMICS OF MEASURED FOLIATIONS

BY CURTIS T. McMULLEN

---

**ABSTRACT.** – This paper studies the behavior of harmonic measured foliations on compact Riemann surfaces. Cascades in the dynamics of such a foliation can occur as its relative periods are varied. We show that in the case of genus 2, the bifurcation locus arising from such a variation is a closed, countable set of  $\mathbb{R}$  that embeds in  $\omega^\omega$ .

**RÉSUMÉ.** – Nous étudions le comportement des feuilletages mesurés harmoniques sur les surfaces de Riemann compactes. Quand les périodes relatives varient, on peut observer des cascades dans la dynamique d'un tel feuilletage. Dans le cas du genre 2, on montre que le lieu de bifurcation résultant d'une telle variation est un sous-ensemble dénombrable et fermé de  $\mathbb{R}$ , qui se plonge dans  $\omega^\omega$ .

## 1. Introduction

Let  $X$  be a compact Riemann surface of genus  $g$ . Every harmonic 1-form  $\rho$  on  $X$  is the pullback of a linear form under a suitable *period map*

$$\pi : X \rightarrow E \cong \mathbb{R}^r / \mathbb{Z}^r.$$

The leaves of the associated measured foliation  $\mathcal{F}(\rho)$  of  $X$  map under  $\pi$  into the leaves of an irrational foliation of  $E$ .

When  $r = 2$ , the behavior of  $\mathcal{F}(\rho)$  is strongly influenced by the *degree* of the period map. For example, if  $\mathcal{F}(\rho)$  is periodic then its degree must be zero. At the other extreme, in § 6 we will see:

**THEOREM 1.1.** – *There is no minimal foliation of degree zero.*

The degree depends only on the *absolute periods* of  $\rho$ , given by the real cohomology class  $[\rho] \in H^1(X)$ . In the case of genus  $g = 2$ , there is one important remaining invariant, the *relative period* of  $\rho$  along a path connecting its zeros (see § 7). In § 8 we will show:

---

Research supported in part by the NSF and MSRI. Revised in September 2013.

**THEOREM 1.2.** – *Let  $(X_t, \rho_t)$  be a family of harmonic forms of genus two with constant absolute periods and relative period  $t$ . Assume  $(X_0, \rho_0)$  has degree zero. Then the bifurcation locus*

$$B = \{t : \mathcal{F}(\rho_t) \text{ is not periodic}\}$$

*is a closed, countable subset of  $\mathbb{R}$  which embeds in  $\omega^\omega$ .*




FIGURE 1. Cascades in the dynamics of  $F_t : [0, 1] \rightarrow [0, 1]$ .

### Example: 1-dimensional dynamics

The spikes in Figure 1 give the bifurcation locus for a family  $(X_t, \rho_t)$  that depends on an irrational number  $L \in (0, 1)$ . Here a transversal to  $\mathcal{F}(\rho_t)$  determines an interval exchange map

$$F_t : [0, 1] \rightarrow [0, 1].$$

The map  $F_t$  rotates  $[0, 1]$  by  $t$ , then rotates the subintervals  $[0, L]$  and  $[L, 1]$  each by  $-t$ . (Rotation of  $[0, a]$  by  $b$  is given by  $x \mapsto (x + b) \bmod a$ .)

The period  $N(t)$  of  $F_t(x)$  is finite iff  $t \notin B$ . The figure shows the graph of  $y = \log_2 N(t)$  (calculated using Rauzy induction) in the case  $L = \pi/7$ , whose behavior is typical. Theorem 1.2 shows that structural stability is dense in this and similar families of interval exchange transformations.

### Homological invariants

To place these results in a broader setting, consider a general closed 1-form  $\rho$  on a compact, oriented  $n$ -manifold  $X$ . The Poincaré dual of  $[\rho] \in H^1(X)$  maps to a natural cycle

$$\text{flux}(X, \rho) \in H_{n-1}(E),$$

which records the average distribution of the leaves of  $\mathcal{F}(\rho)$  under the period map  $\pi : X \rightarrow E$ .

The vanishing of flux generalizes the vanishing of degree. As we will see in §4, zero flux encourages the leaves of  $\mathcal{F}(\rho)$  to double back in a manner that is delicate to combine with minimality. Minimal foliations with zero flux in genus three will be discussed in §5. Such examples cannot exist in genus two, as we will see in §6 and §7.

A measurable set  $A \subset X$  is *saturated* if it is a union of leaves of  $\mathcal{F}(\rho)$ . In this case  $d(\rho|_A) = 0$  as a current. We define the *content* of  $\rho$  to be the smallest convex set

$$C(\rho) \subset H^1(X)$$

containing  $[\rho|A]$  for every saturated set. In §2 we show this dynamical invariant is upper semicontinuous, in the sense that

$$\limsup C(\rho_n) \subset C(\rho)$$

whenever  $\rho_n \rightarrow \rho$ .

Similarly, when  $(X, \rho)$  has rank  $r = n$ , we can make sense of the degree (as a real number) for any saturated set, and we define

$$\deg^+(X, \rho) = \sup_A \deg(\pi|A).$$

In the context of Theorem 1.2, in §8 we will show:

**THEOREM 1.3.** – *The quantity  $d(t) = \deg^+(X_t, \rho_t) \geq 0$  is an integer,  $B$  is the locus where  $d(t) > 0$ , and we have*

$$(1.1) \quad \limsup_{s \rightarrow t} d(s) \leq d(t) - 1$$

whenever  $t \in B$ .

We emphasize that  $d(s)$  is bounded by  $d(t) - 1$  (and not just  $d(t)$ ) for all  $s$  near  $t$ . This evaporation of degree under perturbations is one of the main new phenomena we establish in this paper. The properties of  $B$  stated in Theorem 1.2 follow from it immediately, and it plays a key role in the analysis of isoperiodic forms [34].

### Quadratic fields

In §8 we will also show:

**THEOREM 1.4.** – *If the absolute periods of  $\rho_0$  lie in a real quadratic field, then  $B$  is self-similar about every point.*

This means  $B$  is locally invariant under a linear contraction fixing  $a$ , for every  $a \in B$ .

A similar renormalization mechanism causes one bifurcation set to reappear in another. Examples of four such bifurcation sets, with the periods of  $\rho_0$  in  $\mathbb{Q}(\sqrt{5})$ , are shown in Figure 2. Note that a compressed version of each cascade reappears in the next, near  $t = (3 + \sqrt{5})/2 = 2.618\dots$  Using this idea we give explicit examples where  $\omega^k$  embeds in  $B$  (§10). For an example with  $B \cong \omega^\omega$ , see [34, §11].

### Appendix: Teichmüller curves

Much attention has focused on the family of foliations arising from parallel geodesics with varying slopes on the flat surface  $(X, |\omega|)$ . These foliations are given by  $\rho_t = \operatorname{Re}(e^{it}\omega)$ .

The invariants discussed in §2 can also be studied for such families. In the Appendix we show that the (rapidly fluctuating) convex set  $C(\rho_t) \subset H^1(X)$  can be readily determined whenever  $(X, \omega)$  generates a Teichmüller curve. An example associated to billiards in the regular pentagon is shown in Figure 3. This convex set records the cylinder decomposition at each cusp of  $\operatorname{SL}(X, \omega)$ , and can be computed by a continued fraction algorithm in  $\mathbb{Q}(\sqrt{5})$ . For another perspective, we also show:

*The function graphed in Figure 3 arises from the radial limits  $F : S^1 \rightarrow \mathbb{PML}_g$  of a suitable Teichmüller disk  $f : \Delta \rightarrow \mathcal{T}_g$ .*