

Bulletin

de la SOCIÉTÉ MATHÉMATIQUE DE FRANCE

MEROMORPHIC QUOTIENTS FOR SOME HOLOMORPHIC G-ACTIONS

Daniel Barlet

**Tome 146
Fascicule 3**

2018

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Publié avec le concours du Centre national de la recherche scientifique

pages 441-477

Le *Bulletin de la Société Mathématique de France* est un périodique
trimestriel de la Société Mathématique de France.

Fascicule 3, tome 146, septembre 2018

Comité de rédaction

Christine BACHOC
Yann BUGEAUD
Jean-François DAT
Pascal HUBERT
Laurent MANIVEL

Julien MARCHÉ
Kieran O'GRADY
Emmanuel RUSS
Christophe SABOT

Marc HERZLICH (Dir.)

Diffusion

Maison de la SMF	AMS
Case 916 - Luminy	P.O. Box 6248
13288 Marseille Cedex 9	Providence RI 02940
France	USA
commandes@smf.emath.fr	http://www.ams.org

Tarifs

Vente au numéro : 43 € (\$ 64)

Abonnement électronique : 135 € (\$ 202),

avec supplément papier : Europe 179 €, hors Europe 197 € (\$ 296)

Des conditions spéciales sont accordées aux membres de la SMF.

Secrétariat

Bulletin de la Société Mathématique de France
Société Mathématique de France
Institut Henri Poincaré, 11, rue Pierre et Marie Curie
75231 Paris Cedex 05, France
Tél. : (33) 01 44 27 67 99
bulletin@smf.emath.fr • <http://smf.emath.fr>

© *Société Mathématique de France* 2018

Tous droits réservés (article L 122-4 du Code de la propriété intellectuelle). Toute représentation ou reproduction intégrale ou partielle faite sans le consentement de l'éditeur est illicite. Cette représentation ou reproduction par quelque procédé que ce soit constituerait une contrefaçon sanctionnée par les articles L 335-2 et suivants du CPI.

ISSN 0037-9484 (print) 2102-622X (electronic)

Directeur de la publication : Stéphane SEURET

MEROMORPHIC QUOTIENTS FOR SOME HOLOMORPHIC G-ACTIONS

BY DANIEL BARLET

ABSTRACT. — Using mainly tools from previous articles we give necessary and sufficient conditions on the G -orbits' configuration in X in order that a holomorphic action of a connected complex Lie group G on a reduced complex space X admits a *strongly quasi-proper meromorphic quotient*. To show how these conditions can be used, we show, when $G = K.B$ with B a closed connected complex subgroup of G and K a real compact subgroup of G , the existence of a strongly quasi-proper meromorphic quotient for the G -action on X , assuming a slightly stronger condition than the existence of such a quotient for the B -action. We also give a similar result when the connected complex Lie group has the form $G = K.A.K$ where A is a closed connected complex subgroup and K is a compact (real) subgroup.

RÉSUMÉ (*Quotients méromorphes pour certaines G -actions holomorphes*). — En utilisant les résultats de précédents articles, nous donnons des conditions nécessaires et suffisantes sur la configuration des G -orbites dans X pour que l'action holomorphe d'un groupe de Lie complexe connexe sur un espace complexe réduit X admette un *quotient méromorphe fortement quasi-propre*. Pour illustrer l'intérêt de ces conditions, nous montrons, quand $G = K.B$ où B est un sous-groupe connexe complexe fermé et

Texte reçu le 9 mai 2016, accepté le 6 mars 2017.

DANIEL BARLET, Institut Elie Cartan, Géométrie, Université de Lorraine, CNRS UMR 7502 and Institut Universitaire de France.

Mathematical subject classification (2010). — 32M05, 32H04, 32H99, 57S20.

Key words and phrases. — Holomorphic G -action, finite type cycles, strongly quasi-proper map, holomorphic quasi-proper geometrically flat quotient, strongly quasi-proper meromorphic quotient.

By many discussions and interesting suggestions Peter Heinzner helps me during the working out of this paper. I want to thank him for that and also for the nice hospitality of the Mathematic Faculty of Bochum University. This work was partially supported by the CRC/TRR 191 Symplectic Structures in Geometry, Algebra and Dynamics of the Deutsche Forschungsgemeinschaft (DFG).

K un sous-groupe compact réel de G , l'existence d'un quotient méromorphe fortement quasi-propre pour l'action de G sur X sous une hypothèse légèrement plus forte que l'existence d'un tel quotient pour l'action de B sur X . Nous donnons également un résultat analogue quand $G = K.A.K$ où A est un sous-groupe complexe fermé et connexe et K un sous-groupe compact réel de G .

1. Introduction

In this article we explain how the tools developed in [9], [1], [2] and [3] can be applied to produce, in suitable cases, a meromorphic quotient of a holomorphic action of a connected complex Lie group G on a reduced complex space X . This uses the notion of *strongly quasi-proper map* introduced in *loc. cit.* and our first goal is to give three hypotheses, called [H.1], [H.2], [H.3], on the G -orbits' configuration in X which are *equivalent* to the existence of a *strongly quasi-proper meromorphic quotient*, notion defined in the Section 1.2.

The proof of this equivalence is the content of Proposition 2.7.1 and Theorem 2.8.1. Then we give a sufficient condition [H.1str], asking the existence of a G -invariant set $\Omega_1 \subset X$ which is dense, Zariski open and “good” for the action, to satisfy the condition [H.1].

Note that the conditions [H.1], [H.2], [H.3] introduced in Section 2.7 only depend on the G -orbits' configuration in X , but the condition [H.1str] depends on the action of G on X itself.

The existence theorem for a strongly quasi-proper meromorphic quotient under our three assumptions is applied to prove the following result:

THEOREM 1.0.1. — *Assume that we have a holomorphic action of a connected complex Lie group G on a reduced complex space X . Assume that $G = K.B$ where K is a compact (real) subgroup of G and B a connected complex closed subgroup of G . Assume that the action of B on X satisfies the condition [H.1str] on a G -invariant Zariski open dense subset Ω in $X^{(1)}$, and the conditions [H.2] and [H.3]. Then the G -action satisfies [H.1str], [H.2] and [H.3]; so it has a strongly quasi-proper meromorphic quotient.*

A first variant of this result is given by the following theorem.

THEOREM 1.0.2. — *Assume that we have a holomorphic action of a connected complex Lie group G on a reduced complex space X . Assume that $G = K.B$ where K is a compact (real) subgroup of G and B a connected complex closed subgroup of G . Assume that K normalizes B and that the B -action satisfies*

1. This precisely means that there exists a G -invariant dense Zariski open set in X which is a “good open set” for the B -action (see Section 2.5)

the conditions [H.1str], [H.2] and [H.3]. Then the G -action satisfies the conditions [H.1str], [H.2] and [H.3] and so has a strongly quasi-proper meromorphic quotient.

Here is a second result obtained by a similar method.

THEOREM 1.0.3. — *Let G be a complex connected Lie group and assume that there exists a closed connected complex subgroup A and a compact (real) subgroup K such that $G = K.A.K$. Assume that we have a completely holomorphic action of G on an irreducible complex space X and that the action of A on X satisfies the following properties:*

- i) *The hypothesis [H.1str] for the A -action is satisfied on a G -invariant (Zariski good) open set Ω_1 in X .*
- ii) *The hypothesis [H.2] for the A -action is satisfied on a G -invariant open set $\Omega_0 \subset \Omega_1$ in X .*
- iii) *The hypothesis [H.3] holds for the A -action.*

Then [H.1str], [H.2] and [H.3] hold for the action of G on X . So there exists a SQP meromorphic quotient of X for the G -action.

Of course the hypothesis $G = K.A.K$ is more “general” than the case $G = K.B$. But the hypothesis of this last theorem is more restrictive for the action on X of the closed connected complex subgroup A of G : we ask also the G -invariance of the dense open subset Ω_0 of Ω_1 (the open set Ω_0 is defined in the condition [H.2]).

We conclude this article with two results (see Section 3.4) relating the SQP meromorphic quotients for the actions of B and G (resp. of A and G) when they exist:

1. The existence of a holomorphic map $h : Q_B \rightarrow Q_G$ (resp. $Q_A \rightarrow Q_G$) between the corresponding quotients.
2. The existence under the hypotheses of the Theorem 1.0.1 (resp. the Theorem 1.0.3) of a G -invariant dense Zariski open set Ω disjoint from the centers of the modifications, such that the corresponding map $h_\Omega : q_B(\Omega) \rightarrow q_G(\Omega)$ (resp. $h_\Omega : q_A(\Omega) \rightarrow q_G(\Omega)$) is proper.

2. Strongly quasi-proper meromorphic quotients

2.1. Preliminaries. — For the definition of the topology on the space $\mathcal{C}_n^f(X)$ of finite type n -cycles in X and its relationship with the topology of the space $\mathcal{C}_n^{\text{loc}}(X)$ we refer to [4] ch. IV, [2] and [3].

For the convenience of the reader we recall shortly here the definitions of a geometrically f-flat map (f-GF map) and of a strongly quasi-proper map (SQP map) between irreducible complex spaces and we give a short summary on some properties of the SQP maps. For more details on these notions see [2] and [3].